



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 05:55 PM UTC

PDB ID : 5MU2 / pdb_00005mu2
Title : ACC1 Fab fragment in complex with CII583-591 (CG10)
Authors : Dobritsch, D.; Holmdahl, R.; Ge, C.
Deposited on : 2017-01-12
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	218	
1	C	218	
1	E	218	
1	G	218	
1	I	218	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	K	218	
1	M	218	
1	O	218	
2	B	218	
2	D	218	
2	F	218	
2	H	218	
2	J	218	
2	L	218	
2	N	218	
2	P	218	
3	Q	30	
3	R	30	
3	S	30	
3	T	30	
3	U	30	
3	V	30	
3	W	30	
3	X	30	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 27447 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ACC1 Fab fragment heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	216	Total	C	N	O	S	0	0	0
			1629	1028	271	322	8			
1	C	218	Total	C	N	O	S	0	0	0
			1640	1034	273	325	8			
1	E	215	Total	C	N	O	S	0	0	0
			1622	1024	270	320	8			
1	G	211	Total	C	N	O	S	0	0	0
			1601	1011	266	317	7			
1	I	218	Total	C	N	O	S	0	0	0
			1640	1034	273	325	8			
1	K	216	Total	C	N	O	S	0	0	0
			1629	1028	271	322	8			
1	M	216	Total	C	N	O	S	0	0	0
			1629	1028	271	322	8			
1	O	214	Total	C	N	O	S	0	0	0
			1615	1019	269	319	8			

- Molecule 2 is a protein called ACC1 Fab fragment light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	218	Total	C	N	O	S	0	0	0
			1666	1038	282	339	7			
2	D	218	Total	C	N	O	S	0	0	0
			1666	1038	282	339	7			
2	F	217	Total	C	N	O	S	0	0	0
			1660	1035	281	338	6			
2	H	217	Total	C	N	O	S	0	0	0
			1660	1035	281	338	6			
2	J	218	Total	C	N	O	S	0	0	0
			1666	1038	282	339	7			
2	L	218	Total	C	N	O	S	0	0	0
			1666	1038	282	339	7			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	N	218	Total	C	N	O	S	0	0	0
			1666	1038	282	339	7			
2	P	217	Total	C	N	O	S	0	0	0
			1660	1035	281	338	6			

- Molecule 3 is a protein called synthetic peptide containing the CII583-591 epitope of collagen type II.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	X	15	Total	C	N	O	0	0	0
			95	59	18	18			
3	Q	15	Total	C	N	O	0	0	0
			95	59	18	18			
3	R	15	Total	C	N	O	0	0	0
			95	59	18	18			
3	S	15	Total	C	N	O	0	0	0
			95	59	18	18			
3	T	13	Total	C	N	O	0	0	0
			83	52	16	15			
3	U	15	Total	C	N	O	0	0	0
			95	59	18	18			
3	V	12	Total	C	N	O	0	0	0
			76	47	15	14			
3	W	9	Total	C	N	O	0	0	0
			57	35	12	10			

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	37	Total	O	0	0
			37	37		
4	B	38	Total	O	0	0
			38	38		
4	C	18	Total	O	0	0
			18	18		
4	D	64	Total	O	0	0
			64	64		
4	E	42	Total	O	0	0
			42	42		
4	F	24	Total	O	0	0
			24	24		
4	G	21	Total	O	0	0
			21	21		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	H	12	Total 12	O 12	0	0
4	I	37	Total 37	O 37	0	0
4	J	25	Total 25	O 25	0	0
4	K	17	Total 17	O 17	0	0
4	L	28	Total 28	O 28	0	0
4	M	15	Total 15	O 15	0	0
4	N	19	Total 19	O 19	0	0
4	O	10	Total 10	O 10	0	0
4	P	22	Total 22	O 22	0	0
4	X	1	Total 1	O 1	0	0
4	Q	3	Total 3	O 3	0	0
4	R	2	Total 2	O 2	0	0
4	S	2	Total 2	O 2	0	0
4	T	3	Total 3	O 3	0	0
4	U	1	Total 1	O 1	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ACC1 Fab fragment heavy chain




- Molecule 1: ACC1 Fab fragment heavy chain



- Molecule 1: ACC1 Fab fragment heavy chain





- Molecule 1: ACC1 Fab fragment heavy chain




- Molecule 1: ACC1 Fab fragment heavy chain





- Molecule 1: ACC1 Fab fragment heavy chain



- Molecule 1: ACC1 Fab fragment heavy chain



- Molecule 1: ACC1 Fab fragment heavy chain



- Molecule 2: ACC1 Fab fragment light chain




- Molecule 2: ACC1 Fab fragment light chain



- Molecule 2: ACC1 Fab fragment light chain




- Molecule 2: ACC1 Fab fragment light chain




- Molecule 2: ACC1 Fab fragment light chain

Chain J:  95% 5%





- Molecule 2: ACC1 Fab fragment light chain

Chain L:  94% 5%




- Molecule 2: ACC1 Fab fragment light chain

Chain N:  94% 5%




- Molecule 2: ACC1 Fab fragment light chain

Chain P:  93% 6% %




- Molecule 3: synthetic peptide containing the CII583-591 epitope of collagen type II

Chain X:  3% 43% 7% 50%





- Molecule 3: synthetic peptide containing the CII583-591 epitope of collagen type II

Chain Q:  7% 40% 10% 50%





- Molecule 3: synthetic peptide containing the CII583-591 epitope of collagen type II


Chain R:  7% 40% 10% 50%





- Molecule 3: synthetic peptide containing the CII583-591 epitope of collagen type II




- Molecule 3: synthetic peptide containing the CII583-591 epitope of collagen type II




- Molecule 3: synthetic peptide containing the CII583-591 epitope of collagen type II



- Molecule 3: synthetic peptide containing the CII583-591 epitope of collagen type II



- Molecule 3: synthetic peptide containing the CII583-591 epitope of collagen type II



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	148.86Å 156.01Å 156.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.09 – 2.70 49.09 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.9 (49.09-2.70) 99.9 (49.09-2.70)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.20 (at 2.69Å)	Xtriage
Refinement program	REFMAC 5.8.0155	Depositor
R, R_{free}	0.206 , 0.228 (Not available) , 0.205	Depositor DCC
R_{free} test set	4865 reflections (4.84%)	wwPDB-VP
Wilson B-factor (Å ²)	46.1	Xtriage
Anisotropy	0.194	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 29.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.018 for -h,l,k 0.018 for -l,-k,-h 0.019 for k,h,-l 0.012 for k,l,h 0.012 for l,h,k	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	27447	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HYP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.65	0/1669	0.83	2/2277 (0.1%)
1	C	0.67	0/1681	0.86	3/2295 (0.1%)
1	E	0.70	0/1662	0.84	3/2267 (0.1%)
1	G	0.64	0/1640	0.80	2/2237 (0.1%)
1	I	0.70	0/1681	0.82	2/2295 (0.1%)
1	K	0.67	0/1669	0.78	0/2277
1	M	0.63	0/1669	0.83	3/2277 (0.1%)
1	O	0.64	0/1654	0.86	4/2255 (0.2%)
2	B	0.65	0/1703	0.77	0/2310
2	D	0.72	0/1703	0.82	0/2310
2	F	0.66	0/1697	0.79	0/2302
2	H	0.63	0/1697	0.78	0/2302
2	J	0.66	0/1703	0.78	0/2310
2	L	0.64	0/1703	0.77	0/2310
2	N	0.63	0/1703	0.77	0/2310
2	P	0.70	0/1697	0.83	2/2302 (0.1%)
3	Q	0.83	0/79	0.85	0/102
3	R	0.89	0/80	0.92	0/106
3	S	0.93	0/80	1.07	0/106
3	T	0.81	0/76	0.94	0/100
3	U	0.95	0/80	0.86	0/106
3	V	0.78	0/69	0.96	0/92
3	W	0.87	0/57	1.20	0/75
3	X	0.88	0/80	0.91	0/106
All	All	0.67	0/27532	0.81	21/37429 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1
1	I	0	1
1	K	0	1
2	B	0	1
2	J	0	1
2	L	0	1
2	N	0	1
All	All	0	7

There are no bond length outliers.

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	O	69	ARG	NE-CZ-NH2	-11.07	109.23	119.20
1	M	69	ARG	NE-CZ-NH2	-10.68	109.59	119.20
1	C	69	ARG	NE-CZ-NH2	-10.63	109.63	119.20
2	P	192	ARG	NE-CZ-NH2	10.13	128.32	119.20
1	C	69	ARG	NE-CZ-NH1	9.84	131.34	121.50
1	O	69	ARG	NE-CZ-NH1	9.79	131.29	121.50
1	M	69	ARG	NE-CZ-NH1	9.59	131.09	121.50
1	E	69	ARG	NE-CZ-NH2	8.67	127.01	119.20
1	I	69	ARG	NE-CZ-NH2	8.54	126.88	119.20
1	G	69	ARG	NE-CZ-NH2	8.43	126.79	119.20
1	A	69	ARG	NE-CZ-NH2	8.02	126.42	119.20
1	G	69	ARG	NE-CZ-NH1	-7.29	114.21	121.50
1	E	69	ARG	NE-CZ-NH1	-7.16	114.34	121.50
1	I	69	ARG	NE-CZ-NH1	-7.13	114.37	121.50
1	A	69	ARG	NE-CZ-NH1	-7.05	114.45	121.50
1	E	119	THR	CA-CB-CG2	6.84	122.13	110.50
1	M	69	ARG	CD-NE-CZ	6.76	133.87	124.40
1	C	69	ARG	CD-NE-CZ	6.64	133.69	124.40
1	O	69	ARG	CD-NE-CZ	6.62	133.67	124.40
2	P	192	ARG	NE-CZ-NH1	-6.43	115.07	121.50
1	O	131	CYS	CA-C-O	-6.14	114.37	120.82

There are no chirality outliers.

All (7) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	217	GLU	Peptide
1	C	135	THR	Peptide
1	I	135	THR	Peptide

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
2	J	216	ASN	Peptide
1	K	52	ARG	Sidechain
2	L	216	ASN	Peptide
2	N	216	ASN	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1629	0	1589	2	0
1	C	1640	0	1600	10	0
1	E	1622	0	1583	1	0
1	G	1601	0	1563	6	0
1	I	1640	0	1600	7	0
1	K	1629	0	1589	1	1
1	M	1629	0	1589	2	0
1	O	1615	0	1576	2	1
2	B	1666	0	1597	7	0
2	D	1666	0	1597	12	0
2	F	1660	0	1593	5	0
2	H	1660	0	1593	5	0
2	J	1666	0	1597	5	0
2	L	1666	0	1597	7	0
2	N	1666	0	1597	10	0
2	P	1660	0	1593	6	0
3	Q	95	0	95	1	0
3	R	95	0	96	1	0
3	S	95	0	95	1	0
3	T	83	0	85	0	0
3	U	95	0	95	0	0
3	V	76	0	79	0	0
3	W	57	0	60	1	0
3	X	95	0	96	0	0
4	A	37	0	0	0	0
4	B	38	0	0	0	0
4	C	18	0	0	1	0
4	D	64	0	0	0	0
4	E	42	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	F	24	0	0	0	0
4	G	21	0	0	0	0
4	H	12	0	0	0	0
4	I	37	0	0	0	0
4	J	25	0	0	0	0
4	K	17	0	0	0	0
4	L	28	0	0	0	0
4	M	15	0	0	0	0
4	N	19	0	0	0	0
4	O	10	0	0	0	0
4	P	22	0	0	0	0
4	Q	3	0	0	0	0
4	R	2	0	0	0	0
4	S	2	0	0	0	0
4	T	3	0	0	0	0
4	U	1	0	0	0	0
4	X	1	0	0	0	0
All	All	27447	0	26154	63	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:M:174:LEU:HD13	2:N:164:LEU:HD11	1.61	0.79
2:F:32:TYR:HA	1:I:161:SER:O	1.90	0.72
1:O:174:LEU:HD13	2:P:164:LEU:HD11	1.74	0.68
1:A:174:LEU:HD13	2:B:164:LEU:HD11	1.75	0.67
2:J:204:THR:O	2:L:33:GLY:N	2.30	0.65
1:C:174:LEU:HB2	2:D:164:LEU:HD11	1.79	0.65
2:F:30:ASP:HB3	1:I:164:SER:OG	1.96	0.64
1:I:61:ASN:HD21	2:P:7:SER:HB2	1.64	0.63
1:C:75:ASP:OD1	2:N:203:LYS:HE3	2.00	0.62
2:D:73:THR:OG1	1:I:193:SER:OG	2.17	0.61
2:P:155:ASP:OD1	2:P:194:ASN:OD1	2.19	0.61
1:G:154:THR:HG22	1:G:201:ALA:HB3	1.83	0.59
2:J:164:LEU:C	2:J:164:LEU:HD23	2.29	0.58
2:D:192:ARG:O	2:D:192:ARG:HG3	2.04	0.56
2:D:73:THR:OG1	1:I:193:SER:CB	2.54	0.55
2:D:204:THR:O	1:G:1:GLU:N	2.37	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:P:140:LEU:HD13	2:P:179:MET:HE2	1.91	0.53
2:D:140:LEU:HD13	2:D:179:MET:HE2	1.91	0.53
2:H:140:LEU:HD13	2:H:179:MET:HE2	1.90	0.52
1:C:174:LEU:HD13	2:D:164:LEU:HD11	1.91	0.52
2:N:140:LEU:HD13	2:N:179:MET:HE2	1.91	0.52
2:L:140:LEU:HD13	2:L:179:MET:HE2	1.91	0.52
1:E:31:ASP:O	3:R:14:GLY:HA3	2.10	0.51
2:D:73:THR:OG1	1:I:193:SER:HB3	2.10	0.51
1:C:78:ARG:NH1	2:N:203:LYS:O	2.44	0.51
2:J:140:LEU:HD13	2:J:179:MET:HE2	1.93	0.51
2:B:140:LEU:HD13	2:B:179:MET:HE2	1.94	0.50
1:C:124:VAL:HG21	1:C:209:VAL:HG11	1.93	0.50
1:G:154:THR:CG2	1:G:201:ALA:HB3	2.43	0.49
1:C:134:THR:HG21	4:C:312:HOH:O	2.13	0.48
1:C:57:ASN:HB3	2:N:11:LEU:HD23	1.95	0.48
1:C:174:LEU:HD22	2:D:164:LEU:CD1	2.44	0.48
2:F:140:LEU:HD13	2:F:179:MET:HE2	1.96	0.48
2:J:204:THR:O	2:L:33:GLY:CA	2.63	0.47
1:A:131:CYS:HB3	2:B:218:CYS:HB3	1.73	0.46
2:H:140:LEU:HD13	2:H:179:MET:CE	2.46	0.46
2:B:140:LEU:HD13	2:B:179:MET:CE	2.47	0.45
1:M:106:TRP:CE3	2:N:48:PRO:HD2	2.51	0.45
2:P:140:LEU:HD13	2:P:179:MET:CE	2.47	0.44
1:G:126:PRO:O	2:H:125:SER:HB3	2.18	0.44
2:L:140:LEU:HD13	2:L:179:MET:CE	2.46	0.44
1:O:31:ASP:O	3:W:14:GLY:HA3	2.17	0.44
1:C:57:ASN:CB	2:N:11:LEU:HD23	2.48	0.43
2:N:140:LEU:HD13	2:N:179:MET:CE	2.47	0.43
2:D:140:LEU:HD13	2:D:179:MET:CE	2.48	0.43
2:F:140:LEU:HD13	2:F:179:MET:CE	2.49	0.42
2:L:11:LEU:HD22	2:L:13:VAL:CG2	2.49	0.42
2:B:140:LEU:HD12	2:B:140:LEU:N	2.35	0.42
2:F:11:LEU:HD22	2:F:13:VAL:CG2	2.50	0.41
1:G:31:ASP:O	3:S:14:GLY:HA3	2.20	0.41
2:J:140:LEU:HD13	2:J:179:MET:CE	2.49	0.41
2:D:140:LEU:N	2:D:140:LEU:HD12	2.36	0.41
1:K:174:LEU:HD13	2:L:164:LEU:HD11	2.03	0.41
1:C:31:ASP:O	3:Q:14:GLY:HA3	2.21	0.41
2:L:140:LEU:N	2:L:140:LEU:HD12	2.35	0.41
2:N:11:LEU:HD22	2:N:13:VAL:CG2	2.50	0.41
2:H:140:LEU:N	2:H:140:LEU:HD12	2.35	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:N:140:LEU:HD12	2:N:140:LEU:N	2.36	0.41
2:B:82:VAL:CG1	2:B:110:ILE:HD12	2.51	0.41
2:D:11:LEU:HD22	2:D:13:VAL:CG2	2.51	0.41
1:G:174:LEU:HD13	2:H:164:LEU:HD11	2.02	0.40
1:I:61:ASN:ND2	2:P:7:SER:HB2	2.35	0.40
2:B:11:LEU:HD22	2:B:13:VAL:CG2	2.52	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:K:28:THR:OG1	1:O:164:SER:OG[3_555]	2.01	0.19

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	212/218 (97%)	206 (97%)	6 (3%)	0	100	100
1	C	216/218 (99%)	209 (97%)	7 (3%)	0	100	100
1	E	211/218 (97%)	205 (97%)	6 (3%)	0	100	100
1	G	207/218 (95%)	202 (98%)	5 (2%)	0	100	100
1	I	216/218 (99%)	210 (97%)	6 (3%)	0	100	100
1	K	212/218 (97%)	206 (97%)	6 (3%)	0	100	100
1	M	212/218 (97%)	206 (97%)	6 (3%)	0	100	100
1	O	210/218 (96%)	204 (97%)	6 (3%)	0	100	100
2	B	216/218 (99%)	211 (98%)	5 (2%)	0	100	100
2	D	216/218 (99%)	211 (98%)	5 (2%)	0	100	100
2	F	215/218 (99%)	210 (98%)	5 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	H	215/218 (99%)	210 (98%)	5 (2%)	0	100	100
2	J	216/218 (99%)	211 (98%)	5 (2%)	0	100	100
2	L	216/218 (99%)	211 (98%)	5 (2%)	0	100	100
2	N	216/218 (99%)	211 (98%)	5 (2%)	0	100	100
2	P	215/218 (99%)	210 (98%)	5 (2%)	0	100	100
3	Q	11/30 (37%)	11 (100%)	0	0	100	100
3	R	12/30 (40%)	12 (100%)	0	0	100	100
3	S	12/30 (40%)	11 (92%)	1 (8%)	0	100	100
3	T	10/30 (33%)	10 (100%)	0	0	100	100
3	U	12/30 (40%)	12 (100%)	0	0	100	100
3	V	10/30 (33%)	10 (100%)	0	0	100	100
3	W	7/30 (23%)	7 (100%)	0	0	100	100
3	X	12/30 (40%)	12 (100%)	0	0	100	100
All	All	3507/3728 (94%)	3418 (98%)	89 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	184/185 (100%)	176 (96%)	8 (4%)	26	54
1	C	185/185 (100%)	181 (98%)	4 (2%)	45	74
1	E	183/185 (99%)	178 (97%)	5 (3%)	39	69
1	G	181/185 (98%)	176 (97%)	5 (3%)	38	68
1	I	185/185 (100%)	177 (96%)	8 (4%)	26	54
1	K	184/185 (100%)	176 (96%)	8 (4%)	26	54
1	M	184/185 (100%)	179 (97%)	5 (3%)	39	69
1	O	182/185 (98%)	175 (96%)	7 (4%)	29	58

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	188/188 (100%)	181 (96%)	7 (4%)	30	59
2	D	188/188 (100%)	182 (97%)	6 (3%)	34	64
2	F	187/188 (100%)	182 (97%)	5 (3%)	39	69
2	H	187/188 (100%)	180 (96%)	7 (4%)	30	59
2	J	188/188 (100%)	182 (97%)	6 (3%)	34	64
2	L	188/188 (100%)	182 (97%)	6 (3%)	34	64
2	N	188/188 (100%)	183 (97%)	5 (3%)	39	69
2	P	187/188 (100%)	180 (96%)	7 (4%)	30	59
3	Q	7/12 (58%)	7 (100%)	0	100	100
3	R	7/12 (58%)	7 (100%)	0	100	100
3	S	7/12 (58%)	7 (100%)	0	100	100
3	T	7/12 (58%)	7 (100%)	0	100	100
3	U	7/12 (58%)	7 (100%)	0	100	100
3	V	6/12 (50%)	6 (100%)	0	100	100
3	W	5/12 (42%)	4 (80%)	1 (20%)	1	4
3	X	7/12 (58%)	7 (100%)	0	100	100
All	All	3022/3080 (98%)	2922 (97%)	100 (3%)	33	63

All (100) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	GLU
1	A	18	MET
1	A	80	VAL
1	A	131	CYS
1	A	135	THR
1	A	138	SER
1	A	180	LEU
1	A	193	SER
2	B	11	LEU
2	B	43	LYS
2	B	50	PHE
2	B	76	SER
2	B	111	LYS
2	B	150	VAL
2	B	180	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	18	MET
1	C	131	CYS
1	C	180	LEU
1	C	193	SER
2	D	11	LEU
2	D	50	PHE
2	D	76	SER
2	D	111	LYS
2	D	159	ARG
2	D	180	SER
1	E	1	GLU
1	E	18	MET
1	E	119	THR
1	E	180	LEU
1	E	193	SER
2	F	11	LEU
2	F	50	PHE
2	F	76	SER
2	F	111	LYS
2	F	180	SER
1	G	18	MET
1	G	80	VAL
1	G	135	THR
1	G	180	LEU
1	G	193	SER
2	H	11	LEU
2	H	43	LYS
2	H	50	PHE
2	H	76	SER
2	H	96	LYS
2	H	173	LYS
2	H	180	SER
1	I	6	GLU
1	I	18	MET
1	I	54	LYS
1	I	131	CYS
1	I	134	THR
1	I	180	LEU
1	I	189	ASN
1	I	193	SER
2	J	11	LEU
2	J	43	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	J	50	PHE
2	J	76	SER
2	J	126	SER
2	J	180	SER
1	K	6	GLU
1	K	18	MET
1	K	80	VAL
1	K	131	CYS
1	K	138	SER
1	K	180	LEU
1	K	189	ASN
1	K	193	SER
2	L	11	LEU
2	L	43	LYS
2	L	50	PHE
2	L	76	SER
2	L	111	LYS
2	L	180	SER
1	M	6	GLU
1	M	18	MET
1	M	131	CYS
1	M	180	LEU
1	M	193	SER
2	N	11	LEU
2	N	50	PHE
2	N	76	SER
2	N	126	SER
2	N	180	SER
1	O	6	GLU
1	O	18	MET
1	O	67	LYS
1	O	80	VAL
1	O	131	CYS
1	O	180	LEU
1	O	193	SER
2	P	11	LEU
2	P	43	LYS
2	P	50	PHE
2	P	76	SER
2	P	111	LYS
2	P	126	SER
2	P	180	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	W	17	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (44) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	56	ASN
1	A	58	HIS
2	B	17	GLN
2	B	141	ASN
2	B	194	ASN
1	C	56	ASN
1	C	57	ASN
1	C	58	HIS
1	C	167	HIS
2	D	141	ASN
2	D	194	ASN
2	F	17	GLN
2	F	194	ASN
2	F	214	ASN
2	F	216	ASN
1	G	58	HIS
1	G	167	HIS
2	H	17	GLN
2	H	141	ASN
2	H	194	ASN
2	H	214	ASN
1	I	56	ASN
1	I	58	HIS
2	J	17	GLN
2	J	141	ASN
2	J	194	ASN
1	K	39	GLN
1	K	56	ASN
1	K	58	HIS
1	K	86	ASN
2	L	42	GLN
2	L	141	ASN
2	L	194	ASN
1	M	39	GLN
1	M	167	HIS
2	N	17	GLN
2	N	42	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	N	141	ASN
2	N	142	ASN
2	N	194	ASN
1	O	56	ASN
1	O	58	HIS
2	P	142	ASN
2	P	149	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

12 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	HYP	U	24	3	7,8,9	0.40	0	5,10,12	2.37	3 (60%)
3	HYP	R	12	3	7,8,9	1.25	1 (14%)	5,10,12	1.43	1 (20%)
3	HYP	U	12	3	7,8,9	1.26	0	5,10,12	1.12	1 (20%)
3	HYP	R	9	3	7,8,9	0.68	0	5,10,12	2.72	3 (60%)
3	HYP	V	12	3	7,8,9	0.55	0	5,10,12	3.11	2 (40%)
3	HYP	Q	12	3	7,8,9	1.34	1 (14%)	5,10,12	1.20	0
3	HYP	S	9	3	7,8,9	0.60	0	5,10,12	2.65	2 (40%)
3	HYP	X	12	3	7,8,9	1.35	1 (14%)	5,10,12	1.55	0
3	HYP	Q	9	3	7,8,9	0.66	0	5,10,12	2.82	3 (60%)
3	HYP	X	9	3	7,8,9	0.93	0	5,10,12	2.23	2 (40%)
3	HYP	T	12	3	7,8,9	1.36	1 (14%)	5,10,12	1.17	1 (20%)
3	HYP	S	12	3	7,8,9	1.25	1 (14%)	5,10,12	1.33	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HYP	U	24	3	-	0/0/11/13	0/1/1/1
3	HYP	R	12	3	-	0/0/11/13	0/1/1/1
3	HYP	U	12	3	-	0/0/11/13	0/1/1/1
3	HYP	R	9	3	-	0/0/11/13	0/1/1/1
3	HYP	V	12	3	-	0/0/11/13	0/1/1/1
3	HYP	Q	12	3	-	0/0/11/13	0/1/1/1
3	HYP	S	9	3	-	0/0/11/13	0/1/1/1
3	HYP	X	12	3	-	0/0/11/13	0/1/1/1
3	HYP	Q	9	3	-	0/0/11/13	0/1/1/1
3	HYP	X	9	3	-	0/0/11/13	0/1/1/1
3	HYP	T	12	3	-	0/0/11/13	0/1/1/1
3	HYP	S	12	3	-	0/0/11/13	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	T	12	HYP	CD-N	2.34	1.55	1.47
3	Q	12	HYP	CD-N	2.33	1.55	1.47
3	S	12	HYP	CD-N	2.22	1.54	1.47
3	R	12	HYP	CD-N	2.21	1.54	1.47
3	X	12	HYP	CD-N	2.15	1.54	1.47

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	V	12	HYP	CG-CB-CA	6.06	110.76	103.75
3	R	9	HYP	CG-CB-CA	5.17	109.73	103.75
3	S	9	HYP	CB-CG-CD	5.05	108.78	103.16
3	Q	9	HYP	CG-CB-CA	4.96	109.48	103.75
3	X	9	HYP	CB-CG-CD	3.41	106.95	103.16
3	U	24	HYP	CG-CB-CA	3.38	107.66	103.75
3	Q	9	HYP	O-C-CA	-3.09	116.82	124.77
3	V	12	HYP	O-C-CA	-3.07	116.86	124.77
3	U	24	HYP	O-C-CA	-2.88	117.36	124.77
3	U	24	HYP	CB-CG-CD	2.87	106.35	103.16
3	X	9	HYP	CG-CB-CA	-2.84	100.47	103.75
3	S	9	HYP	CG-CB-CA	-2.34	101.05	103.75
3	S	12	HYP	O-C-CA	-2.28	118.91	124.77

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	Q	9	HYP	OD1-CG-CD	2.24	114.88	110.30
3	R	12	HYP	O-C-CA	-2.22	119.05	124.77
3	R	9	HYP	CB-CG-CD	2.19	105.59	103.16
3	U	12	HYP	CB-CG-CD	2.14	105.54	103.16
3	R	9	HYP	O-C-CA	-2.01	119.60	124.77
3	T	12	HYP	O-C-CA	-2.00	119.62	124.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	216/218 (99%)	-0.23	2 (0%) 81 80	28, 50, 75, 99	0
1	C	218/218 (100%)	-0.12	3 (1%) 73 72	29, 55, 81, 99	0
1	E	215/218 (98%)	-0.25	0 100 100	29, 47, 83, 106	0
1	G	211/218 (96%)	0.27	4 (1%) 66 63	28, 74, 121, 131	0
1	I	218/218 (100%)	-0.16	2 (0%) 81 80	23, 43, 91, 105	0
1	K	216/218 (99%)	-0.13	5 (2%) 61 58	35, 54, 79, 103	0
1	M	216/218 (99%)	-0.01	4 (1%) 66 63	39, 58, 79, 98	0
1	O	214/218 (98%)	0.20	6 (2%) 55 51	39, 65, 99, 117	0
2	B	218/218 (100%)	-0.14	1 (0%) 87 86	25, 47, 107, 132	0
2	D	218/218 (100%)	-0.31	1 (0%) 87 86	25, 43, 65, 78	0
2	F	217/218 (99%)	-0.08	1 (0%) 87 86	29, 56, 104, 127	0
2	H	217/218 (99%)	0.34	3 (1%) 73 72	31, 67, 133, 157	0
2	J	218/218 (100%)	0.03	1 (0%) 87 86	26, 58, 103, 118	0
2	L	218/218 (100%)	-0.03	1 (0%) 87 86	38, 59, 84, 109	0
2	N	218/218 (100%)	-0.03	0 100 100	39, 60, 80, 101	0
2	P	217/218 (99%)	0.01	2 (0%) 81 80	34, 54, 99, 114	0
3	Q	13/30 (43%)	0.44	2 (15%) 5 4	40, 48, 93, 128	0
3	R	13/30 (43%)	0.43	2 (15%) 5 4	38, 46, 90, 94	0
3	S	13/30 (43%)	0.31	1 (7%) 19 17	33, 44, 101, 102	0
3	T	12/30 (40%)	0.73	2 (16%) 4 3	47, 59, 86, 95	0
3	U	13/30 (43%)	0.98	1 (7%) 19 17	58, 65, 93, 93	0
3	V	11/30 (36%)	0.83	1 (9%) 15 12	57, 65, 97, 103	0
3	W	9/30 (30%)	0.42	0 100 100	50, 54, 62, 66	0
3	X	13/30 (43%)	0.40	1 (7%) 19 17	40, 50, 95, 98	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
All	All	3562/3728 (95%)	-0.02	46 (1%) 75 73	23, 55, 103, 157	0

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	U	26	PRO	3.9
1	K	135	THR	3.6
3	T	23	PRO	3.5
3	X	23	PRO	3.4
1	K	132	GLY	3.4
3	V	23	PRO	3.3
1	M	135	THR	3.2
1	O	217	GLY	3.2
1	I	174	LEU	3.1
2	P	33	GLY	3.0
3	R	23	PRO	3.0
1	O	132	GLY	2.9
1	G	174	LEU	2.9
1	A	135	THR	2.9
1	C	218	PRO	2.8
1	K	218	PRO	2.7
1	C	174	LEU	2.7
1	O	174	LEU	2.7
1	G	135	THR	2.7
1	M	218	PRO	2.7
2	B	133	GLY	2.6
1	G	129	PRO	2.6
3	T	11	PRO	2.6
2	H	213	PHE	2.6
1	A	218	PRO	2.5
3	Q	8	PRO	2.5
1	G	117	ALA	2.5
1	C	1	GLU	2.5
3	Q	22	GLY	2.5
1	I	218	PRO	2.4
2	J	206	THR	2.3
2	D	161	ASN	2.3
2	P	32	TYR	2.3
3	S	23	PRO	2.3
1	K	174	LEU	2.2
2	L	218	CYS	2.2
1	M	174	LEU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	O	130	VAL	2.2
1	O	68	GLY	2.1
1	M	132	GLY	2.1
2	H	212	SER	2.1
1	K	217	GLY	2.1
2	F	192	ARG	2.1
3	R	11	PRO	2.1
2	H	200	ALA	2.0
1	O	136	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	HYP	X	9	8/9	0.30	0.20	106,109,115,117	0
3	HYP	S	9	8/9	0.47	0.16	97,112,129,131	0
3	HYP	R	9	8/9	0.60	0.14	99,101,103,103	0
3	HYP	Q	9	8/9	0.68	0.16	94,104,109,110	0
3	HYP	U	24	8/9	0.80	0.13	93,93,94,94	0
3	HYP	U	12	8/9	0.84	0.13	84,90,91,92	0
3	HYP	V	12	8/9	0.85	0.13	77,77,78,78	0
3	HYP	R	12	8/9	0.87	0.14	67,72,77,78	0
3	HYP	Q	12	8/9	0.90	0.11	56,64,72,72	0
3	HYP	S	12	8/9	0.91	0.11	64,69,76,77	0
3	HYP	X	12	8/9	0.91	0.11	69,73,77,79	0
3	HYP	T	12	8/9	0.92	0.15	74,76,81,82	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.