



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 06:16 PM UTC

PDB ID : 7MU2 / pdb_00007mu2
Title : Crystal Structure of WIPI2 in complex with W2IR of ATG16L1
Authors : Strong, L.M.; Flower, T.G.; Buffalo, C.Z.; Hurley, J.H.
Deposited on : 2021-05-14
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	324	5% 85% 9% • 5%
1	C	324	4% 91% 6% • •
2	B	25	24% 68% 16% 16%
2	D	25	8% 48% 52%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 5665 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called WD repeat domain phosphoinositide-interacting protein 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	309	Total	C	N	O	S	0	0	0
			2356	1494	402	443	17			
1	C	316	Total	C	N	O	S	0	0	0
			2422	1534	416	453	19			

- Molecule 2 is a protein called Autophagy-related protein 16-1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	21	Total	C	N	O	0	0	0
			170	100	35	35			
2	D	12	Total	C	N	O	0	0	0
			92	56	17	19			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	206	TYR	-	insertion	UNP E7EVC7
D	206	TYR	-	insertion	UNP E7EVC7


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	309	Total	O	0	0
			309	309		
3	B	18	Total	O	0	0
			18	18		
3	C	292	Total	O	0	0
			292	292		
3	D	6	Total	O	0	0
			6	6		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: WD repeat domain phosphoinositide-interacting protein 2




- Molecule 1: WD repeat domain phosphoinositide-interacting protein 2



- Molecule 2: Autophagy-related protein 16-1



- Molecule 2: Autophagy-related protein 16-1



4 Data and refinement statistics

Property	Value	Source
Space group	I 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	117.77Å 49.08Å 120.13Å 90.00° 95.86° 90.00°	Depositor
Resolution (Å)	38.94 – 1.85 38.94 – 1.85	Depositor EDS
% Data completeness (in resolution range)	94.4 (38.94-1.85) 94.0 (38.94-1.85)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.51 (at 1.27Å)	Xtriage
Refinement program	PHENIX (1.18.2_3874: ???)	Depositor
R, R_{free}	0.183 , 0.217 0.182 , 0.217	Depositor DCC
R_{free} test set	2001 reflections (1.57%)	wwPDB-VP
Wilson B-factor (Å ²)	9.2	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 41.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.014 for l,-k,h	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5665	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.12% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.31	0/2398	0.54	0/3243
1	C	0.30	0/2464	0.53	0/3330
2	B	0.29	0/169	0.47	0/220
2	D	0.27	0/91	0.47	0/120
All	All	0.31	0/5122	0.53	0/6913

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2356	0	2342	23	1
1	C	2422	0	2431	17	0
2	B	170	0	170	4	0
2	D	92	0	90	0	0
3	A	309	0	0	9	4
3	B	18	0	0	1	1
3	C	292	0	0	8	1
3	D	6	0	0	0	0
All	All	5665	0	5033	43	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:134:ARG:NH1	3:A:403:HOH:O	2.09	0.85
1:A:229:ARG:O	3:A:401:HOH:O	1.96	0.84
1:C:54:ASP:OD1	3:C:401:HOH:O	1.99	0.81
1:A:228:LYS:O	3:A:402:HOH:O	2.04	0.74
1:A:225:ARG:NE	3:A:401:HOH:O	2.20	0.73
1:C:260:GLU:OE1	3:C:402:HOH:O	2.06	0.72
1:C:178:ASN:O	3:C:403:HOH:O	2.09	0.71
1:C:167:GLN:OE1	3:C:403:HOH:O	2.11	0.68
1:C:297:ASN:N	3:C:406:HOH:O	2.25	0.68
1:A:109:GLN:HG2	3:A:541:HOH:O	1.93	0.68
1:A:14:GLN:N	3:A:405:HOH:O	2.27	0.67
1:A:207:THR:HA	1:A:225:ARG:HD2	1.78	0.64
1:A:71:ALA:HB1	1:A:81:LEU:HD11	1.80	0.62
1:A:228:LYS:N	3:A:410:HOH:O	2.32	0.61
1:C:209:ILE:HD11	1:C:233:ILE:HD13	1.81	0.60
1:C:350:ALA:O	3:C:405:HOH:O	2.18	0.56
1:C:205:LYS:NZ	3:C:410:HOH:O	2.40	0.55
1:C:71:ALA:HB1	1:C:81:LEU:HD11	1.87	0.54
1:A:44:ASP:OD2	3:A:404:HOH:O	2.18	0.52
2:B:210:GLU:HG2	2:B:212:ASP:HB3	1.93	0.51
1:A:254:VAL:HB	1:A:305:VAL:HG22	1.91	0.51
1:A:97:TYR:HB3	1:A:115:LEU:HD21	1.94	0.50
1:A:57:ASP:HB3	1:A:75:LEU:HD12	1.94	0.49
1:C:11:MET:HE3	1:C:15:LEU:HB2	1.95	0.49
1:C:254:VAL:HB	1:C:305:VAL:HG22	1.94	0.49
2:B:215:ARG:NH2	3:B:303:HOH:O	2.46	0.48
1:A:225:ARG:H	1:A:225:ARG:HG3	1.45	0.46
2:B:211:LYS:HA	2:B:211:LYS:HD3	1.73	0.46
1:A:125:ARG:NH2	1:C:116:GLU:OE2	2.46	0.45
1:A:18:ALA:HA	1:A:28:ALA:O	2.16	0.45
1:A:57:ASP:HB3	1:A:75:LEU:HB2	1.99	0.45
1:A:252:GLU:H	1:A:252:GLU:CD	2.25	0.44
1:C:209:ILE:HD12	1:C:236:LEU:HD21	1.98	0.44
1:A:305:VAL:HG12	1:A:349:CYS:HB2	2.00	0.44
1:C:128:LYS:NZ	3:C:422:HOH:O	2.52	0.43
1:A:137:PRO:HA	1:A:138:PRO:HD3	1.95	0.42
1:A:225:ARG:NH2	3:A:425:HOH:O	2.53	0.42
1:C:305:VAL:HG12	1:C:349:CYS:HB2	2.01	0.42
1:A:115:LEU:HD22	1:A:120:TYR:HE1	1.85	0.41
2:B:210:GLU:N	2:B:210:GLU:OE1	2.53	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:316:CYS:SG	1:A:329:VAL:HG22	2.60	0.40
1:C:123:ASN:HB2	1:C:130:LEU:HD11	2.02	0.40
1:C:11:MET:CG	1:C:15:LEU:H	2.34	0.40

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:441:HOH:O	3:C:609:HOH:O[2_656]	1.91	0.29
3:A:634:HOH:O	3:A:675:HOH:O[4_556]	2.09	0.11
1:A:261:THR:OG1	1:A:261:THR:OG1[2_656]	2.11	0.09
3:A:453:HOH:O	3:A:634:HOH:O[4_546]	2.13	0.07
3:A:684:HOH:O	3:B:318:HOH:O[4_546]	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	305/324 (94%)	299 (98%)	4 (1%)	2 (1%)	18	8
1	C	312/324 (96%)	305 (98%)	7 (2%)	0	100	100
2	B	19/25 (76%)	18 (95%)	1 (5%)	0	100	100
2	D	10/25 (40%)	9 (90%)	1 (10%)	0	100	100
All	All	646/698 (93%)	631 (98%)	13 (2%)	2 (0%)	36	25

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	227	VAL
1	A	228	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	256/276 (93%)	249 (97%)	7 (3%)	39	24
1	C	266/276 (96%)	263 (99%)	3 (1%)	65	57
2	B	16/20 (80%)	16 (100%)	0	100	100
2	D	8/20 (40%)	8 (100%)	0	100	100
All	All	546/592 (92%)	536 (98%)	10 (2%)	51	40

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	43	VAL
1	A	54	ASP
1	A	115	LEU
1	A	225	ARG
1	A	300	ARG
1	A	305	VAL
1	A	357	LEU
1	C	115	LEU
1	C	297	ASN
1	C	305	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	354	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	309/324 (95%)	0.17	15 (4%) 35 38	11, 21, 41, 61	0
1	C	316/324 (97%)	0.18	14 (4%) 39 42	10, 21, 40, 62	0
2	B	21/25 (84%)	0.96	6 (28%) 1 1	14, 28, 73, 82	0
2	D	12/25 (48%)	0.96	2 (16%) 4 4	25, 34, 39, 54	0
All	All	658/698 (94%)	0.21	37 (5%) 30 33	10, 22, 42, 82	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	226	GLY	6.6
1	A	227	VAL	6.2
1	C	227	VAL	5.1
1	A	225	ARG	5.0
1	A	261	THR	4.1
1	A	17	PHE	3.7
1	C	228	LYS	3.6
1	C	226	GLY	3.6
1	C	352	MET	3.5
1	A	300	ARG	3.4
1	C	11	MET	3.2
1	C	12	ALA	3.0
1	A	54	ASP	2.9
1	A	299	GLY	2.9
1	A	230	CYS	2.9
1	A	16	LEU	2.9
1	A	15	LEU	2.7
1	C	13	GLY	2.7
2	B	213	SER	2.7
1	A	260	GLU	2.6
2	B	212	ASP	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	261	THR	2.5
2	B	211	LYS	2.5
2	B	216	ARG	2.5
1	C	135	GLU	2.4
1	C	229	ARG	2.4
2	D	219	ARG	2.4
2	B	215	ARG	2.3
1	A	117	GLU	2.3
2	D	220	LEU	2.3
1	A	53	THR	2.2
1	A	229	ARG	2.2
1	C	298	GLN	2.2
1	C	178	ASN	2.2
2	B	210	GLU	2.2
1	C	297	ASN	2.2
1	C	14	GLN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.