



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 1, 2026 – 01:29 PM UTC

PDB ID : 3MV2 / pdb_00003mv2
Title : Crystal Structure of α -COP in Complex with e-COP
Authors : Hoelz, A.; Hsia, K.C.
Deposited on : 2010-05-03
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	325	2% 42%40%10%6%
1	C	325	2% 45%38%10%7%
1	E	325	2% 45%38%10%7%
2	B	310	% 36%42%14%5%
2	D	310	2% 35%44%14%5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	310	4%35%45%14%5%

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 14273 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Coatomer subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	305	Total	C	N	O	S	0	0	0
			2405	1542	403	449	11			
1	C	303	Total	C	N	O	S	0	0	0
			2388	1531	399	447	11			
1	E	303	Total	C	N	O	S	0	0	0
			2388	1531	399	447	11			

There are 69 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-22	MET	-	expression tag	UNP P53622
A	-21	GLY	-	expression tag	UNP P53622
A	-20	SER	-	expression tag	UNP P53622
A	-19	SER	-	expression tag	UNP P53622
A	-18	HIS	-	expression tag	UNP P53622
A	-17	HIS	-	expression tag	UNP P53622
A	-16	HIS	-	expression tag	UNP P53622
A	-15	HIS	-	expression tag	UNP P53622
A	-14	HIS	-	expression tag	UNP P53622
A	-13	HIS	-	expression tag	UNP P53622
A	-12	SER	-	expression tag	UNP P53622
A	-11	SER	-	expression tag	UNP P53622
A	-10	GLY	-	expression tag	UNP P53622
A	-9	LEU	-	expression tag	UNP P53622
A	-8	GLU	-	expression tag	UNP P53622
A	-7	VAL	-	expression tag	UNP P53622
A	-6	LEU	-	expression tag	UNP P53622
A	-5	PHE	-	expression tag	UNP P53622
A	-4	GLN	-	expression tag	UNP P53622
A	-3	GLY	-	expression tag	UNP P53622
A	-2	PRO	-	expression tag	UNP P53622
A	-1	HIS	-	expression tag	UNP P53622
A	0	MET	-	expression tag	UNP P53622

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-22	MET	-	expression tag	UNP P53622
C	-21	GLY	-	expression tag	UNP P53622
C	-20	SER	-	expression tag	UNP P53622
C	-19	SER	-	expression tag	UNP P53622
C	-18	HIS	-	expression tag	UNP P53622
C	-17	HIS	-	expression tag	UNP P53622
C	-16	HIS	-	expression tag	UNP P53622
C	-15	HIS	-	expression tag	UNP P53622
C	-14	HIS	-	expression tag	UNP P53622
C	-13	HIS	-	expression tag	UNP P53622
C	-12	SER	-	expression tag	UNP P53622
C	-11	SER	-	expression tag	UNP P53622
C	-10	GLY	-	expression tag	UNP P53622
C	-9	LEU	-	expression tag	UNP P53622
C	-8	GLU	-	expression tag	UNP P53622
C	-7	VAL	-	expression tag	UNP P53622
C	-6	LEU	-	expression tag	UNP P53622
C	-5	PHE	-	expression tag	UNP P53622
C	-4	GLN	-	expression tag	UNP P53622
C	-3	GLY	-	expression tag	UNP P53622
C	-2	PRO	-	expression tag	UNP P53622
C	-1	HIS	-	expression tag	UNP P53622
C	0	MET	-	expression tag	UNP P53622
E	-22	MET	-	expression tag	UNP P53622
E	-21	GLY	-	expression tag	UNP P53622
E	-20	SER	-	expression tag	UNP P53622
E	-19	SER	-	expression tag	UNP P53622
E	-18	HIS	-	expression tag	UNP P53622
E	-17	HIS	-	expression tag	UNP P53622
E	-16	HIS	-	expression tag	UNP P53622
E	-15	HIS	-	expression tag	UNP P53622
E	-14	HIS	-	expression tag	UNP P53622
E	-13	HIS	-	expression tag	UNP P53622
E	-12	SER	-	expression tag	UNP P53622
E	-11	SER	-	expression tag	UNP P53622
E	-10	GLY	-	expression tag	UNP P53622
E	-9	LEU	-	expression tag	UNP P53622
E	-8	GLU	-	expression tag	UNP P53622
E	-7	VAL	-	expression tag	UNP P53622
E	-6	LEU	-	expression tag	UNP P53622
E	-5	PHE	-	expression tag	UNP P53622
E	-4	GLN	-	expression tag	UNP P53622

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	-3	GLY	-	expression tag	UNP P53622
E	-2	PRO	-	expression tag	UNP P53622
E	-1	HIS	-	expression tag	UNP P53622
E	0	MET	-	expression tag	UNP P53622

- Molecule 2 is a protein called Coatomer subunit epsilon.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	293	Total	C	N	O	S	0	0	0
			2364	1508	371	480	5			
2	D	293	Total	C	N	O	S	0	0	0
			2364	1508	371	480	5			
2	F	293	Total	C	N	O	S	0	0	0
			2364	1508	371	480	5			

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-13	MET	-	expression tag	UNP P40509
B	-12	GLY	-	expression tag	UNP P40509
B	-11	SER	-	expression tag	UNP P40509
B	-10	SER	-	expression tag	UNP P40509
B	-9	HIS	-	expression tag	UNP P40509
B	-8	HIS	-	expression tag	UNP P40509
B	-7	HIS	-	expression tag	UNP P40509
B	-6	HIS	-	expression tag	UNP P40509
B	-5	HIS	-	expression tag	UNP P40509
B	-4	HIS	-	expression tag	UNP P40509
B	-3	SER	-	expression tag	UNP P40509
B	-2	GLN	-	expression tag	UNP P40509
B	-1	ASP	-	expression tag	UNP P40509
B	0	PRO	-	expression tag	UNP P40509
D	-13	MET	-	expression tag	UNP P40509
D	-12	GLY	-	expression tag	UNP P40509
D	-11	SER	-	expression tag	UNP P40509
D	-10	SER	-	expression tag	UNP P40509
D	-9	HIS	-	expression tag	UNP P40509
D	-8	HIS	-	expression tag	UNP P40509
D	-7	HIS	-	expression tag	UNP P40509
D	-6	HIS	-	expression tag	UNP P40509
D	-5	HIS	-	expression tag	UNP P40509
D	-4	HIS	-	expression tag	UNP P40509

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-3	SER	-	expression tag	UNP P40509
D	-2	GLN	-	expression tag	UNP P40509
D	-1	ASP	-	expression tag	UNP P40509
D	0	PRO	-	expression tag	UNP P40509
F	-13	MET	-	expression tag	UNP P40509
F	-12	GLY	-	expression tag	UNP P40509
F	-11	SER	-	expression tag	UNP P40509
F	-10	SER	-	expression tag	UNP P40509
F	-9	HIS	-	expression tag	UNP P40509
F	-8	HIS	-	expression tag	UNP P40509
F	-7	HIS	-	expression tag	UNP P40509
F	-6	HIS	-	expression tag	UNP P40509
F	-5	HIS	-	expression tag	UNP P40509
F	-4	HIS	-	expression tag	UNP P40509
F	-3	SER	-	expression tag	UNP P40509
F	-2	GLN	-	expression tag	UNP P40509
F	-1	ASP	-	expression tag	UNP P40509
F	0	PRO	-	expression tag	UNP P40509


3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Coatomer subunit alpha





• Molecule 1: Coatomer subunit alpha







• Molecule 1: Coatomer subunit alpha




• Molecule 2: Coatomer subunit epsilon



- Molecule 2: Coatomer subunit epsilon



- Molecule 2: Coatomer subunit epsilon



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	329.13Å 74.38Å 97.25Å 90.00° 102.27° 90.00°	Depositor
Resolution (Å)	20.00 – 2.90 20.00 – 2.90	Depositor EDS
% Data completeness (in resolution range)	85.4 (20.00-2.90) 85.6 (20.00-2.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.15 (at 2.88Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.246 , 0.297 0.246 , 0.299	Depositor DCC
R_{free} test set	2080 reflections (4.39%)	wwPDB-VP
Wilson B-factor (Å ²)	76.7	Xtriage
Anisotropy	0.578	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 81.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	14273	wwPDB-VP
Average B, all atoms (Å ²)	125.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.37% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.71	0/2454	1.13	13/3317 (0.4%)
1	C	0.70	0/2435	1.13	14/3291 (0.4%)
1	E	0.66	0/2435	1.12	12/3291 (0.4%)
2	B	0.68	0/2403	1.19	26/3261 (0.8%)
2	D	0.67	0/2403	1.14	21/3261 (0.6%)
2	F	0.64	0/2403	1.14	21/3261 (0.6%)
All	All	0.68	0/14533	1.14	107/19682 (0.5%)

There are no bond length outliers.

The worst 5 of 107 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	264	GLN	N-CA-C	-10.92	96.31	112.04
2	D	264	GLN	N-CA-C	-10.90	96.34	112.04
2	B	264	GLN	N-CA-C	-10.86	96.40	112.04
1	A	961	LEU	N-CA-C	-10.12	97.23	110.39
1	C	961	LEU	N-CA-C	-10.06	97.31	110.39

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2405	0	2455	169	0
1	C	2388	0	2441	144	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	2388	0	2441	154	0
2	B	2364	0	2319	152	0
2	D	2364	0	2319	173	0
2	F	2364	0	2319	175	0
All	All	14273	0	14294	919	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 32.

The worst 5 of 919 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1166:ASP:OD2	1:A:1169:THR:HB	1.59	1.03
1:E:1166:ASP:OD2	1:E:1169:THR:HB	1.59	1.03
1:C:1166:ASP:OD2	1:C:1169:THR:HB	1.59	1.02
2:D:248:ILE:HD11	2:D:263:ASN:HB3	1.43	0.99
2:F:248:ILE:HD11	2:F:263:ASN:HB3	1.43	0.98

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	303/325 (93%)	256 (84%)	32 (11%)	15 (5%)	1	6
1	C	301/325 (93%)	255 (85%)	31 (10%)	15 (5%)	1	6
1	E	301/325 (93%)	255 (85%)	31 (10%)	15 (5%)	1	6
2	B	291/310 (94%)	227 (78%)	41 (14%)	23 (8%)	1	2
2	D	291/310 (94%)	226 (78%)	44 (15%)	21 (7%)	1	2
2	F	291/310 (94%)	228 (78%)	42 (14%)	21 (7%)	1	2

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	1778/1905 (93%)	1447 (81%)	221 (12%)	110 (6%)	1	3

5 of 110 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	957	THR
1	A	973	ASP
1	A	1114	SER
1	A	1133	ALA
1	A	1186	MET

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	259/276 (94%)	232 (90%)	27 (10%)	7	23
1	C	257/276 (93%)	230 (90%)	27 (10%)	6	22
1	E	257/276 (93%)	230 (90%)	27 (10%)	6	22
2	B	263/279 (94%)	224 (85%)	39 (15%)	3	10
2	D	263/279 (94%)	223 (85%)	40 (15%)	3	9
2	F	263/279 (94%)	224 (85%)	39 (15%)	3	10
All	All	1562/1665 (94%)	1363 (87%)	199 (13%)	4	14

5 of 199 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	201	LEU
1	E	1042	ARG
2	D	225	SER
2	D	271	GLU
1	E	1080	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 35 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	1101	GLN
1	E	1127	ASN
2	F	176	ASN
1	C	979	GLN
1	C	930	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	305/325 (93%)	0.05	5 (1%) 70 62	55, 89, 144, 199	0
1	C	303/325 (93%)	0.15	8 (2%) 57 48	57, 97, 167, 201	0
1	E	303/325 (93%)	0.27	7 (2%) 61 52	98, 157, 196, 201	0
2	B	293/310 (94%)	0.08	3 (1%) 79 73	59, 99, 166, 201	0
2	D	293/310 (94%)	0.12	5 (1%) 69 61	64, 104, 172, 201	0
2	F	293/310 (94%)	0.50	13 (4%) 39 30	106, 179, 201, 201	0
All	All	1790/1905 (93%)	0.19	41 (2%) 61 52	55, 119, 197, 201	0

The worst 5 of 41 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1102	ALA	4.6
1	C	1094	PHE	3.6
2	D	258	THR	3.4
1	A	1129	ALA	3.3
2	F	75	LEU	3.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.