



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:46 AM UTC

PDB ID : 5MVJ / pdb_00005mvj
Title : Structure of DC8E8 Fab at pH 6.5 crystallized in space-group P1
Authors : Skrabana, R.; Novak, M.; Cehlar, O.; Kontsekova, E.
Deposited on : 2017-01-16
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	221	 93% 5% .
1	C	221	 88% 8% . .
1	E	221	 86% 12% .
1	H	221	 87% 10% . .
2	B	219	 89% 11%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	219	 89%10%
2	F	219	 89%10%
2	L	219	 89%11%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 13492 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called antibody Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	H	219	Total	C	N	O	S	0	0	0
			1654	1052	266	328	8			
1	A	219	Total	C	N	O	S	0	1	0
			1662	1057	268	329	8			
1	C	214	Total	C	N	O	S	0	0	0
			1621	1033	259	321	8			
1	E	216	Total	C	N	O	S	0	1	0
			1639	1043	260	327	9			

- Molecule 2 is a protein called antibody kappa chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	219	Total	C	N	O	S	0	0	0
			1697	1056	288	345	8			
2	B	218	Total	C	N	O	S	0	1	0
			1679	1048	281	343	7			
2	D	218	Total	C	N	O	S	0	1	0
			1692	1055	284	346	7			
2	F	218	Total	C	N	O	S	0	0	0
			1680	1047	283	343	7			

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	F	1	Total	Cl	0	0
			1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	H	22	Total	O	0	0
			22	22		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	L	19	Total 19	O 19	0	0
4	A	17	Total 17	O 17	0	0
4	B	15	Total 15	O 15	0	0
4	C	29	Total 29	O 29	0	0
4	D	22	Total 22	O 22	0	0
4	E	22	Total 22	O 22	0	0
4	F	21	Total 21	O 21	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: antibody Fab heavy chain




- Molecule 1: antibody Fab heavy chain




- Molecule 1: antibody Fab heavy chain




- Molecule 1: antibody Fab heavy chain



- Molecule 2: antibody kappa chain




- Molecule 2: antibody kappa chain






• Molecule 2: antibody kappa chain



• Molecule 2: antibody kappa chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	61.22Å 82.26Å 89.31Å 92.32° 95.47° 89.82°	Depositor
Resolution (Å)	34.37 – 2.50 34.37 – 2.50	Depositor EDS
% Data completeness (in resolution range)	86.1 (34.37-2.50) 85.9 (34.37-2.50)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.55 (at 2.22Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.219 , 0.272 0.225 , 0.274	Depositor DCC
R_{free} test set	3942 reflections (3.33%)	wwPDB-VP
Wilson B-factor (Å ²)	28.5	Xtriage
Anisotropy	0.221	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 21.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	0.080 for -h,k,-l	Xtriage
Reported twinning fraction	0.875 for H, K, L 0.125 for -H, K, -L	Depositor
Outliers	0 of 81441 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	13492	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, PCA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.83	0/1702	0.93	0/2325
1	C	0.87	0/1658	0.96	1/2267 (0.0%)
1	E	0.84	0/1676	0.97	0/2291
1	H	0.90	0/1691	0.99	5/2311 (0.2%)
2	B	0.79	0/1718	0.93	1/2331 (0.0%)
2	D	0.88	0/1731	0.98	3/2347 (0.1%)
2	F	0.83	0/1716	0.93	1/2328 (0.0%)
2	L	0.84	0/1733	0.93	1/2349 (0.0%)
All	All	0.85	0/13625	0.95	12/18549 (0.1%)

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	8	GLY	CA-C-N	8.21	128.14	119.85
1	H	8	GLY	C-N-CA	8.21	128.14	119.85
2	B	143	ASN	N-CA-C	7.24	121.38	111.39
2	D	49	SER	CA-C-N	-6.50	113.97	120.21
2	D	49	SER	C-N-CA	-6.50	113.97	120.21
2	D	205	THR	N-CA-C	5.90	120.09	113.01
1	C	6	GLN	N-CA-C	5.28	118.60	110.42
2	L	49	SER	N-CA-C	-5.27	103.50	110.40
1	H	70	LEU	N-CA-C	5.27	117.31	108.73
2	F	217	ASN	N-CA-C	5.11	116.85	111.28
1	H	111	GLY	N-CA-C	-5.08	103.77	112.30
1	H	67	LYS	N-CA-C	-5.01	106.26	112.93

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1662	0	1619	5	1
1	C	1621	0	1574	9	0
1	E	1639	0	1586	10	1
1	H	1654	0	1608	11	0
2	B	1679	0	1607	9	0
2	D	1692	0	1629	8	0
2	F	1680	0	1607	10	0
2	L	1697	0	1637	11	0
3	F	1	0	0	0	0
4	A	17	0	0	0	0
4	B	15	0	0	0	0
4	C	29	0	0	0	0
4	D	22	0	0	0	0
4	E	22	0	0	0	0
4	F	21	0	0	1	0
4	H	22	0	0	0	0
4	L	19	0	0	1	0
All	All	13492	0	12867	69	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:39:GLN:HE22	2:D:44:GLN:HE22	1.26	0.82
2:F:206:SER:OG	2:F:208:SER:O	2.05	0.75
2:F:193:ARG:NH1	4:F:401:HOH:O	2.27	0.68
1:H:60:TYR:HB2	1:H:65:LYS:HE2	1.77	0.65
1:H:62:GLU:HA	1:H:65:LYS:HD2	1.79	0.64
2:L:67:ARG:NH1	4:L:301:HOH:O	2.32	0.63
2:L:160:ARG:NH1	2:L:162:ASN:O	2.33	0.62
1:C:195:TRP:CG	1:C:196:PRO:HA	2.39	0.57
1:H:9:PRO:HD2	1:H:207:PRO:HB2	1.89	0.55
2:D:43:GLN:HB2	2:D:53:LEU:HD11	1.89	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:12:ALA:HA	2:L:110:ASP:O	2.07	0.54
1:C:12:VAL:HG21	1:C:18:VAL:HB	1.88	0.54
2:B:43:GLN:HB2	2:B:53:LEU:HD11	1.90	0.54
2:L:127:SER:O	2:L:131:THR:HG23	2.07	0.54
1:E:103:THR:OG1	1:E:104:SER:N	2.40	0.54
2:B:37:ASN:O	2:B:56:TRP:HA	2.10	0.51
2:B:11:LEU:HD21	2:B:19:VAL:HG22	1.93	0.50
2:B:160:ARG:NH1	2:B:162:ASN:O	2.45	0.50
2:F:45:LYS:HE2	2:F:87:GLU:O	2.11	0.49
1:H:133:PRO:HD3	1:H:145:LEU:HD23	1.94	0.49
1:H:2:VAL:HB	1:H:109:TYR:CE2	2.47	0.49
1:H:9:PRO:CD	1:H:207:PRO:HB2	2.43	0.49
1:H:161:TRP:CZ3	1:H:202:CYS:HB3	2.48	0.49
1:E:142:MET:HE3	1:E:189:THR:HG22	1.94	0.48
2:F:29:LEU:HD12	2:F:77:PHE:CE2	2.49	0.48
1:H:98:ARG:O	1:H:107:MET:HA	2.12	0.48
2:F:141:LEU:HD23	2:F:201:ALA:HB2	1.95	0.48
2:F:11:LEU:HD21	2:F:19:VAL:HG22	1.96	0.47
1:E:36:TRP:CD1	1:E:70:LEU:HD22	2.49	0.47
2:F:37:ASN:O	2:F:56:TRP:HA	2.14	0.47
1:E:54:ARG:O	1:E:74:LYS:NZ	2.48	0.46
1:E:123:THR:CG2	1:E:209:SER:HB3	2.45	0.46
2:D:113:ARG:HD2	2:D:176:SER:HB2	1.97	0.46
1:A:107:MET:HE3	2:B:95:LYS:HE2	1.97	0.46
1:A:161:TRP:CZ3	1:A:202:CYS:HB3	2.50	0.46
1:H:47:TRP:CG	2:L:101:ARG:HB2	2.52	0.45
1:A:141:SER:O	1:A:192:SER:HB2	2.17	0.45
2:D:186:LEU:HD22	2:D:190:GLU:HG2	1.99	0.44
1:E:61:ASN:OD1	1:E:62:GLU:N	2.50	0.44
2:F:141:LEU:CD2	2:F:201:ALA:HB2	2.47	0.44
2:L:37:ASN:O	2:L:56:TRP:HA	2.17	0.44
2:B:11:LEU:HD21	2:B:19:VAL:CG2	2.47	0.44
2:B:154:LYS:HB2	2:B:198:THR:HB	1.99	0.44
1:C:179:SER:C	1:C:181:LEU:H	2.25	0.44
2:D:129:GLN:HE22	2:D:136:SER:CB	2.31	0.44
1:E:12:VAL:O	1:E:118:VAL:HA	2.18	0.43
2:L:45:LYS:HE2	2:L:87:GLU:O	2.18	0.43
2:L:34:THR:O	2:L:35:ARG:HB2	2.19	0.43
1:E:47:TRP:CG	2:F:101:ARG:HB2	2.54	0.43
2:L:95:LYS:HA	2:L:102:THR:O	2.20	0.42
2:F:128:GLU:OE1	2:F:128:GLU:N	2.52	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:161:TRP:CH2	1:C:202:CYS:HB3	2.54	0.42
1:C:195:TRP:CD1	1:C:196:PRO:HA	2.54	0.42
1:H:91:SER:HA	1:H:116:VAL:O	2.19	0.42
1:A:9:PRO:CD	1:A:207:PRO:HB2	2.50	0.42
2:L:145:TYR:CD1	2:L:146:PRO:HA	2.55	0.41
2:D:34:THR:OG1	2:D:36:LYS:HD2	2.21	0.41
2:L:141:LEU:N	2:L:141:LEU:HD12	2.36	0.41
2:B:51[B]:LYS:HE3	2:B:51[B]:LYS:HB3	1.46	0.41
2:D:200:GLU:HG2	2:D:211:VAL:HG22	2.03	0.41
1:C:152:TYR:OH	1:C:175:ALA:HB2	2.21	0.41
2:D:186:LEU:HD22	2:D:190:GLU:CG	2.51	0.41
1:H:142:MET:HE3	1:H:189:THR:HG22	2.01	0.40
1:C:154:PRO:HD2	1:C:208:ALA:CB	2.51	0.40
2:B:141:LEU:N	2:B:141:LEU:HD12	2.36	0.40
1:C:161:TRP:CZ3	1:C:202:CYS:HB3	2.57	0.40
1:E:152:TYR:CE2	1:E:157:VAL:HG13	2.56	0.40
1:E:194:THR:O	1:E:198:GLU:N	2.54	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:216:LYS:NZ	1:E:163:SER:OG[1_454]	2.15	0.05

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	216/221 (98%)	209 (97%)	7 (3%)	0	100	100
1	C	210/221 (95%)	202 (96%)	8 (4%)	0	100	100
1	E	213/221 (96%)	207 (97%)	6 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	215/221 (97%)	203 (94%)	12 (6%)	0	100	100
2	B	217/219 (99%)	210 (97%)	7 (3%)	0	100	100
2	D	217/219 (99%)	208 (96%)	8 (4%)	1 (0%)	24	43
2	F	216/219 (99%)	206 (95%)	10 (5%)	0	100	100
2	L	217/219 (99%)	210 (97%)	7 (3%)	0	100	100
All	All	1721/1760 (98%)	1655 (96%)	65 (4%)	1 (0%)	48	68

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	193	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	188/190 (99%)	183 (97%)	5 (3%)	39	67
1	C	184/190 (97%)	177 (96%)	7 (4%)	29	56
1	E	187/190 (98%)	181 (97%)	6 (3%)	34	62
1	H	187/190 (98%)	180 (96%)	7 (4%)	30	57
2	B	190/195 (97%)	181 (95%)	9 (5%)	23	47
2	D	193/195 (99%)	187 (97%)	6 (3%)	35	62
2	F	190/195 (97%)	184 (97%)	6 (3%)	34	62
2	L	194/195 (100%)	189 (97%)	5 (3%)	40	68
All	All	1513/1540 (98%)	1462 (97%)	51 (3%)	32	60

All (51) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	65	LYS
1	H	84	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	H	123	THR
1	H	145	LEU
1	H	184	LEU
1	H	202	CYS
1	H	218	VAL
2	L	128	GLU
2	L	158	SER
2	L	161	GLN
2	L	165	LEU
2	L	180	MET
1	A	17	SER
1	A	85	SER
1	A	147	CYS
1	A	168	SER
1	A	192	SER
2	B	20	THR
2	B	32	SER
2	B	39	LEU
2	B	69	THR
2	B	165	LEU
2	B	186	LEU
2	B	190	GLU
2	B	207	THR
2	B	208	SER
1	C	12	VAL
1	C	35	SER
1	C	120	SER
1	C	179	SER
1	C	184	LEU
1	C	185	SER
1	C	197	SER
2	D	14	SER
2	D	39	LEU
2	D	54	ILE
2	D	97	SER
2	D	127	SER
2	D	208	SER
1	E	35	SER
1	E	41	THR
1	E	184	LEU
1	E	186	SER
1	E	193	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	203	ASN
2	F	20	THR
2	F	54	ILE
2	F	97	SER
2	F	173	SER
2	F	186	LEU
2	F	218	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	H	77	ASN
2	L	143	ASN
2	L	195	ASN
2	L	215	ASN
1	A	5	GLN
1	A	171	HIS
2	B	31	ASN
2	B	143	ASN
1	C	112	GLN
1	C	178	GLN
2	D	44	GLN
2	D	85	GLN
2	D	129	GLN
2	D	143	ASN
1	E	5	GLN
1	E	39	GLN
1	E	77	ASN
1	E	171	HIS
2	F	44	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	PCA	E	1	1	7,8,9	0.68	0	9,10,12	0.82	0
1	PCA	A	1	1	7,8,9	0.66	0	9,10,12	0.97	0
1	PCA	H	1	1	7,8,9	0.58	0	9,10,12	1.00	0
1	PCA	C	1	1	7,8,9	0.71	0	9,10,12	1.12	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PCA	E	1	1	-	0/0/11/13	0/1/1/1
1	PCA	A	1	1	-	0/0/11/13	0/1/1/1
1	PCA	H	1	1	-	0/0/11/13	0/1/1/1
1	PCA	C	1	1	-	0/0/11/13	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	218/221 (98%)	-1.20	0 100 100	17, 28, 49, 82	1 (0%)
1	C	213/221 (96%)	-1.21	0 100 100	13, 30, 56, 77	0
1	E	215/221 (97%)	-1.20	0 100 100	15, 28, 57, 89	1 (0%)
1	H	218/221 (98%)	-1.24	0 100 100	15, 27, 51, 73	0
2	B	218/219 (99%)	-1.10	0 100 100	10, 39, 64, 78	1 (0%)
2	D	218/219 (99%)	-1.14	0 100 100	14, 34, 76, 96	1 (0%)
2	F	218/219 (99%)	-1.20	0 100 100	12, 31, 64, 77	0
2	L	219/219 (100%)	-1.30	0 100 100	15, 26, 46, 78	0
All	All	1737/1760 (98%)	-1.20	0 100 100	10, 30, 62, 96	4 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	PCA	H	1	8/9	0.98	0.04	31,35,36,37	0
1	PCA	A	1	8/9	0.99	0.05	32,43,46,52	0
1	PCA	C	1	8/9	0.99	0.03	23,23,25,25	0
1	PCA	E	1	8/9	0.99	0.04	36,36,38,45	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	CL	F	301	1/1	0.99	0.04	39,39,39,39	0

6.5 Other polymers [i](#)

There are no such residues in this entry.