



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 12:47 PM UTC

PDB ID : 1MWE / pdb_00001mwe
Title : THE X-RAY STRUCTURE OF A COMPLEX OF TERN N9 INFLUENZA VIRUS NEURAMINIDASE COMPLEXED WITH SIALIC ACID AT 4 DEGREES C REVEALING A SECOND SIALIC ACID BINDING SITE
Authors : Varghese, J.N.
Deposited on : 1997-08-27
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	388	
2	B	9	

2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 3654 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NEURAMINIDASE.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	388	3067	1914	538	592	23	8	0	0

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	9	105	58	2	45	0	0	0

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is N-acetyl-alpha-neuraminic acid (CCD ID: SIA) (formula: C₁₁H₁₉NO₉).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	1	0
			21	11	1	9		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			21	11	1	9		

- Molecule 5 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Ca	0	0
			2	2		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	396	Total	O	0	0
			396	396		

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: NEURAMINIDASE



- Molecule 2: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	I 4 3 2	Depositor
Cell constants a, b, c, α , β , γ	181.04Å 181.04Å 181.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	6.00 – 1.70	Depositor
% Data completeness (in resolution range)	96.6 (6.00-1.70)	Depositor
R_{merge}	0.08	Depositor
R_{sym}	0.09	Depositor
Refinement program	X-PLOR 3.0	Depositor
R, R_{free}	(Not available) , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3654	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, CA, BMA, NAG, SIA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.75	0/3150	1.08	20/4290 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	226	GLN	N-CA-C	11.80	123.69	111.07
1	A	125	ASP	N-CA-C	-8.60	99.99	110.31
1	A	299	ASN	N-CA-C	-7.41	99.55	110.52
1	A	319	SER	N-CA-C	7.36	118.07	109.60
1	A	245	SER	N-CA-C	6.98	120.08	110.24
1	A	297	GLY	N-CA-C	6.41	123.82	111.31
1	A	151	ASP	N-CA-C	5.79	119.53	112.23
1	A	105	ALA	N-CA-C	5.78	117.25	111.07
1	A	180	SER	N-CA-C	5.63	116.82	108.60
1	A	365	THR	N-CA-C	-5.48	102.77	110.50
1	A	368	ILE	N-CA-C	-5.45	106.52	113.22
1	A	453	LEU	N-CA-C	5.45	118.18	110.50
1	A	323	THR	N-CA-C	5.40	120.56	112.94
1	A	223	LEU	N-CA-C	-5.30	102.45	110.24
1	A	434	ASP	N-CA-C	5.29	119.60	113.20
1	A	229	GLU	N-CA-C	5.29	117.90	110.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	280	CYS	N-CA-C	5.14	117.49	109.52
1	A	113	ASP	N-CA-C	5.07	117.46	109.39
1	A	165	SER	N-CA-C	-5.07	102.79	110.39
1	A	90	GLY	N-CA-C	-5.05	103.87	112.51

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	423	TYR	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3067	0	2893	40	0
2	B	105	0	88	2	0
3	A	42	0	38	6	0
4	A	42	0	36	0	0
5	A	2	0	0	0	0
6	A	396	0	0	3	2
All	All	3654	0	3055	47	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:172:ARG:HD3	1:A:209:ARG:NH2	1.78	0.97
3:A:478(A):NAG:O4	3:A:479(B):NAG:C1	2.15	0.95
3:A:480(A):NAG:O7	6:A:1385:HOH:O	1.90	0.89
1:A:168:THR:H	1:A:170:ASN:HD21	1.22	0.87
1:A:87:LEU:H	1:A:233:HIS:HD2	1.17	0.87
1:A:98:HIS:HE1	1:A:419:ARG:HH11	1.29	0.80

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:170:ASN:HB3	6:A:1303:HOH:O	1.88	0.73
1:A:98:HIS:CE1	1:A:419:ARG:HH11	2.08	0.71
1:A:97:TRP:H	1:A:395:GLN:HE22	1.44	0.65
1:A:465:GLU:H	1:A:465:GLU:CD	2.05	0.64
1:A:87:LEU:H	1:A:233:HIS:CD2	2.09	0.63
1:A:221:ASN:HB3	1:A:244:GLY:HA2	1.79	0.63
1:A:84:PHE:CE1	1:A:187:LYS:HD2	2.33	0.63
3:A:478(A):NAG:C4	3:A:479(B):NAG:C1	2.76	0.62
1:A:292:ARG:HE	1:A:294:ASN:HD22	1.47	0.61
1:A:172:ARG:CD	1:A:209:ARG:NH2	2.60	0.59
1:A:292:ARG:HH21	1:A:294:ASN:ND2	2.04	0.55
1:A:168:THR:H	1:A:170:ASN:ND2	1.98	0.55
1:A:170:ASN:HD22	1:A:170:ASN:H	1.55	0.55
1:A:144:HIS:HD2	6:A:1036:HOH:O	1.91	0.53
1:A:168:THR:OG1	1:A:170:ASN:ND2	2.42	0.52
1:A:188:THR:HG22	1:A:207:TYR:CZ	2.45	0.52
1:A:228:SER:HB3	1:A:350:LYS:HE2	1.92	0.51
1:A:437:TRP:CD1	3:A:480(A):NAG:H82	2.46	0.51
1:A:172:ARG:HD3	1:A:209:ARG:HH22	1.68	0.51
3:A:478(A):NAG:O4	3:A:479(B):NAG:N2	2.43	0.51
1:A:364:ARG:HD2	1:A:375:GLU:OE2	2.12	0.49
3:A:478(A):NAG:O4	3:A:479(B):NAG:C2	2.60	0.49
1:A:328:PRO:HG3	1:A:343:GLY:HA3	1.95	0.49
1:A:174:GLU:OE1	1:A:209:ARG:NH2	2.47	0.47
1:A:422:PHE:CZ	1:A:446:MET:HG3	2.50	0.46
1:A:121:TYR:CG	1:A:228:SER:HA	2.51	0.46
1:A:98:HIS:CE1	1:A:419:ARG:NH1	2.80	0.45
1:A:172:ARG:HD2	1:A:174:GLU:OE2	2.17	0.44
1:A:172:ARG:HD3	1:A:209:ARG:CZ	2.45	0.43
1:A:292:ARG:HH21	1:A:294:ASN:HD21	1.66	0.43
1:A:378:LYS:O	1:A:390:PRO:HA	2.19	0.43
2:B:7:MAN:C6	2:B:9:MAN:H5	2.48	0.43
1:A:188:THR:CG2	1:A:207:TYR:CZ	3.02	0.42
1:A:98:HIS:HD2	1:A:99:ILE:O	2.02	0.42
1:A:367:SER:HB2	1:A:400:ASN:HD21	1.84	0.42
1:A:238:PRO:HA	1:A:255:TYR:O	2.20	0.42
1:A:278:CYS:HB3	1:A:289:CYS:HB3	2.02	0.41
2:B:7:MAN:H61	2:B:9:MAN:H5	2.03	0.41
1:A:221:ASN:HB3	1:A:244:GLY:CA	2.49	0.41
1:A:207:TYR:CZ	1:A:259:GLU:HA	2.57	0.40
1:A:315:GLN:HG3	1:A:316:TYR:N	2.36	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:1307:HOH:O	6:A:1308:HOH:O[48_555]	0.47	1.73
6:A:1265:HOH:O	6:A:1265:HOH:O[48_555]	0.80	1.40

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	386/388 (100%)	367 (95%)	19 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	341/341 (100%)	335 (98%)	6 (2%)	51	36

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	170	ASN
1	A	204	VAL
1	A	304	ARG
1	A	333	VAL
1	A	364	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	465	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	98	HIS
1	A	144	HIS
1	A	170	ASN
1	A	221	ASN
1	A	233	HIS
1	A	294	ASN
1	A	296	GLN
1	A	381	ASN
1	A	395	GLN
1	A	400	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

9 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	B	1	2,1	14,14,15	1.39	2 (14%)	17,19,21	1.32	2 (11%)
2	NAG	B	2	2	14,14,15	1.49	2 (14%)	17,19,21	1.70	4 (23%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BMA	B	3	2	11,11,12	0.90	0	15,15,17	0.91	0
2	MAN	B	4	2	11,11,12	1.26	1 (9%)	15,15,17	1.70	4 (26%)
2	MAN	B	5	2	11,11,12	0.89	0	15,15,17	1.07	1 (6%)
2	MAN	B	6	2	11,11,12	2.27	4 (36%)	15,15,17	0.94	1 (6%)
2	MAN	B	7	2	11,11,12	1.12	0	15,15,17	1.32	2 (13%)
2	MAN	B	8	2	11,11,12	2.45	2 (18%)	15,15,17	2.20	4 (26%)
2	MAN	B	9	2	11,11,12	1.78	2 (18%)	15,15,17	1.31	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	0/6/23/26	0/1/1/1
2	BMA	B	3	2	-	0/2/19/22	0/1/1/1
2	MAN	B	4	2	-	0/2/19/22	0/1/1/1
2	MAN	B	5	2	-	0/2/19/22	0/1/1/1
2	MAN	B	6	2	-	0/2/19/22	0/1/1/1
2	MAN	B	7	2	-	0/2/19/22	0/1/1/1
2	MAN	B	8	2	-	0/2/19/22	0/1/1/1
2	MAN	B	9	2	-	0/2/19/22	0/1/1/1

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	8	MAN	C1-C2	7.44	1.70	1.52
2	B	9	MAN	C1-C2	4.59	1.63	1.52
2	B	6	MAN	O5-C5	4.03	1.51	1.43
2	B	6	MAN	C2-C3	3.82	1.58	1.52
2	B	6	MAN	C4-C5	3.75	1.61	1.53
2	B	1	NAG	C4-C5	3.49	1.60	1.53
2	B	2	NAG	C4-C5	-3.27	1.46	1.53
2	B	9	MAN	O5-C1	-3.25	1.38	1.43
2	B	8	MAN	O5-C1	2.88	1.48	1.43
2	B	6	MAN	C4-C3	2.84	1.59	1.52
2	B	4	MAN	C2-C3	2.49	1.56	1.52
2	B	2	NAG	C3-C2	-2.25	1.47	1.52
2	B	1	NAG	C4-C3	-2.01	1.47	1.52

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	8	MAN	C1-O5-C5	5.36	119.38	112.19
2	B	8	MAN	O4-C4-C3	-4.03	100.89	110.38
2	B	7	MAN	O4-C4-C3	-3.58	101.93	110.38
2	B	9	MAN	O4-C4-C3	-3.58	101.95	110.38
2	B	4	MAN	C1-O5-C5	3.53	116.91	112.19
2	B	8	MAN	O5-C1-C2	-3.46	102.54	110.79
2	B	4	MAN	O6-C6-C5	-3.18	100.51	111.33
2	B	2	NAG	C4-C3-C2	-3.03	106.58	111.02
2	B	8	MAN	C1-C2-C3	2.97	113.97	109.64
2	B	2	NAG	O7-C7-C8	-2.95	116.81	122.05
2	B	2	NAG	C8-C7-N2	2.91	120.95	116.12
2	B	1	NAG	C4-C3-C2	-2.81	106.90	111.02
2	B	1	NAG	C1-C2-N2	-2.64	106.28	110.43
2	B	7	MAN	C1-O5-C5	2.55	115.60	112.19
2	B	6	MAN	C1-O5-C5	2.42	115.43	112.19
2	B	9	MAN	C1-O5-C5	2.37	115.36	112.19
2	B	4	MAN	O2-C2-C1	-2.35	103.84	109.22
2	B	2	NAG	O4-C4-C5	-2.10	104.15	109.32
2	B	4	MAN	C2-C3-C4	-2.06	107.24	110.86
2	B	5	MAN	O3-C3-C2	-2.03	105.92	110.05

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	9	MAN	2	0
2	B	7	MAN	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 2 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	SIA	A	2	-	21,21,21	2.13	7 (33%)	24,31,31	1.58	6 (25%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SIA	A	1	-	21,21,21	1.81	4 (19%)	24,31,31	1.45	5 (20%)
3	NAG	A	480(A)	1	14,14,15	2.46	3 (21%)	17,19,21	1.67	3 (17%)
3	NAG	A	478(A)	1	14,14,15	1.12	2 (14%)	17,19,21	2.21	7 (41%)
3	NAG	A	479(B)	-	14,14,15	1.98	3 (21%)	17,19,21	4.44	10 (58%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SIA	A	2	-	-	2/20/38/38	0/1/1/1
4	SIA	A	1	-	-	0/20/38/38	0/1/1/1
3	NAG	A	480(A)	1	-	0/6/23/26	0/1/1/1
3	NAG	A	478(A)	1	-	0/6/23/26	0/1/1/1
3	NAG	A	479(B)	-	-	3/6/23/26	0/1/1/1

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	480(A)	NAG	O7-C7	-7.49	1.06	1.23
4	A	2	SIA	C3-C2	5.94	1.59	1.51
3	A	479(B)	NAG	C4-C5	5.32	1.64	1.53
4	A	1	SIA	C4-C5	5.24	1.57	1.53
4	A	2	SIA	C2-C1	3.95	1.59	1.53
3	A	480(A)	NAG	C1-C2	-3.86	1.47	1.52
3	A	479(B)	NAG	O5-C1	-3.30	1.38	1.43
3	A	479(B)	NAG	C4-C3	3.10	1.60	1.52
4	A	2	SIA	C3-C4	-3.05	1.48	1.53
4	A	2	SIA	O6-C2	2.89	1.46	1.43
4	A	2	SIA	C4-C5	-2.78	1.50	1.53
4	A	1	SIA	O4-C4	-2.57	1.38	1.43
3	A	480(A)	NAG	C4-C3	-2.45	1.46	1.52
4	A	1	SIA	C5-N5	2.26	1.49	1.45
4	A	2	SIA	O2-C2	2.23	1.42	1.39
4	A	1	SIA	C3-C2	2.21	1.54	1.51
4	A	2	SIA	C6-C5	2.13	1.56	1.53
3	A	478(A)	NAG	O5-C5	2.04	1.47	1.43
3	A	478(A)	NAG	C1-C2	-2.00	1.49	1.52

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	479(B)	NAG	C2-N2-C7	13.19	140.57	122.90
3	A	479(B)	NAG	C8-C7-N2	-5.87	106.38	116.12
3	A	479(B)	NAG	O5-C1-C2	-5.25	103.16	111.29
3	A	479(B)	NAG	C4-C3-C2	-5.06	103.60	111.02
3	A	479(B)	NAG	O7-C7-C8	4.29	129.69	122.05
3	A	479(B)	NAG	O3-C3-C2	-3.85	101.40	109.40
3	A	478(A)	NAG	O3-C3-C2	3.72	117.12	109.40
3	A	480(A)	NAG	O7-C7-C8	-3.66	115.53	122.05
3	A	478(A)	NAG	O6-C6-C5	-3.55	99.25	111.33
3	A	478(A)	NAG	C6-C5-C4	-3.47	104.51	113.02
4	A	1	SIA	C3-C2-C1	-3.16	106.97	112.84
3	A	479(B)	NAG	C1-O5-C5	3.07	116.29	112.19
3	A	478(A)	NAG	C3-C4-C5	3.06	115.78	110.23
4	A	2	SIA	O9-C9-C8	-3.05	104.74	111.16
3	A	479(B)	NAG	O3-C3-C4	3.02	117.49	110.38
3	A	480(A)	NAG	O7-C7-N2	3.01	127.31	121.98
4	A	2	SIA	O2-C2-O6	-2.94	102.10	109.51
3	A	480(A)	NAG	C2-N2-C7	-2.92	118.99	122.90
4	A	1	SIA	C6-C5-N5	-2.84	106.38	110.91
3	A	479(B)	NAG	O4-C4-C3	2.60	116.50	110.38
4	A	2	SIA	O2-C2-C3	2.58	113.35	109.44
3	A	478(A)	NAG	C4-C3-C2	-2.54	107.29	111.02
3	A	479(B)	NAG	O5-C5-C6	-2.52	102.75	107.66
3	A	478(A)	NAG	O4-C4-C5	-2.47	103.23	109.32
4	A	2	SIA	C4-C5-N5	-2.47	105.57	110.44
4	A	1	SIA	O10-C10-C11	-2.30	117.96	122.05
3	A	478(A)	NAG	O4-C4-C3	-2.26	105.06	110.38
4	A	2	SIA	O6-C6-C7	-2.12	103.34	106.65
4	A	2	SIA	C3-C2-C1	-2.11	108.92	112.84
4	A	1	SIA	O9-C9-C8	-2.04	106.87	111.16
4	A	1	SIA	O2-C2-C1	-2.04	106.43	110.73

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	479(B)	NAG	C3-C2-N2-C7
3	A	479(B)	NAG	C4-C5-C6-O6
3	A	479(B)	NAG	O5-C5-C6-O6
4	A	2	SIA	C6-C7-C8-O8
4	A	2	SIA	O1B-C1-C2-C3


There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	480(A)	NAG	2	0
3	A	478(A)	NAG	4	0
3	A	479(B)	NAG	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.