



Full wwPDB NMR Structure Validation Report ⓘ

Mar 9, 2026 – 11:47 AM UTC

PDB ID : 2MWP / pdb_00002mwp
BMRB ID : 25348
Title : Solution structure of 53BP1 tandem Tudor domains in complex with a p53K382me2 peptide
Authors : Cui, G.; Botuyan, M.V.; Mer, G.
Deposited on : 2014-11-15

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 91%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	123	
2	B	12	

2 Ensemble composition and analysis

This entry contains 20 models. Model 12 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:1486-A:1495, A:1500-A:1563, A:1568-A:1602 (109)	0.33	12

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 1 single-model cluster was found.

Cluster number	Models
1	2, 9, 10, 11, 12, 13, 14, 17, 19
2	1, 3, 4, 5, 7, 8, 16, 20
3	15, 18
Single-model clusters	6

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2169 atoms, of which 1083 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Tumor suppressor p53-binding protein 1.

Mol	Chain	Residues	Atoms						Trace
1	A	123	Total	C	H	N	O	S	0
			1947	625	965	166	187	4	

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP Q12888
A	-2	HIS	-	expression tag	UNP Q12888
A	-1	MET	-	expression tag	UNP Q12888

- Molecule 2 is a protein called Cellular tumor antigen p53.


Mol	Chain	Residues	Atoms						Trace
2	B	12	Total	C	H	N	O	S	0
			222	66	118	20	17	1	

4 Residue-property plots


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Tumor suppressor p53-binding protein 1



- Molecule 2: Cellular tumor antigen p53




4.2 Scores per residue for each member of the ensemble


Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

- Molecule 1: Tumor suppressor p53-binding protein 1




- Molecule 2: Cellular tumor antigen p53




4.2.2 Score per residue for model 2

- Molecule 1: Tumor suppressor p53-binding protein 1




- Molecule 2: Cellular tumor antigen p53




4.2.3 Score per residue for model 3

- Molecule 1: Tumor suppressor p53-binding protein 1




- Molecule 2: Cellular tumor antigen p53




4.2.4 Score per residue for model 4

- Molecule 1: Tumor suppressor p53-binding protein 1




- Molecule 2: Cellular tumor antigen p53



4.2.5 Score per residue for model 5

- Molecule 1: Tumor suppressor p53-binding protein 1




- Molecule 2: Cellular tumor antigen p53




4.2.6 Score per residue for model 6

- Molecule 1: Tumor suppressor p53-binding protein 1




- Molecule 2: Cellular tumor antigen p53




4.2.7 Score per residue for model 7

- Molecule 1: Tumor suppressor p53-binding protein 1




- Molecule 2: Cellular tumor antigen p53




4.2.8 Score per residue for model 8

- Molecule 1: Tumor suppressor p53-binding protein 1




- Molecule 2: Cellular tumor antigen p53




4.2.9 Score per residue for model 9

- Molecule 1: Tumor suppressor p53-binding protein 1




- Molecule 2: Cellular tumor antigen p53




4.2.10 Score per residue for model 10

- Molecule 1: Tumor suppressor p53-binding protein 1




- Molecule 2: Cellular tumor antigen p53




4.2.11 Score per residue for model 11

- Molecule 1: Tumor suppressor p53-binding protein 1




- Molecule 2: Cellular tumor antigen p53




4.2.12 Score per residue for model 12 (medoid)

- Molecule 1: Tumor suppressor p53-binding protein 1




- Molecule 2: Cellular tumor antigen p53




4.2.13 Score per residue for model 13

- Molecule 1: Tumor suppressor p53-binding protein 1




- Molecule 2: Cellular tumor antigen p53



4.2.14 Score per residue for model 14

- Molecule 1: Tumor suppressor p53-binding protein 1




- Molecule 2: Cellular tumor antigen p53




4.2.15 Score per residue for model 15

- Molecule 1: Tumor suppressor p53-binding protein 1




- Molecule 2: Cellular tumor antigen p53




4.2.16 Score per residue for model 16

- Molecule 1: Tumor suppressor p53-binding protein 1




- Molecule 2: Cellular tumor antigen p53




4.2.17 Score per residue for model 17

- Molecule 1: Tumor suppressor p53-binding protein 1




- Molecule 2: Cellular tumor antigen p53




4.2.18 Score per residue for model 18

- Molecule 1: Tumor suppressor p53-binding protein 1




- Molecule 2: Cellular tumor antigen p53




4.2.19 Score per residue for model 19

- Molecule 1: Tumor suppressor p53-binding protein 1




- Molecule 2: Cellular tumor antigen p53



4.2.20 Score per residue for model 20


- Molecule 1: Tumor suppressor p53-binding protein 1

Chain A: 75% 13% 11%



- Molecule 2: Cellular tumor antigen p53

Chain B: 100%



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 200 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	
Amber	structure solution	
Amber	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1638
Number of shifts mapped to atoms	1638
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	91%

6 Model quality

6.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MLY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.87±0.01	0±0/909 (0.0± 0.0%)	1.39±0.02	4±1/1224 (0.3± 0.1%)
All	All	0.87	0/18180 (0.0%)	1.39	80/24480 (0.3%)

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	1501	PHE	CA-CB-CG	7.14	120.94	113.80	10	7
1	A	1536	ASP	N-CA-C	6.50	124.19	109.81	10	4
1	A	1599	GLN	N-CA-CB	-6.35	102.14	110.59	10	20
1	A	1561	HIS	CB-CG-CD2	-6.31	123.00	131.20	18	20
1	A	1550	ASP	CA-C-N	5.75	132.53	121.54	10	3
1	A	1550	ASP	C-N-CA	5.75	132.53	121.54	10	3
1	A	1537	PRO	N-CA-C	-5.56	101.76	110.50	5	10
1	A	1553	PHE	CA-CB-CG	5.46	119.26	113.80	4	1
1	A	1521	ASP	CA-CB-CG	5.32	117.92	112.60	14	2
1	A	1596	LEU	CA-C-N	5.22	127.53	120.38	19	3
1	A	1596	LEU	C-N-CA	5.22	127.53	120.38	19	3
1	A	1592	GLN	CA-C-N	5.16	125.81	120.03	13	2
1	A	1592	GLN	C-N-CA	5.16	125.81	120.03	13	2

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	889	891	891	2±1
2	B	0	0	0	0±0
All	All	17780	17820	17820	33

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:1540:LEU:HD12	1:A:1540:LEU:H	0.58	1.58	1	20
1:A:1534:LEU:HD23	1:A:1602:LEU:HD22	0.50	1.82	2	4
1:A:1493:ALA:HB2	1:A:1532:ILE:HG23	0.49	1.84	17	7
1:A:1530:LYS:HD2	1:A:1530:LYS:H	0.46	1.71	10	1
1:A:1538:ILE:HB	1:A:1570:TYR:CE2	0.40	2.52	20	1

6.3 Torsion angles

6.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	109/123 (89%)	101±1 (92±1%)	8±2 (7±2%)	1±1 (1±1%)	18	68
2	B	0	-	-	-	-	-
All	All	2180/2700 (81%)	2011 (92%)	151 (7%)	18 (1%)	18	68

All 4 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	1487	VAL	9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	1536	ASP	4
1	A	1486	PHE	3
1	A	1551	GLU	2

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	93/103 (90%)	84±1 (91±1%)	9±1 (9±1%)	10 56
2	B	0	-	-	-
All	All	1860/2280 (82%)	1686 (91%)	174 (9%)	10 56

All 21 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	1505	LYS	20
1	A	1530	LYS	20
1	A	1534	LEU	20
1	A	1542	THR	20
1	A	1532	ILE	19
1	A	1553	PHE	17
1	A	1602	LEU	16
1	A	1589	SER	12
1	A	1492	VAL	8
1	A	1525	CYS	5
1	A	1579	LYS	3
1	A	1547	LEU	3
1	A	1521	ASP	2
1	A	1595	ARG	2
1	A	1510	VAL	1
1	A	1514	LYS	1
1	A	1515	TYR	1
1	A	1574	LYS	1
1	A	1554	SER	1
1	A	1500	TYR	1
1	A	1596	LEU	1

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
2	MLY	B	382	2	9,10,11	0.87±0.06	0±0 (0±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
2	MLY	B	382	2	6,11,13	0.94±0.15	0±0 (0±0%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MLY	B	382	2	-	0±0,8,9,11	-

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 91% for the well-defined parts and 88% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1638
Number of shifts mapped to atoms	1638
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	4

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	132	-0.40 ± 0.07	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	119	0.07 ± 0.14	None needed (< 0.5 ppm)
$^{13}\text{C}'$	105	-0.04 ± 0.21	None needed (< 0.5 ppm)
^{15}N	109	0.21 ± 0.23	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 91%, i.e. 1412 atoms were assigned a chemical shift out of a possible 1544. 0 out of 22 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	538/552 (97%)	224/227 (99%)	210/218 (96%)	104/107 (97%)
Sidechain	749/840 (89%)	511/544 (94%)	234/261 (90%)	4/35 (11%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	125/152 (82%)	65/72 (90%)	60/77 (78%)	0/3 (0%)
Overall	1412/1544 (91%)	800/843 (95%)	504/556 (91%)	108/145 (74%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 88%, i.e. 1620 atoms were assigned a chemical shift out of a possible 1841. 0 out of 23 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	602/681 (88%)	256/281 (91%)	237/268 (88%)	109/132 (83%)
Sidechain	877/984 (89%)	597/636 (94%)	274/306 (90%)	6/42 (14%)
Aromatic	141/176 (80%)	73/85 (86%)	68/86 (79%)	0/5 (0%)
Overall	1620/1841 (88%)	926/1002 (92%)	579/660 (88%)	115/179 (64%)

7.1.4 Statistically unusual chemical shifts [i](#)


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	1542	THR	HG1	5.62	0.08 – 2.19	21.2
1	A	1545	THR	HG1	5.53	0.08 – 2.19	20.8
1	A	1571	SER	HB3	1.51	2.49 – 5.20	-8.6
1	A	1494	LYS	HB2	0.57	0.58 – 2.97	-5.1

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:



Random coil index (RCI) for chain B:

