



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:59 PM UTC

PDB ID : 4MY4 / pdb_00004my4
Title : Crystal structure of phosphoglycerate mutase from Staphylococcus aureus.
Authors : Roychowdhury, A.; Kundu, A.; Gujar, A.; Bose, M.; Das, A.K.
Deposited on : 2013-09-27
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	513	 85% 11% ...

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4327 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 2,3-bisphosphoglycerate-independent phosphoglycerate mutase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	503	Total	C	N	O	S	0	2	0
			3953	2488	669	785	11			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	HIS	-	expression tag	UNP Q2G029
A	-6	HIS	-	expression tag	UNP Q2G029
A	-5	HIS	-	expression tag	UNP Q2G029
A	-4	HIS	-	expression tag	UNP Q2G029
A	-3	HIS	-	expression tag	UNP Q2G029
A	-2	HIS	-	expression tag	UNP Q2G029
A	-1	GLY	-	expression tag	UNP Q2G029
A	0	SER	-	expression tag	UNP Q2G029

- Molecule 2 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Mn	0	0
			2	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	372	Total	O	0	0
			372	372		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 2,3-bisphosphoglycerate-independent phosphoglycerate mutase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	73.21Å 81.75Å 89.18Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.69 – 2.00 19.69 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.2 (19.69-2.00) 99.2 (19.69-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.04 (at 2.01Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.164 , 0.216 0.174 , 0.223	Depositor DCC
R_{free} test set	1818 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	20.8	Xtriage
Anisotropy	0.417	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 37.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4327	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.16	0/4034	1.18	23/5465 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	260	ARG	NE-CZ-NH2	-9.72	110.45	119.20
1	A	260	ARG	CD-NE-CZ	7.58	135.01	124.40
1	A	260	ARG	NE-CZ-NH1	6.72	128.22	121.50
1	A	350	ARG	NE-CZ-NH2	-6.63	113.23	119.20
1	A	185	ARG	NE-CZ-NH2	-6.57	113.29	119.20
1	A	334	VAL	CG1-CB-CG2	6.15	124.32	110.80
1	A	260	ARG	CG-CD-NE	-6.05	98.69	112.00
1	A	42	THR	N-CA-CB	-5.90	100.56	111.53
1	A	78	SER	N-CA-C	5.89	117.70	111.28
1	A	17	ARG	NE-CZ-NH2	-5.83	113.95	119.20
1	A	35	ARG	NE-CZ-NH2	5.67	124.30	119.20
1	A	115	GLY	N-CA-C	5.53	118.65	110.63
1	A	144	VAL	N-CA-C	5.50	115.77	107.80
1	A	350	ARG	NE-CZ-NH1	5.46	126.96	121.50
1	A	42	THR	OG1-CB-CG2	5.30	119.91	109.30
1	A	185	ARG	NE-CZ-NH1	5.28	126.78	121.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	35	ARG	NE-CZ-NH1	-5.23	116.27	121.50
1	A	17	ARG	NE-CZ-NH1	5.21	126.71	121.50
1	A	206	ARG	NE-CZ-NH2	-5.15	114.56	119.20
1	A	481	LEU	CB-CG-CD1	5.15	126.15	110.70
1	A	470	VAL	N-CA-CB	-5.12	102.36	111.92
1	A	257	ARG	CA-C-N	-5.01	113.39	118.85
1	A	257	ARG	C-N-CA	-5.01	113.39	118.85

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	260	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3953	0	3814	45	1
2	A	2	0	0	0	0
3	A	372	0	0	21	4
All	All	4327	0	3814	45	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:193:LYS:H	1:A:263:GLN:HE22	1.12	0.96
1:A:80:THR:HG21	3:A:748:HOH:O	1.68	0.92
1:A:206:ARG:HD2	3:A:865:HOH:O	1.74	0.86
1:A:16:ASN:HD22	1:A:44:GLN:HE21	1.27	0.82
1:A:185:ARG:HD3	3:A:739:HOH:O	1.82	0.80
1:A:316:GLN:HE21	1:A:347:LYS:H	1.28	0.79

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:125:HIS:HD2	1:A:127:LYS:H	1.36	0.72
1:A:59:MET:HE1	3:A:706:HOH:O	1.91	0.69
1:A:221:GLU:HG3	3:A:782:HOH:O	1.92	0.68
1:A:193:LYS:H	1:A:263:GLN:NE2	1.89	0.67
1:A:42:THR:HG23	3:A:803:HOH:O	1.93	0.67
1:A:212:THR:HG21	3:A:1010:HOH:O	1.94	0.66
1:A:212:THR:HG23	3:A:866:HOH:O	1.95	0.66
1:A:474:GLU:HG2	3:A:769:HOH:O	1.95	0.66
1:A:217:LYS:NZ	3:A:862:HOH:O	2.32	0.62
1:A:94:ASN:HD22	1:A:97:LEU:H	1.48	0.60
1:A:59:MET:CE	3:A:706:HOH:O	2.50	0.59
1:A:88:ASP:OD2	1:A:90:ASP:OD2	2.19	0.59
1:A:42:THR:CG2	3:A:803:HOH:O	2.52	0.57
1:A:221:GLU:HB3	3:A:863:HOH:O	2.05	0.57
1:A:206:ARG:HD3	1:A:245:VAL:HB	1.87	0.57
1:A:218:GLU:HB2	3:A:861:HOH:O	2.04	0.56
1:A:59:MET:CE	3:A:911:HOH:O	2.55	0.53
1:A:17:ARG:HD2	3:A:802:HOH:O	2.09	0.52
1:A:59:MET:HE3	3:A:911:HOH:O	2.09	0.52
1:A:212:THR:CG2	3:A:866:HOH:O	2.53	0.51
1:A:316:GLN:HE21	1:A:347:LYS:N	2.04	0.51
1:A:350:ARG:HH22	1:A:385:ASP:CG	2.19	0.51
1:A:31:PRO:O	1:A:35:ARG:HG2	2.11	0.51
1:A:327:GLU:HA	1:A:354:ILE:O	2.11	0.50
1:A:316:GLN:NE2	1:A:347:LYS:H	2.05	0.49
1:A:16:ASN:HD22	1:A:44:GLN:NE2	2.06	0.48
1:A:88:ASP:OD2	1:A:88:ASP:C	2.56	0.47
1:A:83:ASN:OD1	1:A:128:HIS:HE1	1.98	0.47
1:A:409:ILE:O	1:A:413:GLU:HG3	2.16	0.46
1:A:247:ASP:HA	1:A:282:LYS:O	2.17	0.45
1:A:318:ASN:O	1:A:320:LEU:HD13	2.17	0.45
1:A:219:GLY:HA3	1:A:236:PHE:CZ	2.52	0.44
1:A:153:ARG:HD3	3:A:951:HOH:O	2.16	0.43
1:A:441:ASN:C	1:A:441:ASN:OD1	2.61	0.43
1:A:350:ARG:NH2	1:A:385:ASP:OD1	2.53	0.41
1:A:193:LYS:N	1:A:263:GLN:HE22	1.95	0.41
1:A:42:THR:HG21	3:A:822:HOH:O	2.20	0.41
1:A:125:HIS:HE1	3:A:973:HOH:O	2.04	0.41
1:A:324:ARG:HB3	1:A:334:VAL:HG13	2.02	0.40

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:842:HOH:O	3:A:861:HOH:O[2_445]	1.32	0.88
3:A:839:HOH:O	3:A:863:HOH:O[2_445]	1.67	0.53
1:A:308:ASN:ND2	3:A:862:HOH:O[2_445]	1.98	0.22
3:A:841:HOH:O	3:A:862:HOH:O[2_445]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	503/513 (98%)	489 (97%)	12 (2%)	2 (0%)	30 27

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	441	ASN
1	A	439	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	422/433 (98%)	398 (94%)	24 (6%)	18 15

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	18	GLU
1	A	42	THR
1	A	93	GLU
1	A	108	ASP
1	A	137	LYS
1	A	141	VAL
1	A	162	LYS
1	A	185	ARG
1	A	196	GLU
1	A	206	ARG
1	A	212	THR
1	A	225	ASN
1	A	320	LEU
1	A	334	VAL
1	A	388	LEU
1	A	391	LEU
1	A	425	LYS
1	A	428	ASP
1	A	470	VAL
1	A	474	GLU
1	A	481	LEU
1	A	486	LEU
1	A	488	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	16	ASN
1	A	21	HIS
1	A	29	ASN
1	A	32	ASN
1	A	77	GLN
1	A	94	ASN
1	A	103	HIS
1	A	107	HIS
1	A	125	HIS
1	A	128	HIS
1	A	178	GLN
1	A	241	GLN
1	A	263	GLN
1	A	270	ASN
1	A	308	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	316	GLN
1	A	319	ASN
1	A	459	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	503/513 (98%)	-0.53	0 100 100	12, 20, 37, 53	2 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MN	A	601	1/1	1.00	0.01	15,15,15,15	0
2	MN	A	602	1/1	1.00	0.01	10,10,10,10	0

6.5 Other polymers [i](#)

There are no such residues in this entry.