



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:12 PM UTC

PDB ID : 4MYB / pdb_00004myb
Title : Crystal Structure of Francisella tularensis 2-C-methyl-D-erythritol 4-phosphate cytidyltransferase (IspD)
Authors : Noble, S.M.; Tsang, A.K.; Couch, R.D.
Deposited on : 2013-09-27
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	261	26% 66% 16% • 17%

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 1791 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 2-C-methyl-D-erythritol 4-phosphate cytidylyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	217	Total	C	N	O	S	0	0	0
			1738	1114	291	328	5			

There are 33 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	VAL	-	expression tag	UNP Q2A280
A	230	LYS	-	expression tag	UNP Q2A280
A	231	GLY	-	expression tag	UNP Q2A280
A	232	GLU	-	expression tag	UNP Q2A280
A	233	LEU	-	expression tag	UNP Q2A280
A	234	ASN	-	expression tag	UNP Q2A280
A	235	SER	-	expression tag	UNP Q2A280
A	236	LYS	-	expression tag	UNP Q2A280
A	237	LEU	-	expression tag	UNP Q2A280
A	238	GLU	-	expression tag	UNP Q2A280
A	239	GLY	-	expression tag	UNP Q2A280
A	240	LYS	-	expression tag	UNP Q2A280
A	241	PRO	-	expression tag	UNP Q2A280
A	242	ILE	-	expression tag	UNP Q2A280
A	243	PRO	-	expression tag	UNP Q2A280
A	244	ASN	-	expression tag	UNP Q2A280
A	245	PRO	-	expression tag	UNP Q2A280
A	246	LEU	-	expression tag	UNP Q2A280
A	247	LEU	-	expression tag	UNP Q2A280
A	248	GLY	-	expression tag	UNP Q2A280
A	249	LEU	-	expression tag	UNP Q2A280
A	250	ASP	-	expression tag	UNP Q2A280
A	251	SER	-	expression tag	UNP Q2A280
A	252	THR	-	expression tag	UNP Q2A280
A	253	ARG	-	expression tag	UNP Q2A280
A	254	THR	-	expression tag	UNP Q2A280
A	255	GLY	-	expression tag	UNP Q2A280

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	256	HIS	-	expression tag	UNP Q2A280
A	257	HIS	-	expression tag	UNP Q2A280
A	258	HIS	-	expression tag	UNP Q2A280
A	259	HIS	-	expression tag	UNP Q2A280
A	260	HIS	-	expression tag	UNP Q2A280
A	261	HIS	-	expression tag	UNP Q2A280


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	53	Total	O	0	0
			53	53		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 2-C-methyl-D-erythritol 4-phosphate cytidylyltransferase



4 Data and refinement statistics

Property	Value	Source
Space group	F 41 3 2	Depositor
Cell constants a, b, c, α , β , γ	200.41 Å 200.41 Å 200.41 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.56 – 2.40 30.56 – 2.40	Depositor EDS
% Data completeness (in resolution range)	98.0 (30.56-2.40) 97.9 (30.56-2.40)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.14 (at 2.20 Å)	Xtriage
Refinement program	PHENIX 1.8_1069	Depositor
R, R_{free}	0.255 , 0.266 0.258 , 0.264	Depositor DCC
R_{free} test set	903 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	42.1	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 67.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	1791	wwPDB-VP
Average B, all atoms (Å ²)	73.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.58	0/1771	0.96	4/2403 (0.2%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	64	ASN	O-C-N	5.13	128.90	121.97
1	A	45	ASN	CA-C-N	5.12	124.84	119.82
1	A	45	ASN	C-N-CA	5.12	124.84	119.82
1	A	65	SER	N-CA-C	5.10	116.52	111.07

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1738	0	1733	27	0
2	A	53	0	0	1	0
All	All	1791	0	1733	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:119:ASP:OD1	1:A:123:GLN:NE2	2.25	0.69
1:A:92:VAL:O	1:A:92:VAL:HG13	1.97	0.63
1:A:2:SER:OG	2:A:343:HOH:O	2.16	0.61
1:A:64:ASN:HA	1:A:66:LEU:HB3	1.89	0.55
1:A:9:PRO:HD2	1:A:105:HIS:O	2.07	0.54
1:A:92:VAL:O	1:A:92:VAL:CG1	2.56	0.52
1:A:109:ARG:NH2	1:A:163:GLN:OE1	2.40	0.49
1:A:218:LYS:O	1:A:222:GLU:HG2	2.13	0.49
1:A:65:SER:N	1:A:66:LEU:HA	2.27	0.48
1:A:22:PRO:HB2	1:A:24:GLN:HG2	1.96	0.48
1:A:188:LYS:HD3	1:A:188:LYS:O	2.14	0.47
1:A:32:LYS:O	1:A:37:ASN:ND2	2.50	0.45
1:A:133:ILE:HG21	1:A:193:ALA:HB2	1.97	0.45
1:A:224:ALA:O	1:A:227:GLN:HB3	2.16	0.45
1:A:32:LYS:HB3	1:A:37:ASN:OD1	2.17	0.45
1:A:62:TRP:CD1	1:A:75:VAL:HG11	2.52	0.44
1:A:156:ARG:HB2	1:A:157:ASP:HB3	1.99	0.44
1:A:8:ILE:HD12	1:A:39:LEU:HD23	1.99	0.44
1:A:164:THR:HA	1:A:165:PRO:C	2.43	0.44
1:A:129:SER:OG	1:A:202:ASN:O	2.25	0.43
1:A:34:ILE:HD13	1:A:215:ILE:HD12	2.01	0.43
1:A:28:LEU:HA	1:A:28:LEU:HD23	1.73	0.42
1:A:13:ILE:HA	1:A:14:GLY:HA3	1.63	0.42
1:A:135:ALA:HB3	1:A:205:VAL:HG13	2.02	0.41
1:A:102:VAL:O	1:A:167:LEU:HD12	2.20	0.41
1:A:66:LEU:O	1:A:68:TYR:N	2.54	0.41
1:A:27:LYS:HA	1:A:33:THR:HA	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	211/261 (81%)	201 (95%)	10 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	193/230 (84%)	190 (98%)	3 (2%)	55	76

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	92	VAL
1	A	140	GLU
1	A	164	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	30	ASN
1	A	64	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	217/261 (83%)	1.36	69 (31%) 1 1	31, 66, 133, 164	0

All (69) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	226	TRP	9.5
1	A	65	SER	6.7
1	A	36	ASP	6.5
1	A	61	PHE	5.9
1	A	66	LEU	5.8
1	A	33	THR	5.5
1	A	227	GLN	5.4
1	A	157	ASP	5.4
1	A	28	LEU	5.2
1	A	29	ASN	5.1
1	A	158	ASN	5.0
1	A	11	ALA	5.0
1	A	222	GLU	5.0
1	A	26	TYR	4.7
1	A	55	ILE	4.7
1	A	13	ILE	4.6
1	A	228	LEU	4.5
1	A	62	TRP	4.4
1	A	37	ASN	4.3
1	A	20	ASP	4.1
1	A	63	ASN	4.0
1	A	80	GLU	3.9
1	A	144	GLN	3.8
1	A	34	ILE	3.8
1	A	67	TYR	3.7
1	A	217	THR	3.7
1	A	110	PRO	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	25	TYR	3.4
1	A	155	ALA	3.4
1	A	64	ASN	3.4
1	A	221	LEU	3.3
1	A	154	LEU	3.3
1	A	224	ALA	3.3
1	A	216	THR	3.2
1	A	70	HIS	3.2
1	A	68	TYR	3.2
1	A	156	ARG	3.1
1	A	31	GLY	3.1
1	A	218	LYS	3.1
1	A	23	LYS	3.0
1	A	27	LYS	3.0
1	A	225	ASN	2.9
1	A	12	GLY	2.9
1	A	21	ILE	2.8
1	A	213	ILE	2.8
1	A	139	TYR	2.7
1	A	30	ASN	2.7
1	A	40	VAL	2.6
1	A	69	ASN	2.6
1	A	32	LYS	2.6
1	A	136	VAL	2.6
1	A	35	LEU	2.5
1	A	10	ALA	2.5
1	A	108	ALA	2.5
1	A	137	ARG	2.5
1	A	111	CYS	2.4
1	A	223	TYR	2.4
1	A	215	ILE	2.4
1	A	14	GLY	2.3
1	A	77	ASN	2.3
1	A	24	GLN	2.2
1	A	219	ASP	2.2
1	A	107	ALA	2.2
1	A	38	THR	2.2
1	A	209	SER	2.2
1	A	91	ASN	2.1
1	A	44	ASP	2.1
1	A	142	VAL	2.0
1	A	114	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.