



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 12:47 AM UTC

PDB ID : 5MY6 / pdb_00005my6
Title : Crystal structure of a HER2-Nb complex
Authors : Sterckx, Y.G.-J.; D'Huyvetter, M.; Devoogdt, N.
Deposited on : 2017-01-25
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3416 (2.26-2.22)
Clashscore	190562	3556 (2.26-2.22)
Ramachandran outliers	187476	3500 (2.26-2.22)
Sidechain outliers	187428	3501 (2.26-2.22)
RSRZ outliers	180081	3415 (2.26-2.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	622	10% 78% 12% 11%
2	B	115	6% 95% 5%
3	C	2	100%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5581 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Receptor tyrosine-protein kinase erbB-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	556	Total	C	N	O	S	0	5	0
			4282	2658	769	806	49			

- Molecule 2 is a protein called Nanobody 2Rs15d.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	115	Total	C	N	O	S	0	0	0
			875	545	150	174	6			

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	297	Total 297	O 297	0	0
6	B	73	Total 73	O 73	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Receptor tyrosine-protein kinase erbB-2



- Molecule 2: Nanobody 2Rs15d



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	66.52Å 113.52Å 137.43Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.79 – 2.25 47.79 – 2.25	Depositor EDS
% Data completeness (in resolution range)	98.1 (47.79-2.25) 98.2 (47.79-2.25)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.50 (at 2.24Å)	Xtriage
Refinement program	PHENIX 1.10_2155	Depositor
R, R_{free}	0.185 , 0.220 0.192 , 0.226	Depositor DCC
R_{free} test set	2477 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	39.5	Xtriage
Anisotropy	0.120	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 44.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5581	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.21	0/4376	0.44	0/5954
2	B	0.20	0/892	0.45	0/1205
All	All	0.21	0/5268	0.44	0/7159

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4282	0	4059	44	0
2	B	875	0	832	4	0
3	C	28	0	25	0	0
4	A	14	0	13	0	0
5	A	12	0	16	0	0
6	A	297	0	0	2	0
6	B	73	0	0	1	0
All	All	5581	0	4945	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:275:ASN:HD22	1:A:278:THR:H	1.16	0.94
1:A:275:ASN:ND2	1:A:278:THR:H	1.73	0.85
1:A:581:ALA:HB1	1:A:596:CYS:H	1.52	0.74
1:A:168:LEU:H	1:A:213:GLN:HE22	1.37	0.73
2:B:82:MET:HE1	2:B:111:VAL:HG21	1.77	0.67
1:A:57:GLN:OE1	1:A:81:GLN:NE2	2.30	0.64
1:A:106:GLN:NE2	6:A:802:HOH:O	2.31	0.64
1:A:92:ARG:HB3	1:A:138:ARG:HG3	1.79	0.63
1:A:574:VAL:HG12	1:A:575:THR:H	1.64	0.63
1:A:66:PRO:HD2	1:A:69:ALA:HB2	1.80	0.62
1:A:119:ASN:HB2	1:A:158:ASN:HD22	1.68	0.57
1:A:574:VAL:HG12	1:A:575:THR:N	2.19	0.57
1:A:514:LEU:HD21	1:A:542:GLU:HG3	1.87	0.56
1:A:284:ASN:HD22	1:A:285:PRO:HD2	1.71	0.54
1:A:306:THR:HG22	1:A:331:CYS:HB3	1.91	0.52
1:A:160:GLN:NE2	1:A:190:ARG:HD3	2.25	0.52
2:B:19:LYS:HE3	6:B:209:HOH:O	2.11	0.50
1:A:456:ARG:HA	1:A:521:TRP:CD1	2.47	0.50
1:A:502:ASP:HA	1:A:505:VAL:HG12	1.94	0.49
1:A:147:GLU:HB2	1:A:242:ALA:HA	1.95	0.49
1:A:92:ARG:NH2	1:A:133:SER:O	2.46	0.49
1:A:329:GLN:OE1	1:A:434:ARG:NH2	2.48	0.47
1:A:90:GLN:NE2	1:A:121:ASP:OD1	2.46	0.47
1:A:231:LEU:HB2	1:A:233:THR:HG22	1.96	0.46
1:A:160:GLN:HE21	1:A:190:ARG:HD3	1.79	0.46
1:A:365:ALA:HA	1:A:399:VAL:HG22	1.98	0.46
1:A:478:HIS:HD2	6:A:1061:HOH:O	1.99	0.46
1:A:475:CYS:SG	1:A:499:ARG:HD3	2.56	0.46
1:A:87:ALA:HA	1:A:117:LEU:O	2.16	0.45
1:A:550:LEU:HD23	1:A:551:PRO:HA	1.99	0.45
1:A:119:ASN:HD22	1:A:158:ASN:ND2	2.15	0.45
1:A:395:GLU:H	1:A:395:GLU:CD	2.25	0.45
1:A:341:VAL:HG13	1:A:371:PHE:CD1	2.53	0.44
1:A:300:PRO:HG2	1:A:303:TYR:CD1	2.53	0.44
1:A:63:THR:HA	1:A:87:ALA:O	2.17	0.44
1:A:119:ASN:H	1:A:158:ASN:ND2	2.16	0.44
1:A:24:GLN:HG3	1:A:25:VAL:HG23	2.00	0.43
1:A:381:PHE:CZ	1:A:416:PRO:HD3	2.54	0.43
1:A:454:GLY:HA2	1:A:527:GLN:OE1	2.17	0.43
1:A:575:THR:HG22	1:A:576:CYS:SG	2.59	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:553:GLU:OE1	1:A:560:CYS:HB3	2.19	0.43
1:A:157:ARG:NH1	2:B:44:GLU:HB3	2.35	0.42
1:A:90:GLN:HE22	1:A:120:GLY:C	2.28	0.42
2:B:87:PRO:HG3	2:B:115:SER:HB3	2.02	0.41
1:A:552:ARG:C	1:A:553:GLU:HG3	2.45	0.40
1:A:140:LEU:HD23	1:A:140:LEU:HA	1.91	0.40
1:A:217:ARG:HB2	1:A:226:ARG:NH1	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	553/622 (89%)	530 (96%)	23 (4%)	0	100	100
2	B	113/115 (98%)	113 (100%)	0	0	100	100
All	All	666/737 (90%)	643 (96%)	23 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	472/536 (88%)	468 (99%)	4 (1%)	73	80

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	94/96 (98%)	94 (100%)	0	100	100
All	All	566/632 (90%)	562 (99%)	4 (1%)	76	81

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	187	ASN
1	A	249	HIS
1	A	338	CYS
1	A	531	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	75	GLN
1	A	81	GLN
1	A	90	GLN
1	A	158	ASN
1	A	160	GLN
1	A	213	GLN
1	A	239	GLN
1	A	257	HIS
1	A	275	ASN
1	A	284	ASN
1	A	429	GLN
1	A	570	GLN
2	B	71	GLN
2	B	81	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	C	1	3,1	14,14,15	0.41	0	17,19,21	0.41	0
3	NAG	C	2	3	14,14,15	0.23	0	17,19,21	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	3,1	-	4/6/23/26	0/1/1/1
3	NAG	C	2	3	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	2	NAG	O5-C5-C6-O6
3	C	2	NAG	C4-C5-C6-O6
3	C	1	NAG	C4-C5-C6-O6
3	C	1	NAG	O5-C5-C6-O6
3	C	1	NAG	C8-C7-N2-C2
3	C	1	NAG	O7-C7-N2-C2

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	GOL	A	705	-	5,5,5	0.39	0	5,5,5	0.36	0
5	GOL	A	704	-	5,5,5	0.38	0	5,5,5	0.35	0
4	NAG	A	703	-	14,14,15	1.47	2 (14%)	17,19,21	1.04	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	705	-	-	3/4/4/4	-
5	GOL	A	704	-	-	3/4/4/4	-
4	NAG	A	703	-	-	2/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	703	NAG	O5-C1	3.99	1.50	1.43
4	A	703	NAG	C1-C2	3.03	1.56	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	703	NAG	C1-O5-C5	-2.13	109.34	112.19

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	704	GOL	O1-C1-C2-C3
5	A	705	GOL	O1-C1-C2-C3
4	A	703	NAG	O5-C5-C6-O6
5	A	705	GOL	O1-C1-C2-O2
4	A	703	NAG	C4-C5-C6-O6
5	A	704	GOL	O1-C1-C2-O2
5	A	704	GOL	O2-C2-C3-O3
5	A	705	GOL	O2-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	556/622 (89%)	0.22	62 (11%) 10 9	16, 42, 112, 158	5 (0%)
2	B	115/115 (100%)	-0.17	7 (6%) 27 25	26, 36, 70, 107	0
All	All	671/737 (91%)	0.16	69 (10%) 12 11	16, 41, 108, 158	5 (0%)

All (69) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	597	VAL	5.8
1	A	337	PRO	5.7
1	A	132	ALA	4.6
1	A	334	CYS	4.5
1	A	557	ALA	4.2
1	A	317	LEU	4.1
1	A	587	CYS	4.0
1	A	555	VAL	4.0
1	A	595	PHE	3.9
1	A	585	VAL	3.9
1	A	325	GLU	3.8
1	A	122	PRO	3.8
1	A	580	GLU	3.7
1	A	579	PRO	3.7
1	A	577	PHE	3.6
1	A	576	CYS	3.6
1	A	314	VAL	3.5
1	A	572	GLY	3.5
1	A	586	ALA	3.5
2	B	1	GLN	3.5
1	A	338	CYS	3.4
1	A	340	ARG	3.4
1	A	331	CYS	3.4
1	A	550	LEU	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	574	VAL	3.3
1	A	324	ALA	3.2
2	B	42	GLY	3.1
1	A	551	PRO	3.1
2	B	115	SER	3.1
1	A	307	ASP	3.0
1	A	562	PRO	3.0
1	A	141	GLN	2.9
1	A	575	THR	2.9
1	A	327	GLY	2.9
1	A	558	ARG	2.9
1	A	505	VAL	2.9
1	A	540	CYS	2.8
1	A	308	VAL	2.8
1	A	121	ASP	2.8
1	A	565	PRO	2.7
2	B	41	PRO	2.7
1	A	532	SER	2.7
1	A	315	CYS	2.7
1	A	578	GLY	2.7
2	B	44	GLU	2.6
1	A	506	GLY	2.6
1	A	573	SER	2.6
1	A	596	CYS	2.5
1	A	571	ASN	2.5
1	A	568	GLN	2.5
1	A	583	GLN	2.5
1	A	563	CYS	2.4
1	A	502	ASP	2.4
1	A	318	HIS	2.4
1	A	354[A]	ARG	2.3
1	A	564	HIS	2.3
1	A	316	PRO	2.3
1	A	581	ALA	2.3
1	A	333	LYS	2.3
1	A	569	PRO	2.2
1	A	508	GLY	2.2
1	A	554	TYR	2.2
2	B	40	SER	2.2
1	A	319	ASN	2.2
2	B	43	ARG	2.2
1	A	187	ASN	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	547	LEU	2.1
1	A	320	GLN	2.0
1	A	567	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	C	2	14/15	0.75	0.17	73,90,97,97	0
3	NAG	C	1	14/15	0.89	0.12	46,53,75,79	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	A	703	14/15	0.58	0.20	81,98,110,113	0
5	GOL	A	705	6/6	0.82	0.19	81,86,93,104	0
5	GOL	A	704	6/6	0.87	0.17	54,62,69,74	0

6.5 Other polymers [i](#)

There are no such residues in this entry.