



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 09:51 PM UTC

PDB ID : 5MYP / pdb_00005myp
Title : Structure of apo-TbALDH3
Authors : Zoltner, M.; Zhang, N.; Horn, D.; Field, M.C.
Deposited on : 2017-01-27
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	546	
1	B	546	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 8406 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Aldehyde dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	485	Total	C	N	O	S	0	20	0
			3945	2499	696	731	19			
1	B	485	Total	C	N	O	S	0	22	0
			3959	2508	699	733	19			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	expression tag	UNP Q583M9
A	2	SER	-	expression tag	UNP Q583M9
A	3	MET	-	expression tag	UNP Q583M9
A	4	ALA	-	expression tag	UNP Q583M9
A	19	VAL	THR	conflict	UNP Q583M9
A	30	ASP	ASN	conflict	UNP Q583M9
A	235	THR	SER	conflict	UNP Q583M9
A	323	GLN	HIS	conflict	UNP Q583M9
A	354	ASN	ASP	conflict	UNP Q583M9
A	465	LEU	PHE	conflict	UNP Q583M9
A	541	SER	GLY	conflict	UNP Q583M9
B	1	GLY	-	expression tag	UNP Q583M9
B	2	SER	-	expression tag	UNP Q583M9
B	3	MET	-	expression tag	UNP Q583M9
B	4	ALA	-	expression tag	UNP Q583M9
B	19	VAL	THR	conflict	UNP Q583M9
B	30	ASP	ASN	conflict	UNP Q583M9
B	235	THR	SER	conflict	UNP Q583M9
B	323	GLN	HIS	conflict	UNP Q583M9
B	354	ASN	ASP	conflict	UNP Q583M9
B	465	LEU	PHE	conflict	UNP Q583M9
B	541	SER	GLY	conflict	UNP Q583M9

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	252	Total	O	0	0
			252	252		
3	B	238	Total	O	0	0
			238	238		

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	87.72Å 63.95Å 91.75Å 90.00° 98.64° 90.00°	Depositor
Resolution (Å)	22.90 – 1.95 22.90 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.2 (22.90-1.95) 99.2 (22.90-1.95)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.70 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.200 , 0.223 0.207 , 0.225	Depositor DCC
R_{free} test set	3580 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	17.7	Xtriage
Anisotropy	0.380	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 34.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.018 for l,-k,h	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	8406	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.52	0/4014	0.76	0/5429
1	B	0.52	0/4028	0.76	0/5449
All	All	0.52	0/8042	0.76	0/10878

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3945	0	4034	14	0
1	B	3959	0	4049	20	0
2	A	12	0	16	0	0
3	A	252	0	0	0	0
3	B	238	0	0	0	0
All	All	8406	0	8099	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:36:LEU:HD21	1:B:95:VAL:HG21	1.81	0.62
1:B:191[B]:LYS:NZ	1:B:191[B]:LYS:HB3	2.17	0.60
1:A:47:LEU:HB2	1:A:84:VAL:HG11	1.83	0.60
1:B:47:LEU:HB2	1:B:84:VAL:HG11	1.84	0.59
1:A:36:LEU:HD21	1:A:95:VAL:HG21	1.84	0.58
1:A:167:LEU:HD23	1:A:178:ILE:HD11	1.88	0.56
1:B:191[B]:LYS:HB3	1:B:191[B]:LYS:HZ3	1.71	0.56
1:B:167:LEU:HD23	1:B:178:ILE:HD11	1.88	0.54
1:B:191[B]:LYS:NZ	1:B:191[B]:LYS:CB	2.74	0.49
1:B:351:ILE:HD12	1:B:371:LEU:HD23	1.95	0.48
1:A:424:VAL:HG11	1:B:469:ILE:HD12	1.95	0.48
1:A:323[B]:GLN:HA	1:A:323[B]:GLN:OE1	2.12	0.47
1:B:271:MET:HE3	1:B:274:VAL:HG21	1.97	0.47
1:A:460:MET:HE1	1:B:428:ALA:HB2	1.98	0.46
1:B:65[A]:ARG:CZ	1:B:65[A]:ARG:HA	2.46	0.45
1:B:302[A]:LYS:HD3	1:B:302[A]:LYS:HA	1.91	0.43
1:A:260:ILE:HG23	1:A:393:LEU:HD23	2.01	0.43
1:B:260:ILE:HG23	1:B:393:LEU:HD23	2.00	0.43
1:B:352:LEU:CB	1:B:355:ILE:HD11	2.49	0.42
1:A:469:ILE:HD12	1:B:424:VAL:HG11	2.02	0.42
1:A:215[A]:LYS:HD2	1:B:207[A]:GLU:OE1	2.20	0.42
1:A:352:LEU:CB	1:A:355:ILE:HD11	2.50	0.42
1:B:245:GLU:O	1:B:286:MET:HE1	2.19	0.42
1:A:140:GLY:HA3	1:A:449:SER:HB3	2.00	0.42
1:A:65[A]:ARG:CZ	1:A:65[A]:ARG:HA	2.49	0.42
1:A:296[A]:LYS:HA	1:A:296[A]:LYS:HD2	1.84	0.41
1:B:351:ILE:CD1	1:B:371:LEU:HD23	2.51	0.41
1:A:245:GLU:O	1:A:286:MET:HE1	2.21	0.41
1:B:140:GLY:HA3	1:B:449:SER:HB3	2.02	0.41
1:B:65[A]:ARG:HD3	1:B:254:ASN:HD21	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	503/546 (92%)	494 (98%)	7 (1%)	2 (0%)	30	21
1	B	505/546 (92%)	496 (98%)	7 (1%)	2 (0%)	30	21
All	All	1008/1092 (92%)	990 (98%)	14 (1%)	4 (0%)	30	21

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	445	HIS
1	B	445	HIS
1	A	423	VAL
1	B	423	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	433/459 (94%)	429 (99%)	4 (1%)	70	70
1	B	435/459 (95%)	433 (100%)	2 (0%)	81	82
All	All	868/918 (95%)	862 (99%)	6 (1%)	78	76

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	151	LYS
1	A	191[A]	LYS
1	A	191[B]	LYS
1	A	490	LYS
1	B	151	LYS
1	B	254	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	41	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	48	ASN
1	A	455	HIS
1	B	62	HIS
1	B	136	GLN
1	B	196	HIS
1	B	254	ASN
1	B	359	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	A	601	-	5,5,5	0.31	0	5,5,5	0.26	0
2	GOL	A	602	-	5,5,5	0.41	0	5,5,5	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	601	-	-	0/4/4/4	-
2	GOL	A	602	-	-	3/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	602	GOL	C1-C2-C3-O3
2	A	602	GOL	O2-C2-C3-O3
2	A	602	GOL	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	485/546 (88%)	0.23	11 (2%) 61 68	6, 18, 30, 53	20 (4%)
1	B	485/546 (88%)	0.31	29 (5%) 27 32	6, 17, 33, 58	22 (4%)
All	All	970/1092 (88%)	0.27	40 (4%) 41 48	6, 17, 31, 58	42 (4%)

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	469	ILE	6.0
1	B	6	ALA	6.0
1	B	468	SER	4.8
1	A	6	ALA	4.4
1	B	490	LYS	3.9
1	B	489	LEU	3.6
1	A	466	PHE	3.5
1	B	466	PHE	3.3
1	B	484	VAL	3.3
1	B	300[A]	LEU	3.2
1	A	305[A]	GLN	3.2
1	A	469	ILE	3.2
1	B	289	ASP	3.2
1	B	471	THR	3.1
1	A	490	LYS	3.1
1	B	487	SER	2.9
1	A	289	ASP	2.8
1	B	488	LEU	2.7
1	A	489	LEU	2.7
1	A	10	GLU	2.6
1	B	465	LEU	2.6
1	B	302[A]	LYS	2.5
1	B	406	ARG	2.4
1	B	85	TRP	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	486	ASN	2.3
1	B	89	GLU	2.3
1	A	465	LEU	2.3
1	B	485	LEU	2.3
1	B	479	THR	2.3
1	B	472	VAL	2.2
1	B	297	GLY	2.2
1	A	107	ALA	2.2
1	B	483[A]	ARG	2.2
1	B	470	ASP	2.2
1	B	355	ILE	2.1
1	B	403	ARG	2.1
1	A	355	ILE	2.1
1	B	356	ASP	2.1
1	B	108	LEU	2.1
1	B	65[A]	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GOL	A	602	6/6	0.65	0.18	43,43,44,44	0
2	GOL	A	601	6/6	0.84	0.13	24,25,26,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.