



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 17, 2026 – 08:39 PM UTC

PDB ID : 6MYS / pdb_00006mys
Title : Avian mitochondrial complex II with Atpenin A5 bound, sidechain outside
Authors : Berry, E.A.; Huang, L.-S.
Deposited on : 2018-11-02
Resolution : 2.37 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.37 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7164 (2.40-2.36)
Clashscore	190562	7722 (2.40-2.36)
Ramachandran outliers	187476	7626 (2.40-2.36)
Sidechain outliers	187428	7627 (2.40-2.36)
RSRZ outliers	180081	7170 (2.40-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	621	3% 87% 12% .
2	B	252	2% 85% 10% 5% .
3	C	140	4% 91% 8% .
4	D	103	4% 95% . .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	OAA	A	1003	-	-	X	-
8	UNL	B	1006	-	-	-	X

2 Entry composition

There are 16 unique types of molecules in this entry. The entry contains 9014 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Succinate dehydrogenase [ubiquinone] flavoprotein subunit, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	612	Total	C	N	O	S	0	2	0
			4737	2963	848	897	29			

- Molecule 2 is a protein called Succinate dehydrogenase [ubiquinone] iron-sulfur subunit, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	240	Total	C	N	O	S	0	0	0
			1928	1219	326	361	22			

- Molecule 3 is a protein called Succinate dehydrogenase cytochrome b, large subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	139	Total	C	N	O	S	0	0	0
			1076	707	178	186	5			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	139	ALA	SER	conflict	UNP D0VWW3
C	140	MET	GLU	conflict	UNP D0VWW3

- Molecule 4 is a protein called Succinate dehydrogenase [ubiquinone] cytochrome b small subunit, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	102	Total	C	N	O	S	0	0	0
			771	508	122	138	3			

- Molecule 5 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 6 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	K	0	0
			1	1		

- Molecule 7 is OXALOACETATE ION (CCD ID: OAA) (formula: C₄H₃O₅).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			9	4	5		

- Molecule 8 is UNKNOWN LIGAND (CCD ID: UNL) (formula:).


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
8	A	14	Total	C	N	O	S	0	0
			48	19	7	19	3		
8	B	6	Total	C	N	O	S	0	0
			27	12	5	8	2		
8	C	6	Total	C	O	S		0	0
			23	13	9	1			
8	D	14	Total	C	O			0	0
			65	37	28				

- Molecule 9 is FE2/S2 (INORGANIC) CLUSTER (CCD ID: FES) (formula: Fe₂S₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	B	1	Total	Fe	S	0	0
			4	2	2		

- Molecule 10 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe₄S₄).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	B	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 11 is FE3-S4 CLUSTER (CCD ID: F3S) (formula: Fe_3S_4).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
11	B	1	Total	Fe	S	0	0
			7	3	4		

- Molecule 12 is 3-[(2S,4S,5R)-5,6-DICHLORO-2,4-DIMETHYL-1-OXOHEXYL]-4-HYDROXY-5,6-DIMETHOXY-2(1H)-PYRIDINONE (CCD ID: AT5) (formula: $\text{C}_{15}\text{H}_{21}\text{Cl}_2\text{NO}_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
14	C	1	Total	C	O	0	0
			34	23	11		

- Molecule 15 is 1,2-Distearoyl-sn-glycerophosphoethanolamine (CCD ID: 3PE) (formula: $C_{41}H_{82}NO_8P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
15	D	1	Total	C	N	O	P	0	0
			36	26	1	8	1		


- Molecule 16 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
16	A	45	Total 45	O 45	0	0
16	B	52	Total 52	O 52	0	0
16	C	17	Total 17	O 17	0	0
16	D	9	Total 9	O 9	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Succinate dehydrogenase [ubiquinone] flavoprotein subunit, mitochondrial



- Molecule 2: Succinate dehydrogenase [ubiquinone] iron-sulfur subunit, mitochondrial




- Molecule 3: Succinate dehydrogenase cytochrome b, large subunit



- Molecule 4: Succinate dehydrogenase [ubiquinone] cytochrome b small subunit, mitochondrial





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	71.68Å 84.42Å 293.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.75 – 2.37 38.75 – 2.37	Depositor EDS
% Data completeness (in resolution range)	99.3 (38.75-2.37) 90.4 (38.75-2.37)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.80 (at 2.37Å)	Xtriage
Refinement program	PHENIX (DEV_3150: ???)	Depositor
R, R_{free}	0.211 , 0.237 0.213 , 0.238	Depositor DCC
R_{free} test set	2144 reflections (2.92%)	wwPDB-VP
Wilson B-factor (Å ²)	44.6	Xtriage
Anisotropy	0.697	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 44.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9014	wwPDB-VP
Average B, all atoms (Å ²)	74.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SF4, AT5, K, HEM, OAA, UMQ, FES, MLZ, UNL, FAD, 3PE, F3S

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.09	0/4838	0.29	0/6549
2	B	0.11	0/1970	0.31	0/2656
3	C	0.09	0/1094	0.25	0/1486
4	D	0.08	0/794	0.20	0/1089
All	All	0.10	0/8696	0.28	0/11780

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4737	0	4619	46	0
2	B	1928	0	1921	14	0
3	C	1076	0	1116	5	0
4	D	771	0	763	3	0
5	A	53	0	30	5	0
6	A	1	0	0	0	0
7	A	9	0	2	6	0
8	A	48	0	0	2	0
8	B	27	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	C	23	0	0	0	0
8	D	65	0	0	0	0
9	B	4	0	0	0	0
10	B	8	0	0	0	0
11	B	7	0	0	0	0
12	C	23	0	20	1	0
13	C	41	0	24	0	0
14	C	34	0	44	0	0
15	D	36	0	46	2	0
16	A	45	0	0	0	0
16	B	52	0	0	1	0
16	C	17	0	0	0	0
16	D	9	0	0	0	0
All	All	9014	0	8585	69	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:297:ARG:HH22	7:A:1003:OAA:C2	1.69	1.05
1:A:297:ARG:HH22	7:A:1003:OAA:H22	1.27	0.98
1:A:297:ARG:NH2	7:A:1003:OAA:H22	1.89	0.86
1:A:297:ARG:HH22	7:A:1003:OAA:C1	1.97	0.76
1:A:60:ALA:HB3	1:A:154:GLY:HA3	1.70	0.73
3:C:6:LYS:HA	3:C:9:MET:HE2	1.70	0.72
1:A:401:ALA:N	1:A:402:SER:HA	2.15	0.62
1:A:422[A]:ARG:NH1	8:A:1014:UNL:O	2.32	0.62
3:C:88:LYS:NZ	3:C:140:MET:O	2.35	0.57
3:C:92:VAL:HG21	3:C:137:ILE:HG21	1.86	0.57
1:A:117:PHE:HA	1:A:149:VAL:HG22	1.86	0.57
4:D:44:THR:HG21	4:D:80:PHE:HB2	1.88	0.56
2:B:10:ARG:NH2	16:B:1104:HOH:O	2.38	0.55
1:A:63:GLY:HA2	1:A:153:THR:HG21	1.87	0.55
2:B:23:LYS:HB2	2:B:26:ASP:HB2	1.90	0.54
2:B:210:PHE:HA	2:B:214:ARG:HG2	1.89	0.54
1:A:253:HIS:NE2	7:A:1003:OAA:O5	2.37	0.54
1:A:304:MET:HE2	1:A:361:PRO:HD3	1.89	0.54
3:C:107:VAL:HA	3:C:110:MET:HE2	1.90	0.54
1:A:13:PRO:HG2	1:A:200:THR:HG22	1.90	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:8:THR:O	2:B:38:ASN:ND2	2.42	0.53
2:B:18:ARG:NH1	2:B:33:TYR:OH	2.41	0.53
1:A:310:GLU:OE1	1:A:312:ARG:NH2	2.43	0.52
1:A:401:ALA:HB3	1:A:403:VAL:N	2.26	0.51
4:D:97:VAL:HG21	15:D:201:3PE:H252	1.92	0.50
1:A:50:LEU:HD21	1:A:228:THR:HG21	1.93	0.50
1:A:57:THR:OG1	5:A:1001:FAD:O2A	2.23	0.50
1:A:184:GLU:HB3	1:A:189:ARG:HE	1.77	0.50
1:A:243:PRO:HB3	1:A:586:TYR:CZ	2.46	0.50
1:A:189:ARG:HD2	1:A:439:PRO:HB2	1.93	0.50
1:A:401:ALA:HB3	1:A:403:VAL:H	1.75	0.49
1:A:25:GLY:HA2	5:A:1001:FAD:H1B	1.94	0.49
1:A:553:ASP:OD1	1:A:553:ASP:N	2.42	0.49
1:A:239:ARG:HG2	1:A:589:VAL:HG21	1.96	0.47
8:A:1017:UNL:O	2:B:64:SER:OG	2.29	0.47
1:A:235:ALA:HB1	1:A:530:LEU:HD12	1.95	0.47
1:A:297:ARG:NH2	7:A:1003:OAA:C2	2.51	0.47
1:A:87:SER:HB2	1:A:405:GLY:HA3	1.97	0.47
1:A:460:ASP:OD1	1:A:507:LYS:NZ	2.32	0.46
1:A:60:ALA:HA	5:A:1001:FAD:C6	2.46	0.46
1:A:32:ARG:NH2	1:A:109:GLU:OE2	2.45	0.45
1:A:99:MET:HA	1:A:419:VAL:HG11	1.98	0.45
1:A:397:GLU:OE1	5:A:1001:FAD:O3'	2.29	0.45
2:B:103:PRO:HD2	2:B:166:THR:HB	1.99	0.45
1:A:268:CYS:HB3	1:A:325:LEU:HD21	1.99	0.44
2:B:12:LYS:NZ	2:B:95:LYS:O	2.44	0.44
2:B:164:CYS:HB2	2:B:221:CYS:HB2	1.99	0.44
2:B:81:ASN:O	2:B:224:THR:OG1	2.35	0.44
5:A:1001:FAD:H9	5:A:1001:FAD:H1'1	1.79	0.43
1:A:129:ALA:HB2	1:A:144:HIS:CD2	2.53	0.43
1:A:116:PRO:HD2	1:A:156:SER:HB3	2.01	0.43
1:A:597:LYS:HG2	1:A:598:TYR:H	1.84	0.43
1:A:489:LEU:HD13	1:A:540:GLU:HA	2.01	0.42
12:C:202:AT5:H8	12:C:202:AT5:O1	2.19	0.42
1:A:216:TYR:HB3	1:A:231:GLY:HA3	2.02	0.42
2:B:65:CYS:O	2:B:66:ARG:HG3	2.20	0.42
1:A:287:TYR:OH	1:A:307:GLU:OE2	2.32	0.41
2:B:176:ASP:HB3	3:C:13:TRP:CZ2	2.56	0.41
1:A:230:THR:HG21	1:A:524:GLU:HB3	2.03	0.41
1:A:271:GLU:HB3	1:A:332:GLN:NE2	2.36	0.41
1:A:103:ALA:HA	1:A:415:LEU:HD11	2.03	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:D:101:TRP:CH2	15:D:201:3PE:H352	2.56	0.41
1:A:51:PHE:HB3	1:A:54:ARG:HG2	2.02	0.41
1:A:456:LEU:HD13	1:A:522:THR:HG22	2.02	0.41
1:A:220:TYR:CG	1:A:363:VAL:HG21	2.55	0.40
1:A:253:HIS:HB2	1:A:364:HIS:HB2	2.02	0.40
2:B:178:TYR:CD1	2:B:214:ARG:HB2	2.56	0.40
1:A:251:GLN:HB2	1:A:367:MET:SD	2.62	0.40
2:B:52:LYS:HA	2:B:56:ASP:O	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	612/621 (99%)	584 (95%)	28 (5%)	0	100	100
2	B	238/252 (94%)	229 (96%)	7 (3%)	2 (1%)	16	23
3	C	136/140 (97%)	135 (99%)	1 (1%)	0	100	100
4	D	100/103 (97%)	99 (99%)	1 (1%)	0	100	100
All	All	1086/1116 (97%)	1047 (96%)	37 (3%)	2 (0%)	43	56

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	64	SER
2	B	66	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	497/506 (98%)	494 (99%)	3 (1%)	78	88
2	B	216/219 (99%)	214 (99%)	2 (1%)	70	84
3	C	116/117 (99%)	115 (99%)	1 (1%)	70	84
4	D	78/79 (99%)	78 (100%)	0	100	100
All	All	907/921 (98%)	901 (99%)	6 (1%)	76	87

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	72	GLU
1	A	130	PHE
1	A	191	VAL
2	B	34	GLU
2	B	217	THR
3	C	91	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	144	HIS
1	A	320	HIS
2	B	144	GLN
3	C	69	HIS
3	C	105	HIS
3	C	118	GLN
4	D	69	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	MLZ	C	15	3	8,9,10	0.65	0	4,9,11	1.54	1 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MLZ	C	15	3	-	1/7/8/10	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	15	MLZ	CM-NZ-CE	2.89	120.06	112.01

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	15	MLZ	C-CA-CB-CG

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 50 ligands modelled in this entry, 1 is monoatomic and 40 are unknown - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
13	HEM	C	201	4,3	48,48,50	1.39	7 (14%)	66,80,82	1.15	5 (7%)
15	3PE	D	201	-	35,35,50	1.09	2 (5%)	38,40,55	1.12	3 (7%)
10	SF4	B	1002	2	0,12,12	-	-	-	-	-
12	AT5	C	202	-	20,23,23	1.34	3 (15%)	21,32,32	1.56	4 (19%)
14	UMQ	C	203	-	35,35,35	0.41	0	46,46,46	0.70	0
11	F3S	B	1003	2	0,9,9	-	-	-	-	-
9	FES	B	1001	2	0,4,4	-	-	-	-	-
5	FAD	A	1001	-	58,58,58	1.08	6 (10%)	85,89,89	1.46	13 (15%)
7	OAA	A	1003	-	8,8,8	2.55	3 (37%)	8,10,10	1.80	3 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
13	HEM	C	201	4,3	-	1/10/50/54	-
15	3PE	D	201	-	-	4/39/39/54	-
12	AT5	C	202	-	-	9/20/22/22	0/1/1/1
14	UMQ	C	203	-	-	5/20/60/60	0/2/2/2
10	SF4	B	1002	2	-	-	0/6/5/5
11	F3S	B	1003	2	-	-	0/3/3/3
9	FES	B	1001	2	-	-	0/1/1/1
5	FAD	A	1001	-	-	4/34/50/50	0/6/6/6
7	OAA	A	1003	-	-	3/8/8/8	-

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	1003	OAA	O3-C3	5.07	1.33	1.23
15	D	201	3PE	O31-C31	4.19	1.45	1.33
15	D	201	3PE	O21-C21	4.14	1.46	1.34
13	C	201	HEM	FE-NB	3.60	2.06	1.94
7	A	1003	OAA	C2-C3	-3.58	1.37	1.51
13	C	201	HEM	FE-ND	3.54	2.05	1.94
12	C	202	AT5	C6-C1	3.10	1.49	1.40
13	C	201	HEM	FE-NC	3.00	2.05	1.95
5	A	1001	FAD	C5A-C6A	2.95	1.49	1.41
13	C	201	HEM	FE-NA	2.91	2.04	1.95
5	A	1001	FAD	C4-N3	-2.65	1.33	1.38
12	C	202	AT5	C6-C5	2.53	1.49	1.44
5	A	1001	FAD	P-O3P	2.25	1.61	1.59
12	C	202	AT5	C5-N4	-2.23	1.34	1.38
5	A	1001	FAD	C8A-N7A	2.20	1.35	1.31
7	A	1003	OAA	C2-C1	2.14	1.54	1.51
5	A	1001	FAD	PA-O3P	2.06	1.61	1.59
13	C	201	HEM	CMB-C2B	2.05	1.55	1.50
13	C	201	HEM	CMA-C3A	2.05	1.55	1.50
13	C	201	HEM	CAC-C3C	2.04	1.55	1.50
5	A	1001	FAD	C5A-N7A	-2.04	1.35	1.39

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
12	C	202	AT5	C1-C6-C7	4.98	123.67	118.11
5	A	1001	FAD	C5A-C4A-N3A	-4.95	119.90	126.72
5	A	1001	FAD	C4-N3-C2	-4.16	118.26	125.64
15	D	201	3PE	O21-C21-C22	3.76	119.62	111.48
5	A	1001	FAD	C4X-C4-N3	3.38	121.85	113.25
5	A	1001	FAD	N3A-C4A-N9A	3.33	132.83	127.17
5	A	1001	FAD	N9A-C8A-N7A	-3.33	109.22	113.94
5	A	1001	FAD	C4A-N9A-C8A	2.81	108.69	105.74
5	A	1001	FAD	C4'-C3'-C2'	-2.79	108.93	113.57
5	A	1001	FAD	C2A-N3A-C4A	2.74	118.53	111.83
13	C	201	HEM	C1B-NB-C4B	2.65	108.35	105.21
5	A	1001	FAD	O4-C4-C4X	-2.65	119.54	126.53
7	A	1003	OAA	O3-C3-C2	2.54	124.32	120.41
13	C	201	HEM	C4D-ND-C1D	2.53	108.21	105.21
5	A	1001	FAD	C4A-C5A-N7A	-2.53	107.69	110.58
12	C	202	AT5	C6-C5-N4	2.52	122.05	115.10
15	D	201	3PE	O31-C31-C32	2.49	119.42	111.83
7	A	1003	OAA	O2-C1-C2	2.45	122.09	114.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	1001	FAD	C5A-N7A-C8A	2.34	107.13	103.45
15	D	201	3PE	C2-O21-C21	-2.32	112.25	117.80
13	C	201	HEM	C3D-C4D-ND	-2.20	107.76	110.17
13	C	201	HEM	C2A-C1A-NA	-2.15	107.77	110.15
5	A	1001	FAD	C4X-C10-N1	-2.12	119.40	124.59
7	A	1003	OAA	O4-C4-C3	-2.08	119.22	121.81
12	C	202	AT5	C12-C13-CL12	-2.08	107.68	109.91
5	A	1001	FAD	C4-C4X-N5	2.07	121.07	118.21
12	C	202	AT5	C8-C11-C12	-2.06	111.40	115.09
13	C	201	HEM	C3B-C4B-NB	-2.05	107.54	109.90

There are no chirality outliers.

All (26) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
5	A	1001	FAD	N10-C1'-C2'-O2'
5	A	1001	FAD	N10-C1'-C2'-C3'
7	A	1003	OAA	C2-C3-C4-O4
7	A	1003	OAA	C2-C3-C4-O5
12	C	202	AT5	C11-C12-C13-CL12
12	C	202	AT5	C17-C12-C13-CL12
12	C	202	AT5	C12-C13-C14-CL16
15	D	201	3PE	C22-C21-O21-C2
15	D	201	3PE	O22-C21-O21-C2
12	C	202	AT5	C5-C6-C7-O5
12	C	202	AT5	C1-C6-C7-C8
14	C	203	UMQ	CI-CJ-CK-CL
12	C	202	AT5	C5-C6-C7-C8
12	C	202	AT5	C17-C12-C13-C14
5	A	1001	FAD	PA-O3P-P-O5'
14	C	203	UMQ	O5-C5-C6-O6
7	A	1003	OAA	O3-C3-C4-O4
12	C	202	AT5	C11-C12-C13-C14
12	C	202	AT5	C1-C6-C7-O5
13	C	201	HEM	C2A-CAA-CBA-CGA
5	A	1001	FAD	P-O3P-PA-O2A
14	C	203	UMQ	CA-CB-CC-CD
15	D	201	3PE	C22-C23-C24-C25
14	C	203	UMQ	C5'-C4'-O1-C1
14	C	203	UMQ	C3'-C4'-O1-C1
15	D	201	3PE	C25-C26-C27-C28



There are no ring outliers.


4 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
15	D	201	3PE	2	0
12	C	202	AT5	1	0
5	A	1001	FAD	5	0
7	A	1003	OAA	6	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	612/621 (98%)	0.66	17 (2%) 55 54	28, 74, 110, 132	2 (0%)
2	B	240/252 (95%)	0.38	4 (1%) 69 68	45, 59, 107, 121	0
3	C	138/140 (98%)	0.30	5 (3%) 46 46	51, 69, 102, 119	0
4	D	102/103 (99%)	0.39	4 (3%) 43 43	51, 68, 92, 105	0
All	All	1092/1116 (97%)	0.53	30 (2%) 56 56	28, 71, 107, 132	2 (0%)

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	618	ILE	4.2
1	A	10	THR	4.0
4	D	2	SER	3.8
3	C	2	ALA	3.7
1	A	387	LYS	3.4
4	D	73	TYR	3.4
1	A	439	PRO	2.8
1	A	382	VAL	2.7
3	C	67	PHE	2.7
2	B	59	LEU	2.7
2	B	8	THR	2.6
1	A	587	VAL	2.6
1	A	443	PRO	2.5
4	D	89	TYR	2.5
1	A	89	TRP	2.4
1	A	331	GLN	2.3
3	C	79	LEU	2.2
1	A	575	PHE	2.2
1	A	385	GLU	2.2
4	D	85	TYR	2.2
1	A	619	ARG	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	55	LEU	2.2
1	A	558	ILE	2.1
2	B	133	ASP	2.1
1	A	561	PHE	2.1
3	C	76	SER	2.0
1	A	314	CYS	2.0
3	C	83	LEU	2.0
1	A	211	ILE	2.0
1	A	258	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MLZ	C	15	10/11	0.89	0.12	66,72,84,85	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
8	UNL	D	210	1/-	0.41	0.18	87,87,87,87	0
8	UNL	D	214	1/-	0.64	0.23	75,75,75,75	0
8	UNL	A	1015	1/-	0.71	0.23	106,106,106,106	0
8	UNL	B	1006	3/-	0.73	0.41	87,87,88,90	0
8	UNL	D	215	1/-	0.73	0.19	76,76,76,76	0
8	UNL	D	207	18/-	0.74	0.28	111,129,134,134	0
8	UNL	A	1011	3/-	0.74	0.21	82,82,84,84	0
8	UNL	B	1004	15/-	0.76	0.28	76,87,123,129	15

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
8	UNL	D	211	1/-	0.77	0.16	80,80,80,80	0
8	UNL	C	207	3/-	0.78	0.29	69,69,78,81	0
8	UNL	D	204	3/-	0.79	0.31	70,70,71,73	0
8	UNL	A	1005	15/-	0.79	0.30	111,119,161,162	0
8	UNL	C	204	12/-	0.80	0.21	87,109,114,114	0
8	UNL	A	1006	4/-	0.80	0.14	112,116,117,121	0
8	UNL	D	202	9/-	0.80	0.14	96,99,101,103	0
8	UNL	A	1014	1/-	0.82	0.19	65,65,65,65	0
15	3PE	D	201	36/51	0.82	0.21	77,106,170,176	0
8	UNL	D	213	1/-	0.83	0.16	72,72,72,72	0
8	UNL	D	208	9/-	0.84	0.20	77,98,112,113	0
8	UNL	D	205	2/-	0.86	0.20	54,54,54,57	0
8	UNL	A	1012	1/-	0.87	0.16	91,91,91,91	0
8	UNL	A	1007	4/-	0.87	0.13	83,87,88,91	0
8	UNL	B	1007	3/-	0.88	0.19	57,57,64,68	0
8	UNL	A	1009	4/-	0.88	0.17	80,83,86,97	0
8	UNL	A	1010	1/-	0.88	0.19	60,60,60,60	0
8	UNL	D	206	5/-	0.88	0.16	83,89,97,107	0
8	UNL	A	1008	4/-	0.89	0.15	78,79,79,85	0
8	UNL	C	206	4/-	0.89	0.13	87,88,89,92	0
14	UMQ	C	203	34/34	0.89	0.15	59,85,115,119	0
8	UNL	B	1005	4/-	0.89	0.16	57,61,71,85	0
8	UNL	C	209	1/-	0.90	0.24	86,86,86,86	0
8	UNL	D	209	10/-	0.90	0.16	81,90,98,99	0
8	UNL	C	205	2/-	0.90	0.25	83,83,83,85	0
8	UNL	A	1017	1/-	0.91	0.18	53,53,53,53	0
8	UNL	A	1004	7/-	0.91	0.24	97,100,106,108	0
8	UNL	A	1016	1/-	0.91	0.12	62,62,62,62	0
12	AT5	C	202	23/23	0.92	0.13	49,57,109,121	0
8	UNL	A	1013	1/-	0.93	0.12	61,61,61,61	0
8	UNL	C	208	1/-	0.93	0.22	91,91,91,91	0
8	UNL	D	212	1/-	0.93	0.17	71,71,71,71	0
8	UNL	B	1008	1/-	0.94	0.09	62,62,62,62	0
8	UNL	B	1009	1/-	0.95	0.29	68,68,68,68	0
7	OAA	A	1003	9/9	0.95	0.08	55,63,71,77	0
8	UNL	D	203	3/-	0.95	0.12	62,62,64,65	0
5	FAD	A	1001	53/53	0.97	0.07	48,58,63,66	0
6	K	A	1002	1/1	0.97	0.07	61,61,61,61	1
9	FES	B	1001	4/4	0.98	0.04	50,53,58,63	0
13	HEM	C	201	41/43	0.98	0.08	54,59,66,75	0
10	SF4	B	1002	8/8	0.99	0.04	51,53,55,58	0
11	F3S	B	1003	7/7	0.99	0.06	46,50,56,58	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around 3PE D 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around UMQ C 203:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around AT5 C 202:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around FAD A 1001:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM C 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.