



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 12:48 AM UTC

PDB ID : 4MZD / pdb_00004mzd
Title : High resolution crystal structure of the nisin leader peptidase NisP from *Lactococcus lactis*
Authors : Rao, Z.H.; Xu, Y.Y.; Li, X.; Yang, W.
Deposited on : 2013-09-30
Resolution : 1.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1625 (1.12-1.08)
Clashscore	190562	1656 (1.12-1.08)
Ramachandran outliers	187476	1616 (1.12-1.08)
Sidechain outliers	187428	1612 (1.12-1.08)
RSRZ outliers	180081	1625 (1.12-1.08)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	488	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3296 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nisin leader peptide-processing serine protease NisP.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	356	Total	C	N	O	S	0	11	0
			2842	1792	485	557	8			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	160	GLY	-	expression tag	UNP D9IXC0
A	161	SER	-	expression tag	UNP D9IXC0
A	162	GLY	-	expression tag	UNP D9IXC0
A	163	MET	-	expression tag	UNP D9IXC0
A	164	LYS	-	expression tag	UNP D9IXC0
A	165	GLU	-	expression tag	UNP D9IXC0
A	166	THR	-	expression tag	UNP D9IXC0
A	167	ALA	-	expression tag	UNP D9IXC0
A	168	ALA	-	expression tag	UNP D9IXC0
A	169	ALA	-	expression tag	UNP D9IXC0
A	170	LYS	-	expression tag	UNP D9IXC0
A	171	PHE	-	expression tag	UNP D9IXC0
A	172	GLU	-	expression tag	UNP D9IXC0
A	173	ARG	-	expression tag	UNP D9IXC0
A	174	GLN	-	expression tag	UNP D9IXC0
A	175	HIS	-	expression tag	UNP D9IXC0
A	176	MET	-	expression tag	UNP D9IXC0
A	177	ASP	-	expression tag	UNP D9IXC0
A	178	SER	-	expression tag	UNP D9IXC0
A	179	PRO	-	expression tag	UNP D9IXC0
A	180	ASP	-	expression tag	UNP D9IXC0
A	181	LEU	-	expression tag	UNP D9IXC0
A	182	GLY	-	expression tag	UNP D9IXC0
A	183	THR	-	expression tag	UNP D9IXC0
A	184	ASP	-	expression tag	UNP D9IXC0
A	185	ASP	-	expression tag	UNP D9IXC0
A	186	ASP	-	expression tag	UNP D9IXC0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	187	ASP	-	expression tag	UNP D9IXC0
A	188	LYS	-	expression tag	UNP D9IXC0
A	189	ALA	-	expression tag	UNP D9IXC0
A	190	MET	-	expression tag	UNP D9IXC0
A	191	ALA	-	expression tag	UNP D9IXC0
A	192	ASP	-	expression tag	UNP D9IXC0
A	193	ILE	-	expression tag	UNP D9IXC0
A	194	GLY	-	expression tag	UNP D9IXC0
A	195	SER	-	expression tag	UNP D9IXC0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	454	Total O 454 454	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nisin leader peptide-processing serine protease NisP



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	49.89Å 44.97Å 74.65Å 90.00° 106.82° 90.00°	Depositor
Resolution (Å)	26.95 – 1.10 26.95 – 1.10	Depositor EDS
% Data completeness (in resolution range)	(Not available) (26.95-1.10) 97.7 (26.95-1.10)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.54 (at 1.10Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.1_743)	Depositor
R, R_{free}	0.152 , 0.168 0.145 , 0.161	Depositor DCC
R_{free} test set	6307 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	10.4	Xtriage
Anisotropy	0.347	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 35.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.018 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3296	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.35	0/2905	0.63	0/3936

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2842	0	2754	25	0
2	A	454	0	0	9	1
All	All	3296	0	2754	25	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:386[B]:ASN:ND2	2:A:813:HOH:O	1.93	0.99
1:A:534:PRO:O	2:A:1028:HOH:O	1.84	0.95
1:A:224:SER:N	2:A:1150:HOH:O	2.10	0.83
1:A:465[B]:ASP:OD1	2:A:997:HOH:O	2.07	0.70

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:414:ASP:O	1:A:417[B]:THR:HG22	1.95	0.66
1:A:417[B]:THR:HG21	2:A:766:HOH:O	1.97	0.64
1:A:338:ASN:CG	1:A:339:LEU:H	2.06	0.63
1:A:374:TYR:O	1:A:429:LYS:NZ	2.34	0.60
1:A:386[A]:ASN:ND2	2:A:1112:HOH:O	2.35	0.58
1:A:375:ASP:HA	1:A:429:LYS:HZ2	1.72	0.54
1:A:535:ASN:HA	2:A:1028:HOH:O	2.08	0.54
1:A:547:GLU:CD	1:A:550:GLY:HA2	2.32	0.54
1:A:539:ARG:O	1:A:543[B]:MET:HG3	2.09	0.52
1:A:525:VAL:HG12	1:A:531:ILE:HD12	1.92	0.52
1:A:552:ARG:HD3	2:A:894:HOH:O	2.14	0.48
1:A:363:SER:HB3	1:A:512:SER:HB2	1.97	0.47
1:A:291:PRO:O	2:A:890:HOH:O	2.20	0.46
1:A:525:VAL:CG2	1:A:559:LEU:HG	2.45	0.46
1:A:480:TYR:HB3	1:A:484:LYS:HD3	1.98	0.45
1:A:252:LYS:NZ	1:A:252:LYS:HB3	2.33	0.43
1:A:339:LEU:HD21	1:A:638:ARG:CG	2.49	0.43
1:A:525:VAL:HG21	1:A:559:LEU:HG	2.00	0.43
1:A:512:SER:HB3	1:A:647:ARG:C	2.45	0.41
1:A:309:GLU:HA	1:A:499:THR:O	2.20	0.41
1:A:363:SER:HB3	1:A:512:SER:CB	2.51	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:869:HOH:O	2:A:1073:HOH:O[1_545]	2.08	0.12

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	364/488 (75%)	355 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	305/412 (74%)	302 (99%)	3 (1%)	68	33

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	252	LYS
1	A	429	LYS
1	A	493	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	275	ASN
1	A	319	ASN
1	A	476	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	356/488 (72%)	0.29	25 (7%) 22 26	4, 11, 23, 34	11 (3%)

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	338	ASN	5.0
1	A	548	VAL	4.2
1	A	464	ALA	3.4
1	A	635	ALA	3.4
1	A	549	ASN	3.2
1	A	566	ALA	3.1
1	A	291	PRO	3.1
1	A	292	ASP	3.1
1	A	534	PRO	3.0
1	A	533	ASN	3.0
1	A	224	SER	3.0
1	A	536	GLN	2.9
1	A	550	GLY	2.7
1	A	531	ILE	2.7
1	A	535	ASN	2.6
1	A	453	TYR	2.6
1	A	529	TYR	2.6
1	A	230	TYR	2.4
1	A	638	ARG	2.3
1	A	565	LYS	2.2
1	A	539	ARG	2.1
1	A	552	ARG	2.1
1	A	542	LEU	2.0
1	A	547	GLU	2.0
1	A	248	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.