



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 11:28 AM UTC

PDB ID : 7MZI / pdb_00007mzi
Title : SARS-CoV-2 receptor binding domain bound to Fab WCSL 129
Authors : Pymm, P.; Tan, L.L.; Dietrich, M.H.; Chan, L.J.; Tham, W.H.
Deposited on : 2021-05-24
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	216	
2	H	224	
3	A	205	
4	B	3	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	A	603	-	-	X	-
5	GOL	H	303	-	-	X	-

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 5333 atoms, of which 32 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called WCSL 129 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	212	Total	C	N	O	S	0	0	0
			1563	980	258	321	4			

- Molecule 2 is a protein called WCSL 129 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	220	Total	C	N	O	S	0	2	0
			1627	1024	273	320	10			


- Molecule 3 is a protein called Spike protein S1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	196	Total	C	N	O	S	0	1	0
			1549	990	259	292	8			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	528	GLY	-	expression tag	UNP P0DTC2
A	529	SER	-	expression tag	UNP P0DTC2
A	530	HIS	-	expression tag	UNP P0DTC2
A	531	HIS	-	expression tag	UNP P0DTC2
A	532	HIS	-	expression tag	UNP P0DTC2
A	533	HIS	-	expression tag	UNP P0DTC2
A	534	HIS	-	expression tag	UNP P0DTC2
A	535	HIS	-	expression tag	UNP P0DTC2

- Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	B	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	L	1	Total	C	O		0	0
			6	3	3			
5	L	1	Total	C	O		0	0
			6	3	3			
5	L	1	Total	C	O		0	0
			6	3	3			
5	L	1	Total	C	O		0	0
			6	3	3			
5	H	1	Total	C	O		0	0
			6	3	3			
5	H	1	Total	C	O		0	0
			6	3	3			
5	H	1	Total	C	H	O	0	0
			14	3	8	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	H	1	Total C H O 14 3 8 3	0	0
5	H	1	Total C H O 14 3 8 3	0	0
5	H	1	Total C H O 14 3 8 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0

- Molecule 6 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 13 8 5	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	L	147	Total O 147 147	0	0
7	H	161	Total O 161 161	0	0

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	118	Total 118	O 118	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: WCSL 129 light chain



- Molecule 2: WCSL 129 heavy chain



- Molecule 3: Spike protein S1



- Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	I 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	106.56Å 71.90Å 113.69Å 90.00° 108.32° 90.00°	Depositor
Resolution (Å)	44.55 – 1.85 44.55 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.3 (44.55-1.85) 99.3 (44.55-1.85)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.75 (at 1.86Å)	Xtriage
Refinement program	PHENIX dev_3965	Depositor
R, R_{free}	0.174 , 0.207 0.174 , 0.207	Depositor DCC
R_{free} test set	3616 reflections (5.16%)	wwPDB-VP
Wilson B-factor (Å ²)	25.3	Xtriage
Anisotropy	0.509	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 44.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5333	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, PG4, NAG, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	L	0.48	0/1605	0.66	0/2200
2	H	0.53	0/1665	0.67	0/2266
3	A	0.48	0/1593	0.59	0/2168
All	All	0.50	0/4863	0.64	0/6634

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1563	0	1494	13	0
2	H	1627	0	1583	25	0
3	A	1549	0	1455	34	0
4	B	39	0	34	3	0
5	A	18	0	24	5	0
5	H	42	32	56	11	0
5	L	24	0	32	3	0
6	A	13	0	18	5	0
7	A	118	0	0	2	0
7	H	161	0	0	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	L	147	0	0	4	0
All	All	5301	32	4696	77	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:L:303:GOL:H11	7:L:525:HOH:O	1.60	0.99
1:L:118:THR:HG22	5:L:301:GOL:O2	1.76	0.86
2:H:131:SER:O	2:H:135:THR:HG23	1.77	0.85
3:A:495:TYR:O	5:A:603:GOL:H32	1.80	0.81
3:A:364:ASP:OD2	3:A:367:VAL:HG23	1.82	0.78
2:H:28:THR:OG1	5:H:301:GOL:H32	1.88	0.72
1:L:185:THR:OG1	1:L:188:GLN:HG3	1.89	0.72
1:L:126:SER:O	1:L:130:GLN:HG2	1.90	0.72
2:H:129:ALA:HB3	2:H:218:LYS:HE3	1.74	0.69
5:L:303:GOL:H32	7:L:524:HOH:O	1.90	0.69
7:L:508:HOH:O	5:H:303:GOL:H12	1.92	0.68
3:A:355:ARG:HH22	6:A:601:PG4:H12	1.63	0.64
3:A:403:ARG:NH2	5:A:603:GOL:O3	2.32	0.62
3:A:359:SER:HA	3:A:524:VAL:CG2	2.32	0.60
3:A:339:GLY:O	3:A:343:ASN:HB2	2.01	0.59
3:A:403:ARG:HH21	5:A:603:GOL:C3	2.14	0.59
3:A:359:SER:HA	3:A:524:VAL:HG22	1.84	0.57
1:L:192:HIS:HE1	7:L:531:HOH:O	1.88	0.57
3:A:520:ALA:HB1	3:A:521:PRO:CD	2.36	0.55
3:A:364:ASP:O	3:A:367:VAL:HB	2.06	0.55
2:H:105:ASP:HB3	5:H:307:GOL:O3	2.07	0.54
3:A:393:THR:HG21	3:A:518:LEU:HB2	1.89	0.54
2:H:174:LEU:HB3	5:H:303:GOL:H11	1.90	0.54
2:H:76:LYS:HE3	7:H:453:HOH:O	2.08	0.53
3:A:360:ASN:OD1	3:A:523:THR:HB	2.09	0.53
1:L:198:GLN:NE2	1:L:207:GLU:OE2	2.38	0.52
2:H:83:MET:HE1	2:H:94:TYR:CZ	2.45	0.52
1:L:211:ALA:H	2:H:133:LYS:NZ	2.09	0.51
2:H:174:LEU:H	5:H:303:GOL:H11	1.76	0.50
2:H:102:GLY:HA2	3:A:489:TYR:OH	2.11	0.50
6:A:601:PG4:C4	6:A:601:PG4:H71	2.43	0.49
3:A:388:ASN:HB3	3:A:527:PRO:HD2	1.94	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:116:ASN:OD1	1:L:117:PRO:HD2	2.12	0.49
4:B:2:NAG:H62	4:B:3:BMA:H2	1.93	0.49
3:A:388:ASN:HA	3:A:526:GLY:HA3	1.95	0.48
3:A:447:GLY:HA3	3:A:449:TYR:CE1	2.50	0.47
3:A:429:PHE:CZ	3:A:431:GLY:HA3	2.49	0.47
1:L:119:VAL:O	1:L:208:LYS:HE3	2.14	0.47
1:L:187:GLU:O	1:L:191:SER:HB3	2.15	0.46
2:H:3:GLN:NE2	7:H:404:HOH:O	2.48	0.46
3:A:383:SER:HB3	3:A:386:LYS:HG3	1.98	0.46
1:L:154:ALA:HB2	1:L:195:TYR:CE2	2.51	0.46
2:H:19:ARG:HD2	7:H:529:HOH:O	2.14	0.46
2:H:135:THR:HG22	2:H:140:ALA:CB	2.45	0.46
3:A:369:TYR:CD1	3:A:369:TYR:O	2.68	0.46
1:L:29:ILE:O	1:L:67:LYS:HE3	2.16	0.46
2:H:12:VAL:HG11	2:H:18:LEU:HG	1.96	0.46
6:A:601:PG4:H61	6:A:601:PG4:H42	1.79	0.45
1:L:13:THR:HB	1:L:14:PRO:CD	2.47	0.45
2:H:101:TRP:CD1	3:A:455:LEU:CD2	3.00	0.45
2:H:129:ALA:CB	2:H:218:LYS:HE3	2.44	0.45
5:H:301:GOL:H11	7:H:424:HOH:O	2.17	0.45
4:B:1:NAG:H61	4:B:2:NAG:C7	2.47	0.44
4:B:1:NAG:H82	4:B:1:NAG:O3	2.17	0.44
3:A:357:ARG:NH2	7:A:703:HOH:O	2.49	0.44
2:H:101:TRP:CE3	5:H:307:GOL:H12	2.53	0.44
5:A:602:GOL:C3	7:A:714:HOH:O	2.65	0.44
3:A:355:ARG:HH12	6:A:601:PG4:H12	1.81	0.44
1:L:136:LEU:HD12	1:L:136:LEU:N	2.31	0.43
2:H:101:TRP:CD1	3:A:455:LEU:HD23	2.53	0.43
3:A:417:LYS:HD2	3:A:453:TYR:CD1	2.54	0.43
3:A:383:SER:HB3	3:A:386:LYS:CG	2.49	0.43
2:H:174:LEU:H	5:H:303:GOL:C1	2.32	0.42
2:H:212:ASP:H	5:H:305:GOL:C1	2.31	0.42
3:A:364:ASP:CG	3:A:367:VAL:HG23	2.45	0.42
3:A:381:GLY:C	3:A:382:VAL:HG13	2.45	0.42
2:H:174:LEU:O	5:H:303:GOL:H11	2.20	0.41
2:H:149:TYR:CE2	2:H:154:VAL:HG13	2.55	0.41
3:A:380:TYR:O	3:A:430:THR:HA	2.20	0.41
3:A:495:TYR:O	5:A:603:GOL:C3	2.62	0.41
3:A:444:LYS:HG3	3:A:448:ASN:HB2	2.03	0.41
2:H:68:PHE:CE2	2:H:83:MET:HE3	2.56	0.41
3:A:417:LYS:HE2	3:A:417:LYS:HB3	1.84	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:355:ARG:NH2	6:A:601:PG4:H12	2.32	0.41
3:A:517:LEU:O	3:A:518:LEU:HD23	2.21	0.41
2:H:101:TRP:CE3	5:H:307:GOL:C1	3.03	0.40
2:H:102:GLY:HA2	3:A:489:TYR:CZ	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	210/216 (97%)	204 (97%)	6 (3%)	0	100	100
2	H	220/224 (98%)	217 (99%)	3 (1%)	0	100	100
3	A	195/205 (95%)	185 (95%)	10 (5%)	0	100	100
All	All	625/645 (97%)	606 (97%)	19 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	175/181 (97%)	174 (99%)	1 (1%)	78	73
2	H	180/184 (98%)	177 (98%)	3 (2%)	53	42
3	A	168/177 (95%)	168 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	523/542 (96%)	519 (99%)	4 (1%)	78	67

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	191	SER
2	H	142	LEU
2	H	200[A]	CYS
2	H	200[B]	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	39	GLN
2	H	39	GLN
2	H	82	GLN
2	H	84	ASN
2	H	175	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	B	1	4,3	14,14,15	0.40	0	17,19,21	0.66	1 (5%)
4	NAG	B	2	4	14,14,15	0.25	0	17,19,21	0.39	0
4	BMA	B	3	4	11,11,12	0.66	0	15,15,17	0.94	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	B	1	4,3	-	2/6/23/26	0/1/1/1
4	NAG	B	2	4	-	2/6/23/26	0/1/1/1
4	BMA	B	3	4	-	2/2/19/22	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	3	BMA	C1-O5-C5	2.46	115.48	112.19
4	B	1	NAG	C1-O5-C5	2.27	115.23	112.19

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	B	3	BMA	C4-C5-C6-O6
4	B	3	BMA	O5-C5-C6-O6
4	B	1	NAG	C8-C7-N2-C2
4	B	1	NAG	O7-C7-N2-C2
4	B	2	NAG	C4-C5-C6-O6
4	B	2	NAG	O5-C5-C6-O6

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	2	NAG	2	0
4	B	1	NAG	2	0
4	B	3	BMA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	GOL	L	304	-	5,5,5	0.42	0	5,5,5	0.85	0
5	GOL	H	306	-	5,5,5	0.54	0	5,5,5	0.34	0
5	GOL	H	301	-	5,5,5	0.45	0	5,5,5	0.41	0
5	GOL	H	304	-	5,5,5	0.46	0	5,5,5	0.84	0
5	GOL	A	603	-	5,5,5	0.48	0	5,5,5	0.85	0
5	GOL	L	302	-	5,5,5	0.56	0	5,5,5	0.91	0
5	GOL	A	602	-	5,5,5	0.64	0	5,5,5	0.77	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	PG4	A	601	-	12,12,12	0.57	0	11,11,11	0.42	0
5	GOL	A	604	-	5,5,5	0.66	0	5,5,5	0.37	0
5	GOL	L	301	-	5,5,5	0.49	0	5,5,5	0.60	0
5	GOL	H	307	-	5,5,5	0.65	0	5,5,5	1.03	0
5	GOL	H	303	-	5,5,5	0.57	0	5,5,5	0.94	0
5	GOL	L	303	-	5,5,5	0.71	0	5,5,5	0.66	0
5	GOL	H	305	-	5,5,5	0.63	0	5,5,5	0.86	0
5	GOL	H	302	-	5,5,5	0.60	0	5,5,5	1.22	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	L	304	-	-	2/4/4/4	-
5	GOL	H	306	-	-	2/4/4/4	-
5	GOL	H	301	-	-	2/4/4/4	-
5	GOL	H	304	-	-	2/4/4/4	-
5	GOL	A	603	-	-	1/4/4/4	-
5	GOL	L	302	-	-	2/4/4/4	-
5	GOL	A	602	-	-	0/4/4/4	-
6	PG4	A	601	-	-	4/10/10/10	-
5	GOL	A	604	-	-	4/4/4/4	-
5	GOL	L	301	-	-	2/4/4/4	-
5	GOL	H	307	-	-	2/4/4/4	-
5	GOL	H	303	-	-	4/4/4/4	-
5	GOL	L	303	-	-	2/4/4/4	-
5	GOL	H	305	-	-	3/4/4/4	-
5	GOL	H	302	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (36) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	L	301	GOL	O1-C1-C2-C3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
5	L	304	GOL	O2-C2-C3-O3
5	H	301	GOL	O1-C1-C2-O2
5	H	301	GOL	O1-C1-C2-C3
5	H	302	GOL	O1-C1-C2-C3
5	H	302	GOL	C1-C2-C3-O3
5	H	303	GOL	O1-C1-C2-C3
5	H	305	GOL	O1-C1-C2-C3
5	H	307	GOL	O1-C1-C2-C3
5	A	604	GOL	O1-C1-C2-O2
5	A	604	GOL	O1-C1-C2-C3
5	A	604	GOL	C1-C2-C3-O3
5	L	302	GOL	O1-C1-C2-C3
5	L	303	GOL	C1-C2-C3-O3
5	L	304	GOL	C1-C2-C3-O3
5	H	303	GOL	C1-C2-C3-O3
5	H	304	GOL	O1-C1-C2-C3
5	H	306	GOL	C1-C2-C3-O3
5	L	301	GOL	O1-C1-C2-O2
5	H	303	GOL	O1-C1-C2-O2
5	H	304	GOL	O1-C1-C2-O2
5	H	305	GOL	O1-C1-C2-O2
5	H	307	GOL	O1-C1-C2-O2
6	A	601	PG4	O4-C7-C8-O5
5	H	306	GOL	O2-C2-C3-O3
5	A	604	GOL	O2-C2-C3-O3
6	A	601	PG4	C6-C5-O3-C4
5	L	303	GOL	O2-C2-C3-O3
5	H	302	GOL	O1-C1-C2-O2
5	H	302	GOL	O2-C2-C3-O3
5	H	303	GOL	O2-C2-C3-O3
6	A	601	PG4	O1-C1-C2-O2
5	A	603	GOL	O2-C2-C3-O3
5	H	305	GOL	O2-C2-C3-O3
6	A	601	PG4	O2-C3-C4-O3
5	L	302	GOL	O1-C1-C2-O2

There are no ring outliers.

9 monomers are involved in 24 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	H	301	GOL	2	0
5	A	603	GOL	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	602	GOL	1	0
6	A	601	PG4	5	0
5	L	301	GOL	1	0
5	H	307	GOL	3	0
5	H	303	GOL	5	0
5	L	303	GOL	2	0
5	H	305	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	L	212/216 (98%)	0.03	7 (3%) 49 53	18, 29, 55, 68	0
2	H	220/224 (98%)	-0.04	9 (4%) 41 45	12, 27, 48, 66	2 (0%)
3	A	196/205 (95%)	0.69	34 (17%) 4 3	15, 34, 80, 105	1 (0%)
All	All	628/645 (97%)	0.21	50 (7%) 18 19	12, 29, 65, 105	3 (0%)

All (50) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	A	369	TYR	5.5
1	L	26	TYR	5.0
3	A	335	LEU	5.0
2	H	101	TRP	4.9
1	L	116	ASN	4.6
3	A	384	PRO	4.6
3	A	372	ALA	4.2
3	A	518	LEU	4.1
3	A	387	LEU	4.1
3	A	517	LEU	3.8
3	A	529	SER	3.8
3	A	522	ALA	3.7
3	A	367	VAL	3.7
2	H	135	THR	3.4
2	H	133	LYS	3.3
3	A	371	SER	3.3
3	A	521	PRO	3.1
3	A	392	PHE	3.1
3	A	391	CYS	3.1
3	A	519	HIS	3.1
3	A	362	VAL	3.0
3	A	390	LEU	3.0
3	A	385	THR	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	L	213	THR	2.8
3	A	370	ASN	2.8
3	A	393	THR	2.7
3	A	381	GLY	2.6
3	A	528	GLY	2.6
3	A	363	ALA	2.5
3	A	523	THR	2.5
2	H	220	CYS	2.5
3	A	449	TYR	2.4
2	H	137	GLY	2.4
1	L	130	GLN	2.3
2	H	208	ASN	2.3
1	L	2	SER	2.3
2	H	134	SER	2.2
3	A	373	SER	2.2
1	L	160	LYS	2.2
2	H	132	SER	2.2
3	A	366	SER	2.2
3	A	515	PHE	2.2
3	A	524	VAL	2.2
3	A	364	ASP	2.2
3	A	382	VAL	2.1
1	L	158	PRO	2.1
3	A	444	LYS	2.1
2	H	2	VAL	2.1
3	A	388	ASN	2.0
3	A	336	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	B	1	14/15	-	-	44,68,83,100	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	B	2	14/15	-	-	74,91,111,118	0
4	BMA	B	3	11/12	-	-	95,114,122,126	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	GOL	H	307	6/6	0.66	0.24	49,67,87,104	0
5	GOL	H	306	6/6	0.75	0.13	56,72,87,101	0
5	GOL	H	305	6/6	0.76	0.17	38,58,73,80	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	GOL	A	602	6/6	0.80	0.20	33,54,57,61	0
6	PG4	A	601	13/13	0.81	0.16	44,63,72,77	0
5	GOL	H	303	6/6	0.84	0.16	41,54,57,58	0
5	GOL	L	303	6/6	0.85	0.16	31,52,56,59	0
5	GOL	H	301	6/6	0.86	0.15	27,51,66,66	0
5	GOL	A	603	6/6	0.87	0.12	40,46,67,71	0
5	GOL	H	304	6/6	0.87	0.12	48,59,66,71	0
5	GOL	H	302	6/6	0.88	0.13	33,47,57,66	0
5	GOL	L	302	6/6	0.90	0.12	31,47,52,53	0
5	GOL	A	604	6/6	0.92	0.11	39,47,51,56	0
5	GOL	L	304	6/6	0.92	0.13	27,49,58,60	0
5	GOL	L	301	6/6	0.93	0.11	28,38,48,50	0

6.5 Other polymers [i](#)

There are no such residues in this entry.