



wwPDB EM Validation Summary Report ⓘ

Mar 30, 2026 – 12:05 AM UTC

PDB ID : 1N03 / pdb_00001n03
Title : Model for Active RecA Filament
Authors : VanLoock, M.S.; Yu, X.; Yang, S.; Lai, A.L.; Low, C.; Campbell, M.J.; Egelman, E.H.
Deposited on : 2002-10-10
Resolution : 20.00 Å (reported)
Based on initial model : 1REA

This is a wwPDB EM Validation Summary Report for a publicly released PDB/EMDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 20.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	229148	23984

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	352	85% . 14%
1	B	352	85% . 14%
1	C	352	85% . 14%
1	D	352	85% . 14%
1	E	352	85% . 14%
1	F	352	85% . 14%
1	G	352	85% . 14%

2 Entry composition [i](#)


There are 2 unique types of molecules in this entry. The entry contains 2310 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RecA protein.

Mol	Chain	Residues	Atoms	AltConf	Trace
1	A	303	Total C 303 303	0	303
1	B	303	Total C 303 303	0	303
1	C	303	Total C 303 303	0	303
1	D	303	Total C 303 303	0	303
1	E	303	Total C 303 303	0	303
1	F	303	Total C 303 303	0	303
1	G	303	Total C 303 303	0	303

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).




Mol	Chain	Residues	Atoms					AltConf
2	A	1	Total 27	C 10	N 5	O 10	P 2	0
2	B	1	Total 27	C 10	N 5	O 10	P 2	0
2	C	1	Total 27	C 10	N 5	O 10	P 2	0
2	D	1	Total 27	C 10	N 5	O 10	P 2	0
2	E	1	Total 27	C 10	N 5	O 10	P 2	0
2	F	1	Total 27	C 10	N 5	O 10	P 2	0
2	G	1	Total 27	C 10	N 5	O 10	P 2	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: RecA protein

Chain A:  85% 14%




- Molecule 1: RecA protein

Chain B:  85% 14%




- Molecule 1: RecA protein

Chain C:  85% 14%



- Molecule 1: RecA protein

Chain D:  85% 14%




- Molecule 1: RecA protein

[illegible]

- Chain F: 85% • 14%

GLU	ALA
THR	ILE
ASN	D3
GLU	G71
ASP	K72
PHE	I73
	A98
	F156
	GLY
	GLU
	ILE
	GLY
	ASP
	SER
	HIS
	MET
	G165
	Q194
	ILE
	ARG
	MET
	LYS
	ILE
	GLY
	VAL
	MET
	PHE
	GLY
	ASN
	PRO
	GLU
	THR
	THR
	T210
	F217
	L328
	SER
	ASN
	PRO
	ASN
	ASN
	SER
	THR
	PRO
	ASP
	PHE
	SER
	VAL
	ASP
	ASP
	SER
	GLU
	VAL
	ALA

- Chain G:  85% 14%

ASN	ALA	D3	G71	K72	T73	E156	GLY	GLU	ILE	ILE	ARG	MET	LYS	ILE	GLY	VAL	MET	PHE	GLY	ASN	PRO	GLU	THR	THR	T210	F217	L328	SER	ASN	PRO	ASN	SER	THR	PRO	ASP	PHE	SER	VAL	ASP	SER	GLY	VAL	ALA	GLU	THR
-----	-----	----	-----	-----	-----	------	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

4 Data and refinement statistics

Xtrriage (Phenix) and EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	1.00Å 1.00Å 1.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	(Not available) – 20.00	Depositor
% Data completeness (in resolution range)	(Not available) ((Not available)-20.00)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	unknown	Depositor
R, R_{free}	(Not available) , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	2310	wwPDB-VP
Average B, all atoms (Å ²)	10.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

There are no protein, RNA or DNA chains available to summarize Z scores of covalent bonds and angles.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	303	0	0	3	0
1	B	303	0	0	4	0
1	C	303	0	0	4	0
1	D	303	0	0	4	0
1	E	303	0	0	4	0
1	F	303	0	0	4	0
1	G	303	0	0	3	0
2	A	27	0	12	2	0
2	B	27	0	12	2	0
2	C	27	0	12	2	0
2	D	27	0	12	2	0
2	E	27	0	12	2	0
2	F	27	0	12	2	0
2	G	27	0	12	2	0
All	All	2310	0	84	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 20 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:73:THR:CA	2:F:506:ADP:O1A	2.50	0.60
1:G:73:THR:CA	2:G:507:ADP:O1A	2.50	0.60
1:B:73:THR:CA	2:B:502:ADP:O1A	2.50	0.60
1:C:73:THR:CA	2:C:503:ADP:O1A	2.50	0.60
1:E:73:THR:CA	2:E:505:ADP:O1A	2.50	0.60

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein backbone outliers to report in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ADP	C	503	-	28,29,29	0.82	1 (3%)	43,45,45	1.13	1 (2%)
2	ADP	G	507	-	28,29,29	0.83	1 (3%)	43,45,45	1.13	1 (2%)
2	ADP	B	502	-	28,29,29	0.83	1 (3%)	43,45,45	1.13	1 (2%)
2	ADP	A	501	-	28,29,29	0.83	1 (3%)	43,45,45	1.13	1 (2%)
2	ADP	E	505	-	28,29,29	0.83	1 (3%)	43,45,45	1.13	1 (2%)
2	ADP	D	504	-	28,29,29	0.83	1 (3%)	43,45,45	1.13	1 (2%)
2	ADP	F	506	-	28,29,29	0.83	1 (3%)	43,45,45	1.13	1 (2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADP	C	503	-	-	3/16/32/32	0/3/3/3
2	ADP	G	507	-	-	3/16/32/32	0/3/3/3
2	ADP	B	502	-	-	3/16/32/32	0/3/3/3
2	ADP	A	501	-	-	3/16/32/32	0/3/3/3
2	ADP	E	505	-	-	3/16/32/32	0/3/3/3
2	ADP	D	504	-	-	3/16/32/32	0/3/3/3
2	ADP	F	506	-	-	3/16/32/32	0/3/3/3

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	502	ADP	PB-O2B	-2.14	1.46	1.54
2	A	501	ADP	PB-O2B	-2.13	1.46	1.54
2	G	507	ADP	PB-O2B	-2.13	1.46	1.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	506	ADP	PB-O2B	-2.12	1.46	1.54
2	E	505	ADP	PB-O2B	-2.12	1.46	1.54

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	504	ADP	O4'-C1'-N9	4.12	116.00	108.09
2	F	506	ADP	O4'-C1'-N9	4.11	115.99	108.09
2	E	505	ADP	O4'-C1'-N9	4.11	115.98	108.09
2	G	507	ADP	O4'-C1'-N9	4.11	115.98	108.09
2	B	502	ADP	O4'-C1'-N9	4.11	115.98	108.09

There are no chirality outliers.

5 of 21 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	ADP	PA-O3A-PB-O3B
2	B	502	ADP	PA-O3A-PB-O3B
2	C	503	ADP	PA-O3A-PB-O3B
2	D	504	ADP	PA-O3A-PB-O3B
2	E	505	ADP	PA-O3A-PB-O3B


There are no ring outliers.



7 monomers are involved in 14 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	503	ADP	2	0
2	G	507	ADP	2	0
2	B	502	ADP	2	0
2	A	501	ADP	2	0
2	E	505	ADP	2	0
2	D	504	ADP	2	0
2	F	506	ADP	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.