



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:14 AM UTC

PDB ID : 4N0H / pdb_00004n0h
Title : Crystal structure of S. cerevisiae mitochondrial GatFAB
Authors : Araiso, Y.; Ishitani, R.; Nureki, O.
Deposited on : 2013-10-02
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	464	2% 86% 9% • •
2	B	325	11% 76% 10% • 12%
3	F	160	6% 69% 7% • 22%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6874 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamyl-tRNA(Gln) amidotransferase subunit A, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	446	Total	C	N	O	S	0	8	0
			3425	2193	569	650	13			

- Molecule 2 is a protein called Glutamyl-tRNA(Gln) amidotransferase subunit B, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	285	Total	C	N	O	S	0	1	0
			2210	1403	380	421	6			


There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	330	LEU	-	expression tag	UNP P33893
B	331	GLU	-	expression tag	UNP P33893
B	332	SER	-	expression tag	UNP P33893
B	333	SER	-	expression tag	UNP P33893
B	334	GLY	-	expression tag	UNP P33893
B	335	GLU	-	expression tag	UNP P33893
B	336	ASN	-	expression tag	UNP P33893
B	337	LEU	-	expression tag	UNP P33893
B	338	TYR	-	expression tag	UNP P33893
B	339	PHE	-	expression tag	UNP P33893
B	340	GLN	-	expression tag	UNP P33893

- Molecule 3 is a protein called Glutamyl-tRNA(Gln) amidotransferase subunit F, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	F	125	Total	C	N	O	S	0	0	0
			961	615	167	177	2			

- Molecule 4 is NITRATE ION (CCD ID: NO3) (formula: NO_3).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	N	O	0	0
			4	1	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	179	Total	O	0	0
			179	179		
5	B	57	Total	O	0	0
			57	57		
5	F	38	Total	O	0	0
			38	38		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glutamyl-tRNA(Gln) amidotransferase subunit A, mitochondrial



- Molecule 2: Glutamyl-tRNA(Gln) amidotransferase subunit B, mitochondrial



- Molecule 3: Glutamyl-tRNA(Gln) amidotransferase subunit F, mitochondrial



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	61.95Å 86.30Å 191.23Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.58 – 1.95 30.58 – 1.95	Depositor EDS
% Data completeness (in resolution range)	98.8 (30.58-1.95) 98.7 (30.58-1.95)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.38 (at 1.95Å)	Xtriage
Refinement program	PHENIX 1.7.2 _869	Depositor
R, R_{free}	0.188 , 0.222 0.182 , 0.216	Depositor DCC
R_{free} test set	3723 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	27.6	Xtriage
Anisotropy	0.030	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 56.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6874	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.84% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NO3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.36	0/3511	0.73	0/4757
2	B	0.31	0/2252	0.71	0/3062
3	F	0.30	0/976	0.71	0/1313
All	All	0.34	0/6739	0.72	0/9132

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3425	0	3446	23	0
2	B	2210	0	2159	19	0
3	F	961	0	969	9	0
4	A	4	0	0	0	0
5	A	179	0	0	3	0
5	B	57	0	0	2	0
5	F	38	0	0	0	0
All	All	6874	0	6574	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:436:LEU:HD21	1:A:444:LEU:HD22	1.72	0.72
3:F:51:TRP:H	3:F:51:TRP:CD1	2.08	0.71
3:F:51:TRP:H	3:F:51:TRP:HD1	1.39	0.69
1:A:355:GLU:OE1	5:A:721:HOH:O	2.14	0.66
1:A:243:PHE:HZ	1:A:423:PRO:HG2	1.62	0.63
1:A:418:ASP:HB3	1:A:422:VAL:HG23	1.81	0.62
2:B:261:GLN:HG2	2:B:273:LEU:HD12	1.82	0.61
1:A:168:LYS:NZ	1:A:429:LEU:O	2.37	0.57
2:B:240:GLU:HG3	2:B:278:THR:HB	1.87	0.56
1:A:156:ARG:HH11	1:A:445:GLN:NE2	2.05	0.54
1:A:25:THR:O	1:A:111:ARG:NH1	2.41	0.54
2:B:141:ILE:HG22	2:B:143:GLU:H	1.73	0.53
1:A:377:ASN:ND2	5:A:697:HOH:O	2.36	0.52
1:A:195:LYS:HE2	1:A:198:VAL:HG21	1.92	0.52
1:A:156:ARG:HH11	1:A:445:GLN:HE22	1.58	0.51
1:A:307:TYR:CE2	1:A:326:LYS:HG3	2.46	0.50
2:B:87:MET:HE2	2:B:311:LEU:HD13	1.93	0.50
1:A:308:ARG:NH2	1:A:313:ASP:OD2	2.44	0.49
2:B:320[A]:ARG:NH1	5:B:433:HOH:O	2.23	0.49
3:F:134:ASN:HD21	3:F:137:LYS:HG2	1.78	0.48
2:B:35:ILE:HG21	2:B:225:MET:HE3	1.95	0.48
3:F:134:ASN:HD22	3:F:134:ASN:C	2.21	0.48
3:F:141:GLY:O	3:F:145:GLN:HG3	2.13	0.48
2:B:257:GLU:OE2	2:B:260:ARG:NH1	2.47	0.47
2:B:270:THR:O	2:B:270:THR:OG1	2.29	0.47
2:B:231:LEU:HD12	2:B:239:VAL:HG13	1.96	0.46
1:A:186:LEU:HD21	1:A:426:LEU:HD23	1.97	0.45
1:A:212:LYS:NZ	3:F:150:GLU:OE1	2.43	0.45
1:A:20:LYS:HD2	1:A:121:HIS:CD2	2.52	0.44
2:B:113:GLN:HE22	2:B:118:GLN:HB2	1.83	0.43
2:B:152:GLN:OE1	2:B:154:GLN:NE2	2.52	0.43
2:B:320[A]:ARG:NH2	5:B:442:HOH:O	2.51	0.43
1:A:130:SER:H	1:A:154[A]:SER:HB3	1.83	0.43
2:B:246:ASN:O	2:B:250:ILE:HG13	2.18	0.43
2:B:40:ASN:C	2:B:40:ASN:HD22	2.27	0.43
1:A:197[B]:ASN:ND2	1:A:197[B]:ASN:H	2.16	0.42
1:A:283:PRO:HB2	3:F:107:LEU:HD13	2.02	0.42
3:F:69:PRO:HG3	3:F:97:LEU:HD23	2.01	0.42
2:B:87:MET:HE1	2:B:316:ILE:HD13	2.02	0.41
2:B:102:GLN:HG2	2:B:124:ARG:HB2	2.01	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:F:107:LEU:HD12	3:F:107:LEU:HA	1.90	0.41
1:A:224:LEU:HD23	1:A:224:LEU:HA	1.95	0.41
2:B:102:GLN:NE2	2:B:104:ASP:OD2	2.54	0.41
1:A:264:ILE:HG13	1:A:270:ILE:HD12	2.02	0.41
1:A:200:ARG:NH1	1:A:461:GLU:OE1	2.45	0.41
1:A:350:ASN:HB2	5:A:776:HOH:O	2.21	0.41
2:B:44:LYS:HB3	2:B:181:ASN:O	2.22	0.40
1:A:150:ASP:HB2	1:A:155:VAL:HB	2.03	0.40
2:B:244:LEU:HA	2:B:245:PRO:HD3	1.78	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	448/464 (97%)	439 (98%)	9 (2%)	0	100	100
2	B	278/325 (86%)	276 (99%)	2 (1%)	0	100	100
3	F	119/160 (74%)	116 (98%)	3 (2%)	0	100	100
All	All	845/949 (89%)	831 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	385/407 (95%)	368 (96%)	17 (4%)	25	15
2	B	239/292 (82%)	222 (93%)	17 (7%)	13	4
3	F	99/137 (72%)	91 (92%)	8 (8%)	11	3
All	All	723/836 (86%)	681 (94%)	42 (6%)	19	7

All (42) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	LEU
1	A	14	LEU
1	A	92	LEU
1	A	142	LEU
1	A	154[A]	SER
1	A	154[B]	SER
1	A	157	LEU
1	A	182	TYR
1	A	197[A]	ASN
1	A	197[B]	ASN
1	A	224	LEU
1	A	235	LEU
1	A	264	ILE
1	A	326	LYS
1	A	334	ARG
1	A	416	ILE
1	A	444	LEU
2	B	40	ASN
2	B	71	LEU
2	B	143	GLU
2	B	151	LEU
2	B	172	ILE
2	B	173	THR
2	B	239	VAL
2	B	247	THR
2	B	261	GLN
2	B	270	THR
2	B	273	LEU
2	B	278	THR
2	B	287	VAL
2	B	296	ILE
2	B	320[A]	ARG
2	B	320[B]	ARG
2	B	322	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	F	48	ARG
3	F	51	TRP
3	F	77	LEU
3	F	95	MET
3	F	107	LEU
3	F	134	ASN
3	F	138	LEU
3	F	159	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	121	HIS
1	A	265	ASN
1	A	445	GLN
2	B	36	HIS
2	B	43	ASN
2	B	113	GLN
2	B	118	GLN
2	B	122	HIS
2	B	154	GLN
2	B	246	ASN
3	F	108	HIS
3	F	134	ASN
3	F	147	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NO3	A	501	-	1,3,3	3.18	1 (100%)	0,3,3	-	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	501	NO3	O1-N	3.18	1.40	1.24

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	446/464 (96%)	-0.10	11 (2%) 58 65	12, 30, 56, 96	8 (1%)
2	B	285/325 (87%)	0.57	37 (12%) 7 8	17, 42, 95, 114	1 (0%)
3	F	125/160 (78%)	0.44	10 (8%) 18 21	23, 39, 71, 93	0
All	All	856/949 (90%)	0.20	58 (6%) 23 27	12, 35, 80, 114	9 (1%)

All (58) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	233	ILE	6.1
2	B	27	TYR	5.8
2	B	270	THR	4.8
1	A	410	SER	4.7
2	B	322	LEU	4.6
3	F	51	TRP	4.4
3	F	129	ASN	4.2
2	B	264	LEU	4.0
3	F	130	THR	3.8
3	F	113	GLU	3.8
3	F	131	LYS	3.8
2	B	141	ILE	3.7
2	B	287	VAL	3.7
1	A	43	THR	3.6
1	A	6	SER	3.6
2	B	274	MET	3.5
2	B	327	PRO	3.4
2	B	288	LYS	3.4
2	B	248	SER	3.3
2	B	140	ASP	3.1
2	B	259	GLN	3.0
2	B	273	LEU	3.0
1	A	409	LYS	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	416	ILE	3.0
2	B	171	VAL	2.9
1	A	33	SER	2.8
2	B	246	ASN	2.8
1	A	42	GLU	2.8
2	B	281	TRP	2.8
3	F	109	ASN	2.6
2	B	289	LEU	2.5
2	B	285	SER	2.5
1	A	140	CYS	2.4
2	B	262	VAL	2.4
2	B	296	ILE	2.4
3	F	112	VAL	2.4
2	B	284	SER	2.4
2	B	286	THR	2.3
2	B	172	ILE	2.3
2	B	144	SER	2.3
2	B	247	THR	2.3
2	B	276	PRO	2.3
3	F	58	GLY	2.3
1	A	19	SER	2.3
3	F	108	HIS	2.3
2	B	239	VAL	2.3
2	B	208	ASN	2.3
2	B	232	SER	2.2
1	A	407	GLU	2.2
2	B	138	LEU	2.2
2	B	236	TYR	2.2
2	B	235	GLU	2.1
2	B	237	ALA	2.1
2	B	143	GLU	2.1
2	B	234	ASN	2.1
2	B	280	GLY	2.0
1	A	417	ASN	2.0
3	F	95	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NO3	A	501	4/4	0.95	0.11	42,42,48,48	0

6.5 Other polymers [i](#)

There are no such residues in this entry.