



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 12:25 PM UTC

PDB ID : 5N0S / pdb_00005n0s
Title : Crystal structure of OphA-DeltaC6 mutant Y98A in complex with SAM
Authors : Song, H.; Naismith, J.H.
Deposited on : 2017-02-03
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	410	14% 93%
1	B	410	14% 94%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MLI	B	502	-	-	X	-

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 6895 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Peptide N-methyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	401	Total	C	N	O	S	0	6	0
			3146	1996	540	591	19			
1	B	404	Total	C	N	O	S	0	5	0
			3157	2004	542	593	18			

- Molecule 2 is S-ADENOSYLMETHIONINE (CCD ID: SAM) (formula: $C_{15}H_{22}N_6O_5S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			27	15	6	5	1		
2	B	1	Total	C	N	O	S	0	0
			27	15	6	5	1		

- Molecule 3 is MALONATE ION (CCD ID: MLI) (formula: $C_3H_2O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	3	4		
3	B	1	Total	C	O	0	0
			7	3	4		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	259	Total	O	0	0
			259	259		
4	B	265	Total	O	0	0
			265	265		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Peptide N-methyltransferase



- Molecule 1: Peptide N-methyltransferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	163.91Å 92.52Å 85.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	85.33 – 1.95 85.33 – 1.95	Depositor EDS
% Data completeness (in resolution range)	98.3 (85.33-1.95) 98.3 (85.33-1.95)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.56 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.205 , 0.233 0.211 , 0.238	Depositor DCC
R_{free} test set	4632 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å ²)	35.8	Xtriage
Anisotropy	0.117	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 41.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6895	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 82.87 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.5074e-07. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SAM, MLI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.64	0/3223	0.85	1/4384 (0.0%)
1	B	0.64	0/3234	0.86	0/4400
All	All	0.64	0/6457	0.85	1/8784 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	185	GLY	N-CA-C	6.50	118.12	111.95

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3146	0	3112	19	0
1	B	3157	0	3127	17	0
2	A	27	0	22	0	0
2	B	27	0	22	0	0
3	A	7	0	2	1	0
3	B	7	0	2	4	0
4	A	259	0	0	0	0
4	B	265	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	6895	0	6287	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:216[B]:MET:HB3	1:A:217[B]:PRO:HD2	1.35	1.08
1:A:216[B]:MET:HB3	1:A:217[B]:PRO:CD	1.84	1.03
1:B:216[A]:MET:HG3	1:B:219[A]:GLN:CD	2.15	0.72
1:A:208:VAL:HG22	1:A:248:ILE:HD12	1.78	0.64
1:B:404:VAL:HG12	1:B:406:VAL:HG13	1.79	0.64
1:A:217[B]:PRO:O	1:A:218[B]:HIS:CB	2.46	0.63
1:B:216[A]:MET:HB2	1:B:219[A]:GLN:HG3	1.82	0.62
1:B:267:LEU:HB3	1:B:268:PRO:HA	1.81	0.62
1:A:217[B]:PRO:O	1:A:218[B]:HIS:HB3	2.01	0.60
1:A:266:LEU:HD13	1:A:406:VAL:HG21	1.84	0.58
1:A:217[B]:PRO:O	1:A:218[B]:HIS:CD2	2.58	0.56
1:A:216[B]:MET:CB	1:A:217[B]:PRO:CD	2.63	0.55
1:A:218[B]:HIS:O	1:A:218[B]:HIS:ND1	2.40	0.55
1:A:68:ASN:ND2	1:B:314[B]:GLN:OE1	2.40	0.52
1:B:272:VAL:HG13	1:B:406:VAL:HG12	1.94	0.50
1:A:406:VAL:HG23	1:A:407:ILE:HG13	1.94	0.50
1:B:208:VAL:HG13	1:B:248:ILE:CD1	2.41	0.49
1:A:217[B]:PRO:O	1:A:218[B]:HIS:CG	2.66	0.48
1:A:183:PHE:O	1:B:396:ASN:HB2	2.14	0.48
1:B:208:VAL:HG22	1:B:248:ILE:HD12	1.98	0.45
1:A:208:VAL:HG13	1:A:248:ILE:CD1	2.48	0.43
1:A:400:TRP:CZ2	1:B:172:GLN:HB3	2.53	0.43
3:A:502:MLI:H11	1:B:305:GLN:HB2	2.00	0.43
1:A:104:PHE:CE2	1:A:171:PHE:HB2	2.53	0.43
1:A:305:GLN:CB	3:B:502:MLI:H12	2.48	0.42
1:A:144:ASN:HA	1:A:145:PRO:HA	1.90	0.42
1:B:65:TYR:CZ	1:B:82:LEU:HD11	2.54	0.42
1:A:305:GLN:HB2	3:B:502:MLI:H12	2.01	0.42
1:B:272:VAL:HG13	1:B:406:VAL:CG1	2.49	0.41
1:B:70:LYS:HE3	3:B:502:MLI:H11	2.03	0.41
1:B:134:LEU:C	1:B:134:LEU:HD23	2.46	0.41
1:B:352:VAL:O	1:B:360:ARG:NH2	2.54	0.40
1:B:78:GLN:HE22	3:B:502:MLI:C1	2.35	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	405/410 (99%)	385 (95%)	18 (4%)	2 (0%)	24	16
1	B	407/410 (99%)	389 (96%)	17 (4%)	1 (0%)	43	36
All	All	812/820 (99%)	774 (95%)	35 (4%)	3 (0%)	43	21

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	216[A]	MET
1	A	216[B]	MET
1	B	268	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	337/337 (100%)	336 (100%)	1 (0%)	86	87
1	B	338/337 (100%)	335 (99%)	3 (1%)	70	70
All	All	675/674 (100%)	671 (99%)	4 (1%)	81	79

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	71	SER
1	B	216[A]	MET
1	B	216[B]	MET
1	B	406	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	64	GLN
1	A	165	HIS
1	A	350	GLN
1	B	165	HIS
1	B	284	GLN
1	B	300	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	MLI	B	502	-	6,6,6	1.12	0	7,7,7	0.96	1 (14%)
2	SAM	A	501	-	27,29,29	1.38	4 (14%)	34,42,42	2.14	12 (35%)
2	SAM	B	501	-	27,29,29	1.50	4 (14%)	34,42,42	2.04	11 (32%)
3	MLI	A	502	-	6,6,6	1.10	0	7,7,7	1.80	2 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MLI	B	502	-	-	0/4/4/4	-
2	SAM	A	501	-	-	4/17/33/33	0/3/3/3
2	SAM	B	501	-	-	2/17/33/33	0/3/3/3
3	MLI	A	502	-	-	0/4/4/4	-

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	SAM	C5-C4	4.79	1.47	1.39
2	A	501	SAM	C5-C4	3.91	1.46	1.39
2	B	501	SAM	C8-N7	2.65	1.36	1.31
2	A	501	SAM	C4-N9	-2.58	1.32	1.37
2	B	501	SAM	C4-N9	-2.40	1.32	1.37
2	B	501	SAM	C5-C6	2.31	1.47	1.41
2	A	501	SAM	C5-N7	-2.26	1.35	1.39
2	A	501	SAM	C5-C6	2.22	1.47	1.41

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	SAM	CG-SD-C5'	5.31	116.41	103.43
2	B	501	SAM	CG-SD-C5'	4.79	115.14	103.43
2	A	501	SAM	C5-C4-N3	-4.56	120.44	126.72
2	B	501	SAM	C5-C4-N3	-4.47	120.57	126.72
2	A	501	SAM	N3-C2-N1	-4.00	122.53	128.58
2	A	501	SAM	N3-C4-N9	3.85	133.72	127.17
2	B	501	SAM	N3-C4-N9	3.85	133.71	127.17
2	B	501	SAM	N3-C2-N1	-3.72	122.95	128.58
2	B	501	SAM	C4-N9-C8	3.54	109.45	105.74
2	A	501	SAM	C4-N9-C8	3.31	109.21	105.74

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	SAM	C2-N3-C4	3.28	119.84	111.83
2	B	501	SAM	C2-N3-C4	3.11	119.42	111.83
2	A	501	SAM	C2-N1-C6	2.87	123.44	118.73
2	A	501	SAM	C4-C5-N7	-2.77	107.41	110.58
3	A	502	MLI	O6-C2-C1	-2.67	114.50	122.11
2	B	501	SAM	C4-C5-N7	-2.59	107.62	110.58
2	B	501	SAM	C2-N1-C6	2.48	122.80	118.73
2	A	501	SAM	N9-C8-N7	-2.44	110.47	113.94
3	B	502	MLI	C3-C1-C2	2.39	121.40	112.95
3	A	502	MLI	O7-C2-C1	2.39	121.91	114.51
2	B	501	SAM	C2'-C1'-N9	-2.33	107.51	113.30
2	A	501	SAM	C5-N7-C8	2.30	107.06	103.45
2	B	501	SAM	N9-C8-N7	-2.26	110.72	113.94
2	A	501	SAM	C2'-C1'-N9	-2.09	108.10	113.30
2	A	501	SAM	OXT-C-O	-2.07	119.37	124.08
2	B	501	SAM	C6-C5-N7	2.05	136.04	132.09

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	SAM	C4'-C5'-SD-CE
2	B	501	SAM	OXT-C-CA-CB
2	B	501	SAM	O-C-CA-CB
2	A	501	SAM	OXT-C-CA-CB
2	A	501	SAM	O-C-CA-CB
2	A	501	SAM	CB-CG-SD-C5'


There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	502	MLI	4	0
3	A	502	MLI	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring

in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	401/410 (97%)	0.73	58 (14%) 6 6	15, 41, 85, 117	6 (1%)
1	B	404/410 (98%)	0.85	58 (14%) 6 6	21, 42, 84, 120	5 (1%)
All	All	805/820 (98%)	0.79	116 (14%) 6 6	15, 41, 86, 120	11 (1%)

All (116) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	407	ILE	9.1
1	B	272	VAL	8.6
1	A	272	VAL	7.5
1	B	217[A]	PRO	7.4
1	B	410	ILE	6.8
1	B	393	ALA	6.4
1	B	386	ALA	6.1
1	B	394	SER	6.0
1	A	179	ALA	6.0
1	B	407	ILE	5.9
1	B	409	VAL	5.8
1	B	179	ALA	5.5
1	A	396	ASN	5.5
1	A	268	PRO	5.2
1	A	269	ALA	5.2
1	B	269	ALA	5.0
1	A	386	ALA	5.0
1	B	277	ALA	4.9
1	B	383	LEU	4.9
1	A	267	LEU	4.8
1	A	404	VAL	4.7
1	A	7	THR	4.7
1	B	7	THR	4.6
1	B	69	GLY	4.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	369	TRP	4.5
1	B	289	VAL	4.5
1	B	396	ASN	4.5
1	A	270	GLY	4.3
1	B	408	GLY	4.3
1	B	268	PRO	4.2
1	B	403	VAL	4.1
1	A	277	ALA	4.1
1	A	178	ILE	4.1
1	A	394	SER	4.0
1	A	271	GLN	4.0
1	A	406	VAL	4.0
1	A	69	GLY	3.8
1	A	44	ILE	3.8
1	B	178	ILE	3.8
1	B	186	PHE	3.7
1	A	343	ALA	3.7
1	B	44	ILE	3.7
1	B	395	GLN	3.7
1	A	186	PHE	3.7
1	B	273	PRO	3.7
1	B	389	SER	3.6
1	A	385	ALA	3.6
1	A	395	GLN	3.5
1	A	400	TRP	3.4
1	B	267	LEU	3.4
1	B	406	VAL	3.4
1	B	379	PRO	3.3
1	B	382	LEU	3.2
1	B	343	ALA	3.1
1	A	405	GLY	3.1
1	A	217[A]	PRO	3.1
1	A	308	ASP	3.0
1	B	387	ARG	3.0
1	B	388	GLU	3.0
1	B	270	GLY	3.0
1	A	388	GLU	3.0
1	A	289	VAL	2.9
1	B	404	VAL	2.9
1	B	266	LEU	2.9
1	A	390	GLY	2.9
1	B	219[A]	GLN	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	185	GLY	2.9
1	B	392	GLU	2.9
1	B	377	ASN	2.8
1	A	387	ARG	2.8
1	A	398	PHE	2.8
1	B	366	GLY	2.8
1	A	273	PRO	2.8
1	A	346	ARG	2.7
1	B	308	ASP	2.7
1	A	403	VAL	2.7
1	B	271	GLN	2.7
1	A	259	ASP	2.6
1	A	366	GLY	2.6
1	B	274	ASP	2.6
1	A	238	LYS	2.6
1	A	389	SER	2.6
1	B	364	GLU	2.6
1	B	313	GLU	2.5
1	A	369	TRP	2.5
1	A	370	ALA	2.5
1	B	216[A]	MET	2.5
1	A	393	ALA	2.5
1	B	218[A]	HIS	2.5
1	B	276	LYS	2.5
1	B	385	ALA	2.4
1	A	288	ASP	2.4
1	B	185	GLY	2.4
1	B	400	TRP	2.4
1	A	377	ASN	2.3
1	B	184	THR	2.3
1	A	216[A]	MET	2.3
1	A	391	GLU	2.3
1	A	187	ASP	2.3
1	A	340	ASP	2.3
1	A	342	LYS	2.2
1	A	365	LEU	2.2
1	A	337	ALA	2.2
1	A	274	ASP	2.2
1	A	266	LEU	2.1
1	A	276	LYS	2.1
1	B	381	SER	2.1
1	B	390	GLY	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	392	GLU	2.1
1	B	275	LYS	2.1
1	A	372	ARG	2.1
1	A	219[A]	GLN	2.0
1	A	43	VAL	2.0
1	B	175	CYS	2.0
1	B	336	LYS	2.0
1	B	189	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	MLI	B	502	7/7	0.69	0.18	48,63,66,70	0
3	MLI	A	502	7/7	0.70	0.19	51,61,64,70	0
2	SAM	A	501	27/27	0.94	0.08	27,29,34,37	0
2	SAM	B	501	27/27	0.94	0.08	27,30,36,40	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around SAM A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around SAM B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.