



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 17, 2026 – 06:52 PM UTC

PDB ID : 1N1M / pdb_00001n1m
Title : Human Dipeptidyl Peptidase IV/CD26 in complex with an inhibitor
Authors : Rasmussen, H.B.; Branner, S.; Wiberg, F.C.; Wagtmann, N.R.
Deposited on : 2002-10-18
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	728	0.2% 60% 33% 7%
1	B	728	2% 56% 36% 7%
2	C	3	33% 67%
3	D	3	33% 67%
4	E	2	50% 50%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	G	2	 100%
4	H	2	 50% 50%
4	J	2	 50% 50%
5	F	3	 33% 67%
6	I	2	 50% 50%
7	K	3	 33% 67%
7	L	3	 67% 33%
8	M	2	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	D	1	X	-	-	-
4	NAG	J	1	X	-	-	-
5	NDG	F	2	-	-	X	-
8	NAG	M	1	X	-	-	-
9	NAG	A	782	-	-	X	-

2 Entry composition [i](#)


There are 12 unique types of molecules in this entry. The entry contains 13270 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dipeptidyl peptidase IV SOLUBLE FORM.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	726	Total	C	N	O	S	0	0	0
			5948	3816	980	1126	26			
1	B	728	Total	C	N	O	S	0	0	0
			5964	3827	982	1129	26			

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	3	Total	C	N	O	0	0	0
			38	22	2	14			

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	D	3	Total	C	N	O	0	0	0
			38	22	2	14			

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	E	2	Total	C	N	O	0	0	0
			28	16	2	10			
4	G	2	Total	C	N	O	0	0	0
			28	16	2	10			
4	H	2	Total	C	N	O	0	0	0
			28	16	2	10			
4	J	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	F	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 6 is an oligosaccharide called alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	I	2	Total	C	N	O	0	0	0
			24	14	1	9			

- Molecule 7 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
7	K	3	Total	C	N	O	0	0	0
			39	22	2	15			
7	L	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 8 is an oligosaccharide called 2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
8	M	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 9 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	A	1	Total	C	N	O	0	0
			14	8	1	5		
9	B	1	Total	C	N	O	0	0
			14	8	1	5		
9	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 10 is MERCURY (II) ION (CCD ID: HG) (formula: Hg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	A	2	Total Hg 2 2	0	0
10	B	2	Total Hg 2 2	0	0

- Molecule 11 is 2-AMINO-3-METHYL-1-PYRROLIDIN-1-YL-BUTAN-1-ONE (CCD ID: A3M) (formula: C₉H₁₈N₂O).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
11	A	1	Total C N O 12 9 2 1	0	0
11	B	1	Total C N O 12 9 2 1	0	0

- Molecule 12 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
12	A	488	Total O 488 488	0	0
12	B	443	Total O 443 443	0	0

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Chain A:

The visualization displays a complex pattern of colored squares (green, yellow, red) arranged in a grid-like structure, representing the sequence of Chain A. The pattern is highly structured, with many squares appearing in pairs or groups, suggesting a repeating or complementary nature. The colors correspond to the distribution shown in the bar chart above: 60% green, 33% yellow, and 7% red.

Below the grid, several rows of text labels are provided, corresponding to the sequence positions. These labels include various alphanumeric codes, such as R596, E693, K696, G697, L701, L702, L703, L704, L711, L718, L726, L727, L728, L729, L730, L736, L741, L742, L745, L749, L755, L756, L760, L761, L764, LEU, and PRO. Some labels are highlighted in green, yellow, or red, matching the colors of the squares in the grid above them.

Chain B:

2% 56% 36% 7%



- Molecule 2: 2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C: 33% 67%



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D: 33% 67%



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E: 50% 50%



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  100%

NAG1
NAG2

- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:  50% 50%

NAG1
NAG2

- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J:  50% 50%

NAG1
NAG2

- Molecule 5: alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  33% 67%

NAG1
NDG2
MAN3

- Molecule 6: alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:  50% 50%

NAG1
FUC2

- Molecule 7: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain K:  33% 67%

NAG1
NAG2
EM13

- Molecule 7: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain L:  67% 33%



- Molecule 8: 2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain M:

100%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	119.23Å 123.45Å 131.31Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.98 – 2.50 29.98 – 2.50	Depositor EDS
% Data completeness (in resolution range)	87.2 (29.98-2.50) 87.3 (29.98-2.50)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.13 (at 2.51Å)	Xtriage
Refinement program	CNX 2002	Depositor
R, R_{free}	0.208 , 0.263 0.205 , 0.258	Depositor DCC
R_{free} test set	2987 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	28.7	Xtriage
Anisotropy	0.323	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 41.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.018 for k,h,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	13270	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FUC, BMA, MAN, NDG, HG, NAG, A3M

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	0/6119	0.98	35/8321 (0.4%)
1	B	0.41	0/6136	0.95	28/8344 (0.3%)
All	All	0.42	0/12255	0.97	63/16665 (0.4%)

There are no bond length outliers.

The worst 5 of 63 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	530	LEU	CA-C-N	9.83	126.84	119.66
1	B	530	LEU	C-N-CA	9.83	126.84	119.66
1	A	160	VAL	N-CA-C	9.33	118.99	111.62
1	A	207	VAL	N-CA-C	8.53	119.38	111.45
1	B	91	GLU	N-CA-C	8.00	120.25	110.41

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5948	0	5657	260	0
1	B	5964	0	5673	302	0
2	C	38	0	33	2	0
3	D	38	0	34	7	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	E	28	0	25	0	0
4	G	28	0	25	1	0
4	H	28	0	25	2	0
4	J	28	0	25	4	0
5	F	39	0	33	11	0
6	I	24	0	22	2	0
7	K	39	0	34	0	0
7	L	39	0	34	1	0
8	M	28	0	24	4	0
9	A	14	0	13	8	0
9	B	28	0	26	4	0
10	A	2	0	0	0	0
10	B	2	0	0	0	0
11	A	12	0	18	2	0
11	B	12	0	18	3	0
12	A	488	0	0	15	0
12	B	443	0	0	29	0
All	All	13270	0	11719	581	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 24.

The worst 5 of 581 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:F:2:NDG:H6C2	5:F:3:MAN:H2	1.26	1.13
1:A:581:ARG:CZ	9:A:782:NAG:H61	1.81	1.08
1:A:289:ALA:HB1	1:A:290:PRO:HA	1.33	1.06
1:B:289:ALA:HB1	1:B:290:PRO:HA	1.40	1.02
1:B:172:ILE:HG22	1:B:185:ILE:HD11	1.42	1.00

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	724/728 (100%)	661 (91%)	50 (7%)	13 (2%)	6	12
1	B	726/728 (100%)	654 (90%)	58 (8%)	14 (2%)	6	11
All	All	1450/1456 (100%)	1315 (91%)	108 (7%)	27 (2%)	6	11

5 of 27 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	278	SER
1	B	289	ALA
1	B	399	LYS
1	B	645	GLY
1	B	765	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	651/653 (100%)	606 (93%)	45 (7%)	14	30
1	B	653/653 (100%)	604 (92%)	49 (8%)	12	26
All	All	1304/1306 (100%)	1210 (93%)	94 (7%)	13	28

5 of 94 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	232	GLU
1	B	436	LEU
1	B	239	SER
1	B	310	ARG
1	B	507	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 56 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	761	GLN
1	B	731	GLN
1	B	141	GLN
1	B	718	GLN
1	B	606	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

27 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	C	1	2,1	14,14,15	0.77	0	17,19,21	1.15	1 (5%)
2	NDG	C	2	2	14,14,15	0.91	1 (7%)	17,19,21	1.46	3 (17%)
2	FUC	C	3	2	10,10,11	0.50	0	14,14,16	0.36	0
3	NAG	D	1	3,1	14,14,15	0.79	0	17,19,21	0.74	1 (5%)
3	NAG	D	2	3	14,14,15	0.87	1 (7%)	17,19,21	0.62	0
3	FUC	D	3	3	10,10,11	0.62	0	14,14,16	0.38	0
4	NAG	E	1	4,1	14,14,15	0.50	0	17,19,21	0.88	1 (5%)
4	NAG	E	2	4	14,14,15	0.57	0	17,19,21	0.61	0
5	NAG	F	1	5,1	14,14,15	0.60	0	17,19,21	0.88	1 (5%)
5	NDG	F	2	5	14,14,15	0.85	1 (7%)	17,19,21	1.24	3 (17%)
5	MAN	F	3	5	11,11,12	0.72	0	15,15,17	0.43	0
4	NAG	G	1	4,1	14,14,15	0.60	0	17,19,21	1.03	1 (5%)
4	NAG	G	2	4	14,14,15	0.64	0	17,19,21	0.66	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	H	1	4,1	14,14,15	0.61	0	17,19,21	0.92	1 (5%)
4	NAG	H	2	4	14,14,15	0.68	0	17,19,21	0.62	0
6	NAG	I	1	6,1	14,14,15	0.64	0	17,19,21	0.77	1 (5%)
6	FUC	I	2	6	10,10,11	0.58	0	14,14,16	0.37	0
4	NAG	J	1	4,1	14,14,15	0.77	0	17,19,21	1.00	2 (11%)
4	NAG	J	2	4	14,14,15	0.57	0	17,19,21	0.58	0
7	NAG	K	1	7,1	14,14,15	0.46	0	17,19,21	0.81	1 (5%)
7	NAG	K	2	7	14,14,15	0.56	0	17,19,21	0.81	1 (5%)
7	BMA	K	3	7	11,11,12	0.65	0	15,15,17	0.32	0
7	NAG	L	1	7,1	14,14,15	0.65	0	17,19,21	0.80	1 (5%)
7	NAG	L	2	7	14,14,15	0.89	1 (7%)	17,19,21	1.35	3 (17%)
7	BMA	L	3	7	11,11,12	0.73	0	15,15,17	0.94	1 (6%)
8	NAG	M	1	8,1	14,14,15	0.78	0	17,19,21	1.11	2 (11%)
8	NDG	M	2	8	14,14,15	0.79	1 (7%)	17,19,21	0.92	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	2,1	-	0/6/23/26	0/1/1/1
2	NDG	C	2	2	-	2/6/23/26	0/1/1/1
2	FUC	C	3	2	-	-	0/1/1/1
3	NAG	D	1	3,1	1/1/5/7	0/6/23/26	0/1/1/1
3	NAG	D	2	3	-	1/6/23/26	0/1/1/1
3	FUC	D	3	3	-	-	0/1/1/1
4	NAG	E	1	4,1	-	0/6/23/26	0/1/1/1
4	NAG	E	2	4	-	1/6/23/26	0/1/1/1
5	NAG	F	1	5,1	-	2/6/23/26	0/1/1/1
5	NDG	F	2	5	-	3/6/23/26	0/1/1/1
5	MAN	F	3	5	-	2/2/19/22	0/1/1/1
4	NAG	G	1	4,1	-	1/6/23/26	0/1/1/1
4	NAG	G	2	4	-	3/6/23/26	0/1/1/1
4	NAG	H	1	4,1	-	1/6/23/26	0/1/1/1
4	NAG	H	2	4	-	1/6/23/26	0/1/1/1
6	NAG	I	1	6,1	-	1/6/23/26	0/1/1/1
6	FUC	I	2	6	-	-	0/1/1/1
4	NAG	J	1	4,1	1/1/5/7	2/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	J	2	4	-	1/6/23/26	0/1/1/1
7	NAG	K	1	7,1	-	0/6/23/26	0/1/1/1
7	NAG	K	2	7	-	2/6/23/26	0/1/1/1
7	BMA	K	3	7	-	0/2/19/22	0/1/1/1
7	NAG	L	1	7,1	-	2/6/23/26	0/1/1/1
7	NAG	L	2	7	-	3/6/23/26	0/1/1/1
7	BMA	L	3	7	-	2/2/19/22	0/1/1/1
8	NAG	M	1	8,1	1/1/5/7	2/6/23/26	0/1/1/1
8	NDG	M	2	8	-	2/6/23/26	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	2	NAG	C1-C2	2.76	1.56	1.52
2	C	2	NDG	C1-C2	2.46	1.55	1.52
5	F	2	NDG	C1-C2	2.34	1.55	1.52
8	M	2	NDG	C1-C2	2.24	1.55	1.52
7	L	2	NAG	C1-C2	2.05	1.55	1.52

The worst 5 of 25 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	2	NDG	O5-C1-C2	3.94	117.39	111.29
7	L	2	NAG	C4-C3-C2	-3.58	105.78	111.02
2	C	2	NDG	C1-O5-C5	2.93	116.11	112.19
2	C	1	NAG	C4-C3-C2	2.84	115.18	111.02
8	M	1	NAG	C2-N2-C7	-2.60	119.42	122.90

All (3) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	D	1	NAG	C1
4	J	1	NAG	C1
8	M	1	NAG	C1

5 of 34 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	2	NAG	C3-C2-N2-C7
4	G	2	NAG	C1-C2-N2-C7

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
5	F	2	NDG	C3-C2-N2-C7
7	L	3	BMA	C4-C5-C6-O6
8	M	1	NAG	O5-C5-C6-O6


There are no ring outliers.


19 monomers are involved in 34 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1	NAG	1	0
3	D	2	NAG	6	0
3	D	1	NAG	5	0
5	F	3	MAN	3	0
5	F	2	NDG	9	0
4	H	1	NAG	1	0
3	D	3	FUC	1	0
5	F	1	NAG	6	0
4	G	2	NAG	1	0
2	C	3	FUC	1	0
4	J	1	NAG	3	0
2	C	2	NDG	1	0
4	J	2	NAG	1	0
6	I	2	FUC	1	0
7	L	3	BMA	1	0
8	M	2	NDG	4	0
6	I	1	NAG	1	0
4	H	2	NAG	2	0
8	M	1	NAG	4	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





















Oligosaccharide Chain K



Oligosaccharide Chain L





5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 4 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
9	NAG	A	782	1	14,14,15	0.92	1 (7%)	17,19,21	0.78	0
11	A3M	B	955	-	11,12,12	4.66	5 (45%)	13,16,16	2.28	6 (46%)
11	A3M	A	954	-	11,12,12	4.42	5 (45%)	13,16,16	2.31	6 (46%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
9	NAG	B	769	1	14,14,15	0.66	0	17,19,21	0.79	1 (5%)
9	NAG	B	780	1	14,14,15	0.85	1 (7%)	17,19,21	0.70	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	NAG	A	782	1	-	1/6/23/26	0/1/1/1
11	A3M	B	955	-	-	2/12/19/19	0/1/1/1
11	A3M	A	954	-	-	2/12/19/19	0/1/1/1
9	NAG	B	769	1	-	2/6/23/26	0/1/1/1
9	NAG	B	780	1	-	1/6/23/26	0/1/1/1

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
11	B	955	A3M	O18-C14	9.41	1.39	1.22
11	A	954	A3M	O18-C14	8.76	1.38	1.22
11	B	955	A3M	C14-N3	8.41	1.47	1.35
11	A	954	A3M	C14-N3	8.05	1.46	1.35
11	B	955	A3M	C15-C14	-6.51	1.46	1.53

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
11	B	955	A3M	C4-N3-C2	-5.37	104.27	111.37
11	A	954	A3M	C4-N3-C2	-5.14	104.57	111.37
11	A	954	A3M	O18-C14-N3	-3.53	117.40	121.61
11	A	954	A3M	C2-N3-C14	-3.01	113.58	124.03
11	B	955	A3M	C2-N3-C14	-3.01	113.58	124.03

There are no chirality outliers.

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
9	B	780	NAG	C3-C2-N2-C7
11	A	954	A3M	C15-C14-N3-C2
11	A	954	A3M	O18-C14-N3-C2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
11	B	955	A3M	C15-C14-N3-C2
11	B	955	A3M	O18-C14-N3-C2

There are no ring outliers.

4 monomers are involved in 17 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
9	A	782	NAG	8	0
11	B	955	A3M	3	0
11	A	954	A3M	2	0
9	B	780	NAG	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	726/728 (99%)	-0.08	9 (1%) 76 73	14, 30, 54, 65	0
1	B	728/728 (100%)	0.03	18 (2%) 58 54	13, 31, 61, 88	0
All	All	1454/1456 (99%)	-0.02	27 (1%) 66 62	13, 30, 58, 88	0

The worst 5 of 27 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	39	SER	3.6
1	B	39	SER	3.2
1	B	766	PRO	2.8
1	B	103	ASN	2.6
1	A	342	ALA	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	FUC	D	3	10/11	0.19	0.18	79,80,80,80	0
3	NAG	D	2	14/15	0.25	0.17	78,80,80,80	0
4	NAG	H	2	14/15	0.45	0.16	63,65,65,66	0
5	MAN	F	3	11/12	0.48	0.17	71,72,72,73	0
2	NDG	C	2	14/15	0.55	0.14	65,66,67,67	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
8	NDG	M	2	14/15	0.57	0.16	62,64,65,65	0
6	FUC	I	2	10/11	0.58	0.18	73,73,73,73	0
7	BMA	K	3	11/12	0.59	0.15	58,59,59,59	0
3	NAG	D	1	14/15	0.60	0.15	70,73,77,78	0
5	NDG	F	2	14/15	0.63	0.15	63,65,66,69	0
4	NAG	G	2	14/15	0.66	0.16	69,70,70,70	0
7	BMA	L	3	11/12	0.70	0.13	53,54,55,55	0
4	NAG	E	2	14/15	0.71	0.13	56,58,59,59	0
4	NAG	J	2	14/15	0.71	0.13	58,60,60,61	0
8	NAG	M	1	14/15	0.72	0.14	52,54,57,60	0
7	NAG	K	2	14/15	0.77	0.12	51,53,54,56	0
6	NAG	I	1	14/15	0.78	0.12	68,69,71,72	0
4	NAG	H	1	14/15	0.79	0.12	56,59,60,63	0
4	NAG	G	1	14/15	0.82	0.12	64,65,66,67	0
7	NAG	L	2	14/15	0.84	0.12	46,47,49,51	0
2	FUC	C	3	10/11	0.86	0.10	63,63,63,64	0
4	NAG	E	1	14/15	0.87	0.10	48,49,51,54	0
5	NAG	F	1	14/15	0.87	0.12	50,52,55,59	0
2	NAG	C	1	14/15	0.87	0.09	59,60,62,64	0
4	NAG	J	1	14/15	0.88	0.09	50,52,54,56	0
7	NAG	K	1	14/15	0.90	0.10	39,40,44,48	0
7	NAG	L	1	14/15	0.94	0.08	39,41,42,44	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain G:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain H:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain J:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain F:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain I:**

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain K:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain L:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
9	NAG	A	782	14/15	0.44	0.17	63,65,66,66	0
9	NAG	B	769	14/15	0.61	0.15	62,64,64,64	0
9	NAG	B	780	14/15	0.64	0.14	60,62,63,63	0
11	A3M	A	954	12/12	0.93	0.09	18,19,19,19	0
11	A3M	B	955	12/12	0.93	0.11	22,23,24,24	0
10	HG	A	950	1/1	0.98	0.05	56,56,56,56	0
10	HG	B	952	1/1	0.99	0.06	36,36,36,36	0
10	HG	B	953	1/1	0.99	0.05	60,60,60,60	0
10	HG	A	951	1/1	1.00	0.05	36,36,36,36	0

6.5 Other polymers [i](#)

There are no such residues in this entry.