



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 01:56 PM UTC

PDB ID : 5N1T / pdb_00005n1t
Title : Crystal structure of complex between flavocytochrome c and copper chaperone CopC from *T. paradoxus*
Authors : Osipov, E.M.; Lilina, A.V.; Tikhonova, T.V.; Tsallagov, S.I.; Popov, V.O.
Deposited on : 2017-02-06
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	429	
2	B	102	
3	M	160	
3	W	160	

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 5470 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called flavin-binding subunit of sulfide dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	393	Total	C	N	O	S	0	0	0
			2965	1872	524	557	12			


- Molecule 2 is a protein called Cytochrome C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	75	Total	C	N	O	S	0	0	0
			540	324	100	112	4			

- Molecule 3 is a protein called CopC.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	M	124	Total	C	N	O	0	0	0
			890	566	157	167			
3	W	124	Total	C	N	O	0	0	0
			915	578	164	173			

- Molecule 4 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 5 is HEME C (CCD ID: HEC) (formula: C₃₄H₃₄FeN₄O₄).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 6 is COPPER (II) ION (CCD ID: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	W	1	Total	Cu	0	0
			1	1		

- Molecule 7 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	W	1	Total	C	O	0	0
			6	3	3		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	39	Total	O	0	0
			39	39		
8	B	6	Total	O	0	0
			6	6		
8	M	4	Total	O	0	0
			4	4		
8	W	8	Total	O	0	0
			8	8		

- Molecule 1: flavin-binding subunit of sulfide dehydrogenase





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	39.53Å 138.40Å 155.81Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.60 30.00 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.7 (30.00-2.60) 99.7 (30.00-2.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.07 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.186 , 0.257 0.187 , 0.258	Depositor DCC
R_{free} test set	1386 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	51.7	Xtriage
Anisotropy	0.587	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 39.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5470	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CU, GOL, HEC, FAD, CSS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.92	0/3037	1.16	5/4145 (0.1%)
2	B	0.80	0/547	1.25	2/736 (0.3%)
3	M	0.98	0/911	1.16	0/1250
3	W	0.90	0/936	1.10	2/1281 (0.2%)
All	All	0.92	0/5431	1.16	9/7412 (0.1%)

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	25	ILE	N-CA-C	-6.17	102.91	108.95
1	A	194	PRO	CA-C-N	-6.12	113.46	119.64
1	A	194	PRO	C-N-CA	-6.12	113.46	119.64
3	W	66	THR	N-CA-C	5.82	117.81	110.24
1	A	379	VAL	N-CA-C	-5.63	100.34	108.89
2	B	53	MET	N-CA-C	5.56	117.34	111.28
1	A	262	THR	N-CA-C	5.08	119.58	113.17
1	A	243	MET	N-CA-C	5.05	116.87	111.36
3	W	106	VAL	N-CA-C	5.00	115.32	108.12

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2965	0	2833	35	0
2	B	540	0	504	1	0
3	M	890	0	830	3	0
3	W	915	0	863	10	0
4	A	53	0	30	2	0
5	B	43	0	30	0	0
6	W	1	0	0	0	0
7	W	6	0	8	0	0
8	A	39	0	0	1	0
8	B	6	0	0	0	0
8	M	4	0	0	0	0
8	W	8	0	0	0	0
All	All	5470	0	5098	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:193:CSS:SD	1:A:364:CYS:SG	2.74	0.85
1:A:258:VAL:HG11	1:A:279:LEU:HD13	1.62	0.81
1:A:321:ASP:O	1:A:330:LYS:HE2	1.91	0.70
1:A:156:MET:HG3	1:A:277:VAL:HG22	1.72	0.70
1:A:176:MET:HE3	1:A:276:VAL:HG23	1.80	0.64
3:W:62:LEU:HA	3:W:66:THR:HG21	1.79	0.64
1:A:40:GLY:O	1:A:63:LEU:HD11	1.97	0.63
3:W:19:LEU:HD23	3:W:95:LEU:HD12	1.79	0.63
1:A:62:THR:HG23	1:A:103:ARG:HB3	1.81	0.62
1:A:176:MET:HE2	1:A:275:ASP:HB2	1.81	0.61
1:A:360:TRP:HB2	1:A:380:TYR:HB2	1.83	0.60
1:A:342:CYS:O	1:A:346:VAL:HG13	2.02	0.59
1:A:146:GLU:HB3	1:A:257:ARG:HG3	1.85	0.59
1:A:77:ASN:HB3	1:A:203:LEU:HD12	1.85	0.58
3:W:76:LEU:HD13	3:W:89:LEU:HB3	1.84	0.58
3:W:8:ASP:HA	3:W:9:PRO:C	2.29	0.56
1:A:135:SER:HB2	1:A:319:ILE:HG13	1.89	0.54
1:A:183:VAL:HB	1:A:277:VAL:HB	1.89	0.54
1:A:74:PRO:HG3	4:A:501:FAD:C10	2.38	0.54
1:A:74:PRO:HG3	4:A:501:FAD:C4X	2.40	0.51
1:A:176:MET:HE1	1:A:182:VAL:HG22	1.95	0.48
3:M:76:LEU:HA	3:M:90:HIS:O	2.13	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:51:TYR:CZ	1:A:340:LYS:HE2	2.50	0.47
1:A:330:LYS:NZ	8:A:602:HOH:O	2.48	0.47
1:A:84:ARG:HB3	1:A:88:SER:OG	2.14	0.46
1:A:37:VAL:O	1:A:132:LEU:HD12	2.15	0.46
3:W:16:ALA:O	3:W:122:HIS:NE2	2.49	0.45
1:A:258:VAL:HG11	1:A:279:LEU:CD1	2.39	0.45
1:A:39:ILE:HD12	1:A:134:VAL:HG22	1.99	0.44
3:W:74:VAL:HG11	3:W:95:LEU:CD2	2.48	0.44
3:M:30:VAL:HG22	3:M:106:VAL:HG13	1.99	0.43
2:B:14:CYS:HB3	2:B:25:ILE:HD13	2.00	0.43
1:A:77:ASN:CB	1:A:203:LEU:HD12	2.48	0.43
1:A:255:VAL:HG22	1:A:265:VAL:CG1	2.48	0.43
1:A:290:ARG:HG3	1:A:295:THR:HG21	2.01	0.42
1:A:202:SER:HB3	1:A:368:VAL:O	2.19	0.42
1:A:226:PHE:O	1:A:229:GLN:HB2	2.19	0.42
1:A:132:LEU:HD23	1:A:316:ILE:HG12	2.01	0.42
1:A:42:GLY:HA2	1:A:70:TYR:CE2	2.55	0.42
1:A:129:TYR:CE2	1:A:132:LEU:HD22	2.55	0.41
3:W:56:ASN:O	3:W:57:THR:C	2.64	0.41
3:W:80:LEU:HD12	3:W:80:LEU:HA	1.87	0.41
3:M:42:LEU:HG	3:M:58:LEU:HD22	2.03	0.41
3:W:108:SER:HB3	3:W:110:ASP:OD1	2.22	0.40
1:A:228:LYS:NZ	1:A:366:SER:OG	2.44	0.40
3:W:55:LEU:HG	3:W:119:ALA:HB1	2.03	0.40
1:A:140:LEU:HB3	1:A:281:PRO:HG2	2.03	0.40
1:A:310:SER:C	1:A:312:GLN:H	2.29	0.40
1:A:363:THR:HG23	1:A:377:SER:HB2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	390/429 (91%)	375 (96%)	15 (4%)	0	100	100
2	B	73/102 (72%)	68 (93%)	5 (7%)	0	100	100
3	M	120/160 (75%)	113 (94%)	6 (5%)	1 (1%)	16	34
3	W	120/160 (75%)	113 (94%)	7 (6%)	0	100	100
All	All	703/851 (83%)	669 (95%)	33 (5%)	1 (0%)	48	70

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	M	19	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	301/333 (90%)	284 (94%)	17 (6%)	19	40
2	B	54/74 (73%)	52 (96%)	2 (4%)	30	57
3	M	86/124 (69%)	75 (87%)	11 (13%)	4	8
3	W	92/124 (74%)	86 (94%)	6 (6%)	15	35
All	All	533/655 (81%)	497 (93%)	36 (7%)	14	32

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	62	THR
1	A	92	ASP
1	A	95	GLN
1	A	119	VAL
1	A	127	ILE
1	A	146	GLU
1	A	168	LEU
1	A	169	LEU
1	A	171	ARG
1	A	238	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	243	MET
1	A	250	ILE
1	A	273	ARG
1	A	277	VAL
1	A	290	ARG
1	A	346	VAL
1	A	357	GLU
2	B	28	LEU
2	B	65	ILE
3	M	1	HIS
3	M	28	GLU
3	M	30	VAL
3	M	58	LEU
3	M	66	THR
3	M	74	VAL
3	M	84	SER
3	M	106	VAL
3	M	109	VAL
3	M	114	THR
3	M	122	HIS
3	W	30	VAL
3	W	33	ARG
3	W	52	LEU
3	W	64	VAL
3	W	79	ASP
3	W	83	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	210	HIS
1	A	305	GLN
2	B	10	ASN
2	B	75	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CSS	A	193	1	4,6,7	1.05	0	2,6,8	0.57	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSS	A	193	1	-	0/1/5/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	193	CSS	1	0

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	GOL	W	202	-	5,5,5	0.60	0	5,5,5	0.87	0
4	FAD	A	501	1	58,58,58	1.82	11 (18%)	85,89,89	1.72	21 (24%)
5	HEC	B	901	2	46,50,50	2.55	24 (52%)	58,82,82	2.30	20 (34%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	W	202	-	-	2/4/4/4	-
4	FAD	A	501	1	-	5/34/50/50	0/6/6/6
5	HEC	B	901	2	-	2/14/54/54	-

All (35) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	501	FAD	C9A-C5X	5.79	1.50	1.41
4	A	501	FAD	C5A-C4A	5.28	1.48	1.39
5	B	901	HEC	C2A-C3A	4.68	1.46	1.36
5	B	901	HEC	CHC-C4B	4.64	1.47	1.38
4	A	501	FAD	C8-C7	4.58	1.52	1.40
5	B	901	HEC	CAC-C3C	4.57	1.49	1.35
4	A	501	FAD	P-O3P	4.56	1.64	1.59
5	B	901	HEC	CHB-C4A	4.38	1.47	1.38
5	B	901	HEC	CHD-C4C	4.36	1.46	1.38
5	B	901	HEC	CAB-C3B	4.33	1.49	1.35
5	B	901	HEC	CHA-C1A	3.99	1.46	1.38
4	A	501	FAD	PA-O3P	3.81	1.63	1.59
5	B	901	HEC	CHC-C1C	3.72	1.47	1.39
5	B	901	HEC	CHD-C1D	3.68	1.47	1.39
5	B	901	HEC	C1D-ND	-3.40	1.33	1.39
5	B	901	HEC	C3D-C2D	3.34	1.47	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	901	HEC	C3B-C4B	3.29	1.51	1.46
5	B	901	HEC	C1C-C2C	3.02	1.50	1.43
5	B	901	HEC	C1A-NA	-3.00	1.34	1.39
5	B	901	HEC	C4D-ND	-2.98	1.34	1.39
5	B	901	HEC	C4A-NA	-2.89	1.34	1.39
4	A	501	FAD	C5A-C6A	2.70	1.48	1.41
4	A	501	FAD	C8A-N7A	2.64	1.36	1.31
5	B	901	HEC	C4C-NC	-2.54	1.34	1.39
5	B	901	HEC	CHA-C4D	2.53	1.45	1.39
5	B	901	HEC	C1A-C2A	2.42	1.49	1.45
5	B	901	HEC	C1D-C2D	2.35	1.48	1.43
5	B	901	HEC	CHB-C1B	2.33	1.44	1.39
5	B	901	HEC	C3C-C2C	2.33	1.49	1.41
4	A	501	FAD	C1'-C2'	-2.31	1.49	1.52
4	A	501	FAD	C4-N3	-2.29	1.34	1.38
5	B	901	HEC	C1B-NB	-2.10	1.35	1.39
4	A	501	FAD	C4A-N9A	-2.08	1.33	1.37
4	A	501	FAD	C5A-N7A	-2.05	1.35	1.39
5	B	901	HEC	C1C-NC	-2.05	1.35	1.39

All (41) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	901	HEC	C2A-C1A-NA	6.17	116.28	110.32
5	B	901	HEC	C3D-C4D-ND	5.65	116.43	110.15
5	B	901	HEC	C2B-C1B-NB	4.31	117.05	110.14
4	A	501	FAD	C5A-C4A-N3A	-4.14	121.01	126.72
4	A	501	FAD	N3A-C4A-N9A	3.98	133.94	127.17
4	A	501	FAD	N3A-C2A-N1A	-3.94	122.61	128.58
5	B	901	HEC	C1A-C2A-C3A	-3.89	101.99	107.11
4	A	501	FAD	C4'-C3'-C2'	-3.84	107.19	113.57
5	B	901	HEC	CBA-CAA-C2A	-3.78	102.07	112.53
5	B	901	HEC	CBD-CAD-C3D	-3.74	102.20	112.53
5	B	901	HEC	C3A-C4A-NA	3.68	116.44	109.64
4	A	501	FAD	C4A-N9A-C8A	3.66	109.58	105.74
5	B	901	HEC	C4D-C3D-C2D	-3.41	101.59	106.87
5	B	901	HEC	CMB-C2B-C1B	3.35	130.52	125.42
5	B	901	HEC	C1D-C2D-C3D	-3.35	102.98	106.82
5	B	901	HEC	C2C-C1C-NC	3.18	115.23	110.14
4	A	501	FAD	C2A-N3A-C4A	3.15	119.52	111.83
5	B	901	HEC	C4A-C3A-C2A	-3.11	102.35	106.97
4	A	501	FAD	C2A-N1A-C6A	3.04	123.73	118.73

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	901	HEC	C2D-C1D-ND	2.98	114.92	110.14
5	B	901	HEC	CHC-C4B-NB	-2.93	121.27	124.45
4	A	501	FAD	O3'-C3'-C4'	2.89	115.51	108.93
5	B	901	HEC	CHA-C4D-C3D	-2.85	119.06	125.30
4	A	501	FAD	O4-C4-C4X	-2.80	119.13	126.53
5	B	901	HEC	CHB-C1B-C2B	-2.64	119.76	127.43
5	B	901	HEC	CMD-C2D-C1D	2.62	129.41	125.42
4	A	501	FAD	C4X-C10-N1	-2.61	118.18	124.59
4	A	501	FAD	C4-N3-C2	-2.61	121.00	125.64
4	A	501	FAD	C4-C4X-N5	2.47	121.63	118.21
4	A	501	FAD	C4X-C4-N3	2.46	119.50	113.25
5	B	901	HEC	CHA-C1A-NA	-2.44	121.79	124.45
4	A	501	FAD	O2-C2-N1	-2.44	117.75	121.80
4	A	501	FAD	C5X-N5-C4X	2.43	122.02	118.09
4	A	501	FAD	C4A-C5A-N7A	-2.39	107.85	110.58
4	A	501	FAD	C10-N1-C2	2.29	121.81	116.85
5	B	901	HEC	CHB-C4A-C3A	-2.20	120.92	125.49
4	A	501	FAD	C2'-C1'-N10	2.19	120.54	110.20
4	A	501	FAD	O2P-P-O1P	2.15	122.43	112.44
5	B	901	HEC	CMC-C2C-C3C	2.12	131.54	126.55
4	A	501	FAD	N9A-C8A-N7A	-2.07	111.00	113.94
4	A	501	FAD	C6A-C5A-N7A	2.04	136.01	132.09

There are no chirality outliers.

All (9) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	501	FAD	O2'-C2'-C3'-O3'
7	W	202	GOL	O1-C1-C2-C3
4	A	501	FAD	O4B-C4B-C5B-O5B
4	A	501	FAD	C3B-C4B-C5B-O5B
7	W	202	GOL	O1-C1-C2-O2
5	B	901	HEC	CAD-CBD-CGD-O2D
4	A	501	FAD	C1'-C2'-C3'-O3'
5	B	901	HEC	CAD-CBD-CGD-O1D
4	A	501	FAD	O2'-C2'-C3'-C4'


There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	501	FAD	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	392/429 (91%)	-0.66	1 (0%) 90 88	32, 49, 70, 94	0
2	B	75/102 (73%)	-0.53	0 100 100	42, 56, 76, 83	0
3	M	124/160 (77%)	0.05	2 (1%) 70 66	40, 70, 93, 104	0
3	W	124/160 (77%)	-0.30	2 (1%) 70 66	42, 61, 88, 118	0
All	All	715/851 (84%)	-0.46	5 (0%) 84 82	32, 54, 85, 118	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	M	92	ASP	3.8
3	W	80	LEU	3.2
3	M	84	SER	2.8
1	A	384	ASN	2.6
3	W	17	ALA	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CSS	A	193	7/8	0.99	0.04	33,36,41,41	0

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.


6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
7	GOL	W	202	6/6	0.82	0.12	69,87,90,91	0
5	HEC	B	901	43/43	0.98	0.07	39,46,55,61	0
4	FAD	A	501	53/53	0.98	0.05	32,39,51,55	0
6	CU	W	201	1/1	0.99	0.03	83,83,83,83	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.