



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2026 – 11:00 PM UTC

PDB ID : 3N2W / pdb_00003n2w
Title : Crystal structure of the N-terminal beta-aminopeptidase BapA from *Sphingosinicella xenopeptidilytica*
Authors : Merz, T.; Heck, T.; Geueke, B.; Kohler, H.-P.; Gruetter, M.G.
Deposited on : 2010-05-19
Resolution : 1.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1756 (1.46-1.46)
Clashscore	190562	1795 (1.46-1.46)
Ramachandran outliers	187476	1776 (1.46-1.46)
Sidechain outliers	187428	1776 (1.46-1.46)
RSRZ outliers	180081	1756 (1.46-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	373	% 93% 6% .
1	B	373	4% 89% 8% ..
1	C	373	% 92% 6% .
1	D	373	2% 91% 5% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SO4	A	376	-	-	X	-
3	SO4	C	378	-	-	X	-

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 23824 atoms, of which 11136 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-peptidyl aminopeptidase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	367	Total	C	H	N	O	S	0	10	0
			5523	1711	2798	495	506	13			
1	B	365	Total	C	H	N	O	S	6	21	0
			5556	1731	2792	495	527	11			
1	C	367	Total	C	H	N	O	S	0	12	0
			5503	1709	2784	487	510	13			
1	D	361	Total	C	H	N	O	S	0	8	0
			5354	1666	2698	476	504	10			

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	0	0
			14	3	8	3		
2	A	1	Total	C	H	O	0	0
			14	3	8	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	H	O	0	0
			14	3	8	3		
2	C	1	Total	C	H	O	0	0
			14	3	8	3		
2	C	1	Total	C	H	O	0	0
			14	3	8	3		
2	C	1	Total	C	H	O	0	0
			14	3	8	3		
2	D	1	Total	C	H	O	0	0
			14	3	8	3		
2	D	1	Total	C	H	O	0	0
			14	3	8	3		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	1	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	D	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	456	Total	O	0	0
			456	456		
4	B	430	Total	O	0	0
			430	430		
4	C	438	Total	O	0	0
			438	438		
4	D	417	Total	O	0	0
			417	417		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Beta-peptidyl aminopeptidase




- Molecule 1: Beta-peptidyl aminopeptidase



- Molecule 1: Beta-peptidyl aminopeptidase



- Molecule 1: Beta-peptidyl aminopeptidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	87.36Å 96.72Å 101.42Å 90.00° 108.23° 90.00°	Depositor
Resolution (Å)	48.81 – 1.45 48.81 – 1.45	Depositor EDS
% Data completeness (in resolution range)	98.5 (48.81-1.45) 98.5 (48.81-1.45)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$	-	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.6.1_357)	Depositor
R, R_{free}	0.131 , 0.150 0.130 , 0.148	Depositor DCC
R_{free} test set	6965 reflections (2.50%)	wwPDB-VP
Wilson B-factor (Å ²)	(Not available)	Xtrriage
Anisotropy	(Not available)	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 47.4	EDS
L-test for twinning ¹	$\langle L \rangle =$ (Not available), $\langle L^2 \rangle =$ (Not available)	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	23824	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *(Not available)*

¹Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.70	0/2805	0.85	1/3802 (0.0%)
1	B	0.66	0/2838	0.80	0/3855
1	C	0.65	0/2802	0.81	0/3802
1	D	0.67	0/2727	0.83	0/3703
All	All	0.67	0/11172	0.82	1/15162 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	281	GLY	N-CA-C	5.31	121.01	114.69

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2725	2798	2788	26	0
1	B	2764	2792	2792	40	0
1	C	2719	2784	2774	28	0
1	D	2656	2698	2676	19	0
2	A	12	16	16	1	0
2	B	6	8	8	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	18	24	24	0	0
2	D	12	16	16	1	0
3	A	10	0	0	2	0
3	C	15	0	0	2	0
3	D	10	0	0	1	0
4	A	456	0	0	6	0
4	B	430	0	0	2	0
4	C	438	0	0	5	0
4	D	417	0	0	3	0
All	All	12688	11136	11094	102	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (102) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:74:ASN:HD21	1:C:268:ARG:HH21	1.21	0.88
1:A:268:ARG:HH21	1:C:74:ASN:HD21	1.18	0.88
1:B:229:TRP:CD2	1:B:238[A]:ALA:HB2	2.10	0.87
1:B:268:ARG:HH21	1:D:74:ASN:HD21	1.17	0.84
1:B:74:ASN:HD21	1:D:268:ARG:HH21	1.22	0.84
1:C:313:ILE:HG23	3:C:378:SO4:O4	1.80	0.82
1:D:125:SER:HA	2:D:375:GOL:H11	1.63	0.78
1:C:190[B]:VAL:HG12	1:C:197:ARG:CG	2.14	0.76
1:B:229:TRP:CE2	1:B:238[A]:ALA:HB2	2.20	0.75
1:A:302:PRO:HG3	1:A:308[B]:ARG:HE	1.49	0.75
1:C:190[B]:VAL:CG1	1:C:197:ARG:HG2	2.18	0.73
1:C:190[B]:VAL:HG12	1:C:197:ARG:HD3	1.70	0.72
1:C:190[B]:VAL:HG12	1:C:197:ARG:HG2	1.73	0.70
1:C:190[B]:VAL:HG12	1:C:197:ARG:CD	2.21	0.69
1:A:148:THR:H	1:A:151:HIS:CD2	2.12	0.68
1:B:148:THR:H	1:B:151:HIS:CD2	2.12	0.68
1:B:231[A]:GLU:HG2	1:B:235[A]:ILE:HD13	1.76	0.68
1:A:148:THR:H	1:A:151:HIS:HD2	1.43	0.66
1:B:232[B]:VAL:HG22	1:B:237[B]:ALA:HB2	1.78	0.65
1:B:235[A]:ILE:N	1:B:242[A]:ALA:HA	2.12	0.64
1:B:148:THR:H	1:B:151:HIS:HD2	1.45	0.63
1:A:351:ASN:ND2	4:A:931:HOH:O	2.31	0.63
1:B:237[A]:ALA:O	1:B:238[A]:ALA:HB3	1.98	0.62
1:C:110[B]:MET:HG2	4:C:587:HOH:O	1.99	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:369:ARG:HE	1:C:369:ARG:NH1	1.98	0.62
1:A:285:GLY:HA3	3:A:376:SO4:O2	1.98	0.62
1:A:306[B]:LYS:NZ	4:A:1263:HOH:O	2.32	0.61
1:B:236[B]:VAL:HB	1:B:241[B]:ASP:HB3	1.83	0.60
1:C:347[A]:THR:HG23	4:C:610:HOH:O	2.02	0.60
1:B:232[B]:VAL:O	1:B:233[B]:ASN:HB2	2.02	0.59
1:C:190[B]:VAL:CG1	1:C:197:ARG:HD3	2.33	0.58
1:D:230:PRO:HG3	4:D:774:HOH:O	2.03	0.57
1:B:241[A]:ASP:O	1:B:242[A]:ALA:HB3	2.03	0.57
1:B:233[A]:ASN:H	1:B:234[A]:GLY:HA3	1.69	0.57
1:C:190[B]:VAL:CG1	1:C:197:ARG:CD	2.82	0.56
1:D:285:GLY:HA3	3:D:376:SO4:O4	2.06	0.56
1:C:261:LEU:CD1	1:C:269[B]:MET:SD	2.95	0.55
1:B:232[B]:VAL:O	1:B:233[B]:ASN:CB	2.55	0.54
1:D:232:VAL:HG13	4:D:1194:HOH:O	2.05	0.54
1:D:179[B]:THR:OG1	1:D:211:ARG:HB2	2.08	0.54
1:A:364[A]:ARG:HD3	4:A:1436:HOH:O	2.07	0.53
4:A:1283:HOH:O	1:C:326[B]:ARG:NH1	2.42	0.53
1:A:306[B]:LYS:HA	1:A:306[B]:LYS:CE	2.39	0.52
1:D:160[A]:LYS:HE2	4:D:455:HOH:O	2.09	0.52
1:A:138[B]:ARG:HB3	1:A:138[B]:ARG:NH1	2.24	0.52
1:B:300:VAL:HG21	1:B:308:ARG:NH2	2.24	0.52
1:C:369:ARG:NH1	4:C:539:HOH:O	2.42	0.52
1:B:229:TRP:CG	1:B:238[A]:ALA:HB2	2.45	0.52
1:C:364:ARG:HD3	4:C:1347:HOH:O	2.10	0.52
1:B:232[B]:VAL:HG22	1:B:237[B]:ALA:CB	2.39	0.51
1:B:241[A]:ASP:O	1:B:242[A]:ALA:CB	2.58	0.51
1:B:237[A]:ALA:O	1:B:238[A]:ALA:CB	2.59	0.50
1:D:369:ARG:C	1:D:371:PRO:HD3	2.37	0.49
1:D:232:VAL:HG12	1:D:233:ASN:HD22	1.79	0.48
1:B:235[A]:ILE:HB	1:B:242[A]:ALA:H	1.78	0.48
1:C:110[B]:MET:SD	1:C:129:PRO:HB3	2.54	0.48
1:A:74:ASN:ND2	1:C:272:ARG:HH12	2.12	0.47
1:C:317:ASP:OD2	3:C:378:SO4:S	2.72	0.47
1:A:285:GLY:CA	3:A:376:SO4:O2	2.63	0.47
1:C:306:LYS:HE3	1:C:307:PRO:O	2.15	0.47
1:A:138[B]:ARG:HB3	1:A:138[B]:ARG:HH11	1.80	0.47
1:A:302:PRO:CG	1:A:308[B]:ARG:HE	2.22	0.47
1:B:121:GLU:HG3	4:B:1197:HOH:O	2.14	0.47
1:A:300:VAL:HG21	1:A:308[A]:ARG:NH2	2.29	0.46
1:C:143:PHE:CZ	1:D:127[B]:LEU:HD21	2.50	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:179[A]:THR:HG23	1:B:211:ARG:H	1.81	0.46
1:B:272:ARG:HH12	1:D:74:ASN:ND2	2.13	0.46
1:B:369:ARG:C	1:B:371:PRO:HD3	2.39	0.46
1:C:190[B]:VAL:CG1	1:C:197:ARG:CG	2.82	0.46
1:A:302:PRO:HG3	1:A:308[B]:ARG:NE	2.25	0.45
1:A:272:ARG:HH12	1:C:74:ASN:ND2	2.15	0.45
1:C:364:ARG:NH1	4:C:1279:HOH:O	2.50	0.45
1:D:31:HIS:HD2	1:D:53:THR:OG1	1.99	0.44
1:D:370:PHE:N	1:D:371:PRO:HD3	2.31	0.44
1:C:300:VAL:HG21	1:C:308:ARG:NH2	2.33	0.44
1:B:74:ASN:ND2	1:D:272:ARG:HH12	2.16	0.43
1:D:369:ARG:O	1:D:369:ARG:HD2	2.17	0.43
1:B:234[A]:GLY:O	1:B:235[A]:ILE:C	2.61	0.43
1:B:232[A]:VAL:HG23	1:B:236[A]:VAL:HG21	2.01	0.43
1:B:110[B]:MET:HE3	4:B:419:HOH:O	2.19	0.43
1:B:179[A]:THR:OG1	1:B:211:ARG:HB2	2.19	0.43
1:A:364[B]:ARG:NH2	4:A:1281:HOH:O	2.52	0.43
1:C:369:ARG:C	1:C:371:PRO:HD3	2.44	0.43
1:B:147:LEU:HA	1:B:151:HIS:HD2	1.84	0.42
1:A:110[B]:MET:HG2	4:A:525:HOH:O	2.19	0.42
1:C:190[B]:VAL:HG13	1:C:197:ARG:HG2	2.00	0.42
1:D:211:ARG:HD3	1:D:229:TRP:CH2	2.54	0.42
1:A:127:LEU:HD21	1:B:143:PHE:CZ	2.55	0.42
1:B:232[A]:VAL:HG23	1:B:236[A]:VAL:CG2	2.50	0.42
1:B:242[A]:ALA:O	1:B:243:GLY:O	2.37	0.42
1:B:232[B]:VAL:O	1:B:232[B]:VAL:HG23	2.20	0.42
1:A:326[A]:ARG:CZ	1:A:330:GLN:HE22	2.33	0.42
1:D:105:LEU:C	1:D:105:LEU:HD13	2.45	0.42
1:A:306[B]:LYS:HA	1:A:306[B]:LYS:HE3	2.01	0.41
1:B:235[A]:ILE:O	1:B:242[A]:ALA:N	2.54	0.41
1:B:110[B]:MET:HE3	1:B:110[B]:MET:HB3	1.82	0.41
1:B:136:ASP:O	1:B:140:ASN:HB3	2.20	0.41
1:A:121:GLU:H	1:A:121:GLU:CD	2.28	0.41
2:A:374:GOL:H31	1:D:262:MET:HE1	2.02	0.41
1:B:364:ARG:HD2	1:B:364:ARG:C	2.46	0.41
1:B:125:SER:HB3	1:B:303:LEU:HD11	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	374/373 (100%)	369 (99%)	5 (1%)	0	100	100
1	B	382/373 (102%)	370 (97%)	6 (2%)	6 (2%)	7	1
1	C	375/373 (100%)	371 (99%)	4 (1%)	0	100	100
1	D	365/373 (98%)	360 (99%)	5 (1%)	0	100	100
All	All	1496/1492 (100%)	1470 (98%)	20 (1%)	6 (0%)	43	11

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	233[A]	ASN
1	B	233[B]	ASN
1	B	242[A]	ALA
1	B	242[B]	ALA
1	B	240[A]	PRO
1	B	240[B]	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	282/277 (102%)	281 (100%)	1 (0%)	84	70
1	B	285/277 (103%)	285 (100%)	0	100	100
1	C	283/277 (102%)	282 (100%)	1 (0%)	84	70
1	D	275/277 (99%)	271 (98%)	4 (2%)	57	25

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	1125/1108 (102%)	1119 (100%)	6 (0%)	81 64

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	179	THR
1	C	179	THR
1	D	179[A]	THR
1	D	179[B]	THR
1	D	235	ILE
1	D	369	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	74	ASN
1	A	151	HIS
1	A	330	GLN
1	A	339	GLN
1	B	74	ASN
1	B	151	HIS
1	B	220	GLN
1	B	339	GLN
1	C	74	ASN
1	C	339	GLN
1	C	351	ASN
1	D	31	HIS
1	D	74	ASN
1	D	233	ASN
1	D	339	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	C	377	-	4,4,4	0.27	0	6,6,6	0.73	0
2	GOL	D	374	-	5,5,5	0.44	0	5,5,5	0.88	0
3	SO4	C	379	-	4,4,4	0.33	0	6,6,6	0.46	0
2	GOL	C	374	-	5,5,5	0.52	0	5,5,5	0.85	0
3	SO4	D	377	-	4,4,4	0.25	0	6,6,6	0.08	0
3	SO4	D	376	-	4,4,4	0.31	0	6,6,6	0.50	0
3	SO4	A	377	-	4,4,4	0.26	0	6,6,6	0.13	0
2	GOL	D	375	-	5,5,5	0.40	0	5,5,5	0.39	0
3	SO4	C	378	-	4,4,4	0.29	0	6,6,6	0.43	0
2	GOL	C	376	-	5,5,5	0.48	0	5,5,5	0.17	0
2	GOL	A	374	-	5,5,5	0.58	0	5,5,5	0.90	0
2	GOL	A	375	-	5,5,5	0.33	0	5,5,5	0.36	0
2	GOL	C	375	-	5,5,5	0.36	0	5,5,5	0.44	0
3	SO4	A	376	-	4,4,4	0.27	0	6,6,6	0.29	0
2	GOL	B	374	-	5,5,5	0.48	0	5,5,5	0.98	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	D	374	-	-	0/4/4/4	-
2	GOL	C	374	-	-	0/4/4/4	-
2	GOL	D	375	-	-	1/4/4/4	-
2	GOL	C	376	-	-	2/4/4/4	-
2	GOL	A	374	-	-	0/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	375	-	-	4/4/4/4	-
2	GOL	C	375	-	-	2/4/4/4	-
2	GOL	B	374	-	-	1/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	375	GOL	O1-C1-C2-C3
2	A	375	GOL	C1-C2-C3-O3
2	C	375	GOL	C1-C2-C3-O3
2	A	375	GOL	O1-C1-C2-O2
2	C	375	GOL	O2-C2-C3-O3
2	C	376	GOL	C1-C2-C3-O3
2	A	375	GOL	O2-C2-C3-O3
2	C	376	GOL	O2-C2-C3-O3
2	D	375	GOL	O1-C1-C2-O2
2	B	374	GOL	C1-C2-C3-O3

There are no ring outliers.

5 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	376	SO4	1	0
2	D	375	GOL	1	0
3	C	378	SO4	2	0
2	A	374	GOL	1	0
3	A	376	SO4	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	367/373 (98%)	-0.64	5 (1%) 73 77	6, 14, 29, 52	17 (4%)
1	B	365/373 (97%)	-0.46	15 (4%) 41 44	7, 15, 29, 71	31 (8%)
1	C	367/373 (98%)	-0.65	5 (1%) 73 77	6, 14, 31, 50	20 (5%)
1	D	361/373 (96%)	-0.55	8 (2%) 62 65	7, 14, 32, 61	18 (4%)
All	All	1460/1492 (97%)	-0.58	33 (2%) 61 63	6, 14, 32, 71	86 (5%)

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	242[A]	ALA	8.3
1	D	234	GLY	6.4
1	B	240[A]	PRO	6.2
1	B	239[A]	GLY	6.1
1	D	239	GLY	5.8
1	B	236[A]	VAL	5.7
1	B	241[A]	ASP	5.6
1	B	243	GLY	5.4
1	B	233[A]	ASN	4.8
1	D	233	ASN	4.7
1	B	232[A]	VAL	4.2
1	B	238[A]	ALA	4.2
1	A	124	PHE	3.6
1	A	371	PRO	3.5
1	D	232	VAL	3.1
1	B	194	GLY	3.0
1	D	371	PRO	2.9
1	C	371	PRO	2.8
1	C	124	PHE	2.8
1	D	235	ILE	2.7
1	B	193	ALA	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	124	PHE	2.5
1	B	235[A]	ILE	2.4
1	D	238	ALA	2.3
1	A	369	ARG	2.2
1	B	121	GLU	2.2
1	C	245	PRO	2.1
1	C	195	ASP	2.1
1	A	195	ASP	2.1
1	C	193	ALA	2.1
1	B	371	PRO	2.1
1	D	120	GLU	2.0
1	A	1	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	A	377	5/5	0.66	0.30	27,37,49,57	5
2	GOL	D	375	6/6	0.73	0.19	31,48,61,61	0
2	GOL	C	376	6/6	0.74	0.20	38,48,61,62	0
2	GOL	C	375	6/6	0.77	0.17	32,44,56,56	0
2	GOL	A	375	6/6	0.78	0.18	30,44,58,59	0
3	SO4	C	379	5/5	0.83	0.21	15,26,31,45	5
2	GOL	A	374	6/6	0.94	0.08	18,22,24,25	0
3	SO4	C	378	5/5	0.95	0.20	20,24,29,38	5
3	SO4	D	377	5/5	0.95	0.10	19,24,37,38	5
2	GOL	D	374	6/6	0.96	0.06	16,20,23,23	0
2	GOL	B	374	6/6	0.97	0.05	16,20,22,24	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GOL	C	374	6/6	0.97	0.06	17,21,23,24	0
3	SO4	C	377	5/5	0.98	0.14	12,24,31,33	5
3	SO4	D	376	5/5	0.98	0.17	11,24,27,34	5
3	SO4	A	376	5/5	0.98	0.15	11,19,28,37	5

6.5 Other polymers [i](#)

There are no such residues in this entry.