



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 03:23 AM UTC

PDB ID : 4N2R / pdb_00004n2r
Title : Crystal Structure of the alpha-L-arabinofuranosidase UmAbf62A from
Ustilago maydis in complex with L-arabinofuranose
Authors : Siguier, B.; Dumon, C.; Mourey, L.; Tranier, S.
Deposited on : 2013-10-06
Resolution : 1.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1216 (1.20-1.20)
Clashscore	190562	1265 (1.20-1.20)
Ramachandran outliers	187476	1226 (1.20-1.20)
Sidechain outliers	187428	1226 (1.20-1.20)
RSRZ outliers	180081	1214 (1.20-1.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	334	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	TRS	A	404	-	X	X	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3025 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called alpha-L-arabinofuranosidase UmAbf62A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	306	Total	C	N	O	S	0	26	0
			2558	1638	426	484	10			


There are 23 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	312	GLY	-	expression tag	UNP Q4P6F4
A	313	LEU	-	expression tag	UNP Q4P6F4
A	314	GLU	-	expression tag	UNP Q4P6F4
A	315	GLN	-	expression tag	UNP Q4P6F4
A	316	LYS	-	expression tag	UNP Q4P6F4
A	317	LEU	-	expression tag	UNP Q4P6F4
A	318	ILE	-	expression tag	UNP Q4P6F4
A	319	SER	-	expression tag	UNP Q4P6F4
A	320	GLU	-	expression tag	UNP Q4P6F4
A	321	GLU	-	expression tag	UNP Q4P6F4
A	322	ASP	-	expression tag	UNP Q4P6F4
A	323	LEU	-	expression tag	UNP Q4P6F4
A	324	ASN	-	expression tag	UNP Q4P6F4
A	325	SER	-	expression tag	UNP Q4P6F4
A	326	ALA	-	expression tag	UNP Q4P6F4
A	327	VAL	-	expression tag	UNP Q4P6F4
A	328	ASP	-	expression tag	UNP Q4P6F4
A	329	HIS	-	expression tag	UNP Q4P6F4
A	330	HIS	-	expression tag	UNP Q4P6F4
A	331	HIS	-	expression tag	UNP Q4P6F4
A	332	HIS	-	expression tag	UNP Q4P6F4
A	333	HIS	-	expression tag	UNP Q4P6F4
A	334	HIS	-	expression tag	UNP Q4P6F4

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ca	0	0
			1	1		

- Molecule 3 is beta-L-arabinofuranose (CCD ID: FUB) (formula: C₅H₁₀O₅).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	1
			10	5	5		

- Molecule 4 is alpha-L-arabinofuranose (CCD ID: AHR) (formula: C₅H₁₀O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	1
			10	5	5		

- Molecule 5 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: C₄H₁₂NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			8	4	1	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	438	Total	O	0	1
			438	438		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: alpha-L-arabinofuranosidase UmAbf62A



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	60.08Å 66.07Å 68.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.42 – 1.20 47.42 – 1.20	Depositor EDS
% Data completeness (in resolution range)	95.8 (47.42-1.20) 95.8 (47.42-1.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.03	Depositor
$\langle I/\sigma(I) \rangle$ ¹	9.34 (at 1.20Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.100 , 0.130 0.101 , 0.131	Depositor DCC
R_{free} test set	4098 reflections (4.80%)	wwPDB-VP
Wilson B-factor (Å ²)	7.8	Xtriage
Anisotropy	0.339	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 44.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.017 for -h,l,k	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	3025	wwPDB-VP
Average B, all atoms (Å ²)	11.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FUB, TRS, AHR, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.18	12/2670 (0.4%)	1.03	8/3638 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	151	ARG	NE-CZ	6.91	1.40	1.33
1	A	229[A]	SER	N-CA	6.56	1.54	1.46
1	A	229[B]	SER	N-CA	6.56	1.54	1.46
1	A	268	ASN	CG-OD1	6.15	1.35	1.23
1	A	257	ASN	C-O	-6.12	1.16	1.24
1	A	195	GLU	CA-CB	6.09	1.64	1.53
1	A	104	TRP	C-O	-5.87	1.16	1.24
1	A	312	GLY	N-CA	5.83	1.51	1.45
1	A	230	LEU	C-O	-5.49	1.17	1.24
1	A	151	ARG	CD-NE	5.27	1.53	1.46
1	A	237	GLN	CG-CD	-5.22	1.39	1.52
1	A	65	PRO	C-O	5.11	1.29	1.23

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	237	GLN	CB-CG-CD	5.62	122.15	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	314	GLU	CA-C-O	-5.57	111.33	120.80
1	A	151	ARG	NE-CZ-NH1	5.46	126.96	121.50
1	A	263	ASP	CA-CB-CG	5.40	118.00	112.60
1	A	229[A]	SER	O-C-N	5.22	129.26	123.52
1	A	229[B]	SER	O-C-N	5.22	129.26	123.52
1	A	70	ALA	CA-C-N	5.13	130.12	121.14
1	A	70	ALA	C-N-CA	5.13	130.12	121.14

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	266	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2558	0	2491	41	0
2	A	1	0	0	0	0
3	A	10	0	0	0	0
4	A	10	0	0	0	0
5	A	8	0	10	9	0
6	A	438	0	0	13	0
All	All	3025	0	2501	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:272:THR:C	1:A:273[B]:MET:CA	2.44	0.89
1:A:273[B]:MET:CA	1:A:274:THR:N	2.43	0.81
1:A:104:TRP:N	6:A:1031:HOH:O	2.17	0.77

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:273[B]:MET:CE	6:A:808:HOH:O	2.33	0.76
1:A:180:THR:OG1	5:A:404:TRS:H31	1.87	0.75
1:A:180:THR:H	5:A:404:TRS:H22	1.51	0.75
1:A:76[B]:THR:HG21	6:A:984:HOH:O	1.91	0.70
1:A:53:HIS:HE1	6:A:876:HOH:O	1.75	0.69
1:A:180:THR:N	5:A:404:TRS:H22	2.08	0.68
1:A:152:LYS:HE3	6:A:1053[A]:HOH:O	1.94	0.67
1:A:152:LYS:HE3	1:A:154[A]:TYR:OH	1.94	0.67
1:A:181:ALA:H	5:A:404:TRS:C2	2.09	0.66
1:A:180:THR:H	5:A:404:TRS:H31	1.61	0.65
1:A:53:HIS:HD2	1:A:54:ASP:O	1.81	0.64
1:A:81[A]:THR:H	1:A:121:ASN:HD21	1.46	0.63
1:A:81[B]:THR:H	1:A:121:ASN:HD21	1.45	0.63
1:A:266:ARG:HD3	1:A:273[B]:MET:HE3	1.79	0.63
1:A:181:ALA:H	5:A:404:TRS:H21	1.64	0.61
1:A:151:ARG:NH2	6:A:1053[A]:HOH:O	2.37	0.58
1:A:104:TRP:O	6:A:1031:HOH:O	2.17	0.57
1:A:81[A]:THR:H	1:A:121:ASN:ND2	2.03	0.56
1:A:81[B]:THR:H	1:A:121:ASN:ND2	2.03	0.56
1:A:273[B]:MET:HE1	6:A:808:HOH:O	2.02	0.56
1:A:276:ASP:OD2	1:A:279[A]:ARG:NE	2.20	0.55
1:A:80:MET:HA	1:A:121:ASN:HD21	1.72	0.55
1:A:17[B]:ARG:HG2	6:A:1005:HOH:O	2.08	0.54
1:A:93[A]:THR:O	1:A:96[A]:LYS:HD3	2.08	0.54
1:A:266:ARG:CZ	1:A:273[B]:MET:HE1	2.38	0.54
1:A:180:THR:H	5:A:404:TRS:C3	2.21	0.52
1:A:291:LYS:HD3	1:A:300:PRO:HG3	1.92	0.52
1:A:229[B]:SER:OG	1:A:231:ASP:OD1	2.27	0.52
1:A:151:ARG:CZ	6:A:1053[A]:HOH:O	2.60	0.49
1:A:76[B]:THR:CG2	6:A:984:HOH:O	2.57	0.46
1:A:229[B]:SER:HB2	6:A:1041:HOH:O	2.16	0.45
1:A:181:ALA:CB	5:A:404:TRS:H21	2.47	0.45
1:A:53:HIS:CD2	1:A:54:ASP:O	2.67	0.45
1:A:225:PHE:CE2	1:A:236[B]:VAL:HG22	2.55	0.42
1:A:68:HIS:HD2	6:A:996:HOH:O	2.03	0.42
1:A:266:ARG:HD3	1:A:273[B]:MET:CE	2.47	0.41
1:A:181:ALA:N	5:A:404:TRS:H21	2.32	0.41
1:A:154[A]:TYR:OH	1:A:231:ASP:HB3	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	331/334 (99%)	316 (96%)	14 (4%)	1 (0%)	36	14

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	261	HIS

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	279/279 (100%)	276 (99%)	3 (1%)	65	31

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	279[A]	ARG
1	A	279[B]	ARG
1	A	291	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	53	HIS
1	A	68	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	121	ASN
1	A	207	GLN
1	A	237	GLN
1	A	281	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	TRS	A	404	-	7,7,7	2.79	2 (28%)	9,9,9	2.94	5 (55%)
3	FUB	A	402[A]	-	10,10,10	1.80	2 (20%)	13,14,14	2.56	5 (38%)
4	AHR	A	403[B]	-	10,10,10	0.69	0	13,14,14	2.71	5 (38%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	TRS	A	404	-	-	7/9/9/9	-
3	FUB	A	402[A]	-	-	0/2/18/18	0/1/1/1
4	AHR	A	403[B]	-	-	0/2/18/18	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	404	TRS	O2-C2	-6.68	1.21	1.42
3	A	402[A]	FUB	O1-C1	3.95	1.52	1.39
3	A	402[A]	FUB	C1-C2	2.33	1.55	1.52
5	A	404	TRS	O3-C3	-2.24	1.35	1.42

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	403[B]	AHR	O4-C1-C2	7.05	114.45	104.67
3	A	402[A]	FUB	O1-C1-O4	5.79	118.50	111.12
5	A	404	TRS	C3-C-C2	-4.84	97.77	110.66
4	A	403[B]	AHR	O1-C1-O4	-4.71	105.12	111.12
3	A	402[A]	FUB	O4-C1-C2	-4.16	98.90	104.67
3	A	402[A]	FUB	O4-C4-C5	4.06	117.81	109.22
5	A	404	TRS	O1-C1-C	3.77	121.39	110.88
5	A	404	TRS	O2-C2-C	3.63	120.99	110.88
5	A	404	TRS	O3-C3-C	-3.47	101.19	110.88
5	A	404	TRS	C2-C-N	3.23	116.41	108.17
4	A	403[B]	AHR	O2-C2-C3	2.82	120.85	111.82
3	A	402[A]	FUB	O2-C2-C3	2.76	120.65	111.82
4	A	403[B]	AHR	O4-C4-C3	2.36	109.83	105.15
4	A	403[B]	AHR	O5-C5-C4	-2.22	103.79	111.33
3	A	402[A]	FUB	O5-C5-C4	-2.02	104.44	111.33

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	404	TRS	C1-C-C2-O2
5	A	404	TRS	C3-C-C2-O2
5	A	404	TRS	N-C-C3-O3
5	A	404	TRS	C3-C-C1-O1
5	A	404	TRS	C1-C-C3-O3
5	A	404	TRS	C2-C-C3-O3
5	A	404	TRS	N-C-C2-O2

There are no ring outliers.

1 monomer is involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	404	TRS	9	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	306/334 (91%)	-0.44	10 (3%)	49 49	2, 7, 20, 55	26 (8%)

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	311	LEU	4.0
1	A	104	TRP	3.9
1	A	310	GLY	3.7
1	A	292	VAL	3.3
1	A	309	TYR	3.3
1	A	312	GLY	2.8
1	A	294	SER	2.7
1	A	9	SER	2.4
1	A	313	LEU	2.4
1	A	293	PRO	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	AHR	A	403[B]	10/10	0.87	0.12	14,22,26,27	10
5	TRS	A	404	8/8	0.93	0.10	4,10,29,36	0
3	FUB	A	402[A]	10/10	0.94	0.10	6,11,16,16	10
2	CA	A	401	1/1	1.00	0.01	4,4,4,4	0

6.5 Other polymers [i](#)

There are no such residues in this entry.