



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 05:41 PM UTC

PDB ID : 7N2R / pdb_00007n2r
Title : AS4.3-PRPF3-HLA*B27
Authors : Yang, X.; Jude, K.M.; Garcia, K.C.
Deposited on : 2021-05-29
Resolution : 2.28 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.28 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	9078 (2.30-2.26)
Clashscore	190562	9802 (2.30-2.26)
Ramachandran outliers	187476	9690 (2.30-2.26)
Sidechain outliers	187428	9691 (2.30-2.26)
RSRZ outliers	180081	9085 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	D	204	7% 87% 8% .
2	F	242	4% 81% 18%
3	A	278	10% 83% 10% 7%
4	B	100	8% 87% 13%
5	C	9	89% 11%

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 6612 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called AS4.3 T cell receptor alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	D	195	Total	C	N	O	S	0	2	0
			1512	944	251	310	7			

- Molecule 2 is a protein called AS4.3 T cell receptor beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	F	241	Total	C	N	O	S	0	0	0
			1914	1209	332	367	6			

- Molecule 3 is a protein called Human leukocyte antigen (HLA) B27.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	259	Total	C	N	O	S	0	1	0
			2119	1324	386	403	6			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	67	SER	CYS	conflict	UNP A3F718

- Molecule 4 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	B	100	Total	C	N	O	S	0	0	0
			837	533	141	159	4			


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	initiating methionine	UNP P61769

- Molecule 5 is a protein called Pre-mRNA Processing Factor 3.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	C	9	Total	C	N	O	0	0	0
			69	45	13	11			

- Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	D	1	Total	C	N	O	0	0
			14	8	1	5		
6	D	1	Total	C	N	O	0	0
			14	8	1	5		
6	D	1	Total	C	N	O	0	0
			14	8	1	5		
6	D	1	Total	C	N	O	0	0
			14	8	1	5		
6	F	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 7 is ASPARTIC ACID (CCD ID: ASP) (formula: $C_4H_7NO_4$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	F	1	Total	C	N	O	0	0
			9	4	1	4		

- Molecule 8 is SULFATE ION (CCD ID: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	F	1	Total 5	O 4	S 1	0	0
8	F	1	Total 5	O 4	S 1	0	0
8	A	1	Total 5	O 4	S 1	0	0

- Molecule 9 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	D	17	Total	O	0	0
			17	17		
10	F	17	Total	O	0	0
			17	17		
10	A	26	Total	O	0	0
			26	26		
10	B	3	Total	O	0	0
			3	3		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: AS4.3 T cell receptor alpha chain




- Molecule 2: AS4.3 T cell receptor beta chain



- Molecule 3: Human leukocyte antigen (HLA) B27




- Molecule 4: Beta-2-microglobulin



- Molecule 5: Pre-mRNA Processing Factor 3

Chain C:  89% 11%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	54.68Å 94.42Å 178.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.26 – 2.28 40.26 – 2.28	Depositor EDS
% Data completeness (in resolution range)	99.2 (40.26-2.28) 99.2 (40.26-2.28)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.08 (at 2.29Å)	Xtriage
Refinement program	PHENIX 1.19_4080	Depositor
R, R_{free}	0.248 , 0.288 0.248 , 0.288	Depositor DCC
R_{free} test set	2000 reflections (4.67%)	wwPDB-VP
Wilson B-factor (Å ²)	54.3	Xtriage
Anisotropy	0.484	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 54.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6612	wwPDB-VP
Average B, all atoms (Å ²)	86.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, EDO, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	D	0.15	0/1544	0.34	0/2095
2	F	0.14	0/1965	0.31	0/2679
3	A	0.14	0/2178	0.31	0/2958
4	B	0.12	0/860	0.29	0/1162
5	C	0.19	0/69	0.34	0/91
All	All	0.14	0/6616	0.31	0/8985

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	1512	0	1436	10	0
2	F	1914	0	1818	24	0
3	A	2119	0	1997	16	0
4	B	837	0	803	8	0
5	C	69	0	85	1	0
6	D	56	0	52	0	0
6	F	14	0	13	0	0
7	F	9	0	3	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	A	5	0	0	0	0
8	F	10	0	0	0	0
9	A	4	0	6	0	0
10	A	26	0	0	0	0
10	B	3	0	0	0	0
10	D	17	0	0	0	0
10	F	17	0	0	0	0
All	All	6612	0	6213	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:46:ILE:HG13	2:F:56:LYS:HG2	1.73	0.71
1:D:3:GLN:HG2	1:D:28:ASP:HB3	1.78	0.64
1:D:148:SER:HB2	1:D:153:VAL:HG23	1.78	0.63
4:B:81:ARG:HG3	4:B:92:ILE:HG12	1.82	0.61
3:A:177:GLU:O	3:A:181:ARG:NH1	2.34	0.60
2:F:199:PHE:O	2:F:205:ASN:ND2	2.31	0.59
2:F:152:ASP:HB2	2:F:175:PRO:HG2	1.84	0.59
4:B:73:THR:HG22	4:B:75:LYS:H	1.67	0.58
3:A:230:LEU:HD11	3:A:243:LYS:HE3	1.86	0.57
2:F:4:THR:OG1	2:F:23:SER:OG	2.23	0.56
2:F:20:LEU:HD22	2:F:108:THR:HG21	1.87	0.56
3:A:5:MET:HB2	3:A:168:LEU:HD13	1.89	0.55
2:F:121:PRO:HD3	2:F:229:PRO:HB3	1.90	0.54
2:F:116:LEU:HD13	2:F:216:LEU:HD22	1.89	0.54
1:D:193:ILE:HD11	1:D:197:THR:HG21	1.89	0.54
3:A:69:ALA:HB1	5:C:6:ILE:HD11	1.90	0.53
1:D:69:LYS:NZ	3:A:163:GLU:OE2	2.35	0.52
3:A:230:LEU:HD13	3:A:245:ALA:HB2	1.90	0.52
2:F:119:VAL:HG21	2:F:216:LEU:HD21	1.92	0.52
2:F:201:GLN:HB3	7:F:301:ASP:HA	1.92	0.52
1:D:8:ILE:HG13	1:D:23:ASN:HB2	1.93	0.51
4:B:17:ASN:HA	4:B:72:PRO:HG2	1.94	0.50
1:D:111:PRO:HG3	1:D:160:VAL:HG11	1.93	0.49
2:F:241:ARG:HH21	2:F:242:ALA:HB2	1.77	0.49
2:F:126:VAL:HB	2:F:142:LEU:HD21	1.93	0.48
4:B:21:ASN:OD1	4:B:22:PHE:N	2.46	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:115:GLN:HG3	4:B:60:TRP:CH2	2.49	0.48
2:F:177:LYS:HD2	2:F:185:SER:HB3	1.95	0.48
2:F:134:ILE:HG23	2:F:197:ALA:HB1	1.96	0.48
3:A:13:SER:HB3	3:A:78:LEU:HD13	1.96	0.48
2:F:122:PRO:HB3	2:F:149:PHE:HB3	1.95	0.47
2:F:12:THR:HG21	2:F:80:LEU:HD13	1.97	0.47
3:A:207:GLY:HA2	3:A:240:THR:HB	1.96	0.47
2:F:209:CYS:O	2:F:235:SER:OG	2.28	0.47
3:A:109:LEU:HD22	3:A:161:GLU:HA	1.97	0.46
2:F:126:VAL:HG13	2:F:236:ALA:HB3	1.97	0.45
1:D:6:THR:OG1	1:D:25[A]:SER:OG	2.34	0.45
3:A:117:ALA:HB2	4:B:60:TRP:CE2	2.52	0.45
1:D:138:ASP:OD1	2:F:194:ARG:NH1	2.40	0.45
1:D:38:ARG:HB3	1:D:48:LEU:HD11	1.98	0.45
2:F:210:GLN:HG3	2:F:233:ILE:HG23	1.99	0.44
2:F:141:THR:HA	2:F:194:ARG:HA	1.98	0.44
4:B:37:VAL:HG22	4:B:82:VAL:HG22	2.01	0.43
2:F:31:VAL:HG21	2:F:73:SER:HB2	2.01	0.43
3:A:234:ARG:HD2	4:B:10:TYR:CE1	2.54	0.42
3:A:151:ARG:HB3	3:A:154:GLU:HG3	2.01	0.42
2:F:20:LEU:HB2	2:F:75:LEU:HB3	2.02	0.41
1:D:50:LEU:HD12	2:F:101:THR:HB	2.02	0.41
2:F:118:ASN:ND2	2:F:184:ASP:HB2	2.35	0.41
3:A:204:TRP:CE3	3:A:206:LEU:HD21	2.56	0.40
3:A:202:ARG:HD2	3:A:204:TRP:HE1	1.87	0.40
3:A:204:TRP:HE3	3:A:206:LEU:HD21	1.85	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	193/204 (95%)	188 (97%)	5 (3%)	0	100	100
2	F	239/242 (99%)	228 (95%)	10 (4%)	1 (0%)	30	36
3	A	252/278 (91%)	241 (96%)	11 (4%)	0	100	100
4	B	98/100 (98%)	90 (92%)	8 (8%)	0	100	100
5	C	7/9 (78%)	7 (100%)	0	0	100	100
All	All	789/833 (95%)	754 (96%)	34 (4%)	1 (0%)	48	59

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	F	95	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	174/182 (96%)	174 (100%)	0	100	100
2	F	209/212 (99%)	205 (98%)	4 (2%)	50	66
3	A	220/237 (93%)	216 (98%)	4 (2%)	51	68
4	B	95/95 (100%)	94 (99%)	1 (1%)	65	79
5	C	7/7 (100%)	7 (100%)	0	100	100
All	All	705/733 (96%)	696 (99%)	9 (1%)	61	76

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	F	6	THR
2	F	72	HIS
2	F	170	CYS
2	F	241	ARG
3	A	35	ARG
3	A	121	LYS
3	A	262	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	A	271	THR
4	B	89	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	39	GLN
1	D	95	ASN
2	F	36	GLN
2	F	136	HIS
2	F	166	HIS
3	A	54	GLN
3	A	87	GLN
3	A	224	GLN
3	A	262	GLN
4	B	13	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
9	EDO	A	301	-	3,3,3	0.47	0	2,2,2	0.23	0
6	NAG	D	301	1	14,14,15	0.38	0	17,19,21	0.45	0
7	ASP	F	301	-	7,8,8	1.15	1 (14%)	6,10,10	1.38	1 (16%)
8	SO4	A	302	-	4,4,4	0.24	0	6,6,6	0.07	0
8	SO4	F	304	-	4,4,4	0.24	0	6,6,6	0.08	0
6	NAG	F	302	2	14,14,15	0.29	0	17,19,21	0.40	0
6	NAG	D	304	1	14,14,15	0.34	0	17,19,21	0.53	0
6	NAG	D	303	1	14,14,15	0.40	0	17,19,21	0.49	0
8	SO4	F	303	-	4,4,4	0.24	0	6,6,6	0.07	0
6	NAG	D	302	1	14,14,15	0.21	0	17,19,21	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	EDO	A	301	-	-	0/1/1/1	-
6	NAG	D	301	1	-	0/6/23/26	0/1/1/1
7	ASP	F	301	-	-	2/8/8/8	-
6	NAG	F	302	2	-	2/6/23/26	0/1/1/1
6	NAG	D	304	1	-	2/6/23/26	0/1/1/1
6	NAG	D	303	1	-	0/6/23/26	0/1/1/1
6	NAG	D	302	1	-	1/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	F	301	ASP	OXT-C	-2.22	1.23	1.30

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	F	301	ASP	OXT-C-O	-2.69	117.98	124.08

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	F	301	ASP	C-CA-CB-CG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
6	F	302	NAG	C4-C5-C6-O6
6	F	302	NAG	O5-C5-C6-O6
6	D	304	NAG	C4-C5-C6-O6
6	D	304	NAG	O5-C5-C6-O6
7	F	301	ASP	N-CA-CB-CG
6	D	302	NAG	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	F	301	ASP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	D	195/204 (95%)	0.71	15 (7%)	19	20	34, 72, 119, 158	2 (1%)
2	F	241/242 (99%)	0.65	10 (4%)	41	43	46, 80, 124, 165	0
3	A	259/278 (93%)	0.84	28 (10%)	11	12	36, 78, 150, 179	1 (0%)
4	B	100/100 (100%)	0.92	8 (8%)	18	19	58, 102, 155, 176	0
5	C	9/9 (100%)	0.13	0	100	100	49, 50, 62, 64	0
All	All	804/833 (96%)	0.75	61 (7%)	20	20	34, 80, 143, 179	3 (0%)

All (61) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	A	276	PRO	4.4
1	D	136	PHE	4.0
2	F	142	LEU	3.9
3	A	201	LEU	3.8
3	A	215	LEU	3.7
3	A	200	THR	3.6
3	A	270	LEU	3.5
2	F	242	ALA	3.5
1	D	126	SER	3.3
1	D	133	VAL	3.3
4	B	15	ALA	3.1
1	D	123	LEU	3.1
3	A	248	VAL	3.0
3	A	272	LEU	3.0
3	A	113	TYR	3.0
3	A	205	ALA	2.9
1	D	152	ASP	2.8
3	A	260	HIS	2.8
4	B	1	ILE	2.7
2	F	182	LEU	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	A	179	LEU	2.7
3	A	271	THR	2.7
2	F	181	ALA	2.6
4	B	7	ILE	2.6
3	A	258	THR	2.6
3	A	261	VAL	2.5
1	D	193	ILE	2.5
3	A	266	LEU	2.5
3	A	274	TRP	2.5
3	A	192	HIS	2.5
3	A	40	ALA	2.4
3	A	225	THR	2.4
3	A	187	THR	2.4
3	A	41	ALA	2.3
1	D	132	SER	2.3
2	F	134	ILE	2.3
1	D	114	GLN	2.3
3	A	193	PRO	2.3
4	B	99	MET	2.3
1	D	155	ILE	2.3
2	F	143	VAL	2.3
2	F	132	ALA	2.2
4	B	71	THR	2.2
1	D	201	SER	2.2
4	B	78	TYR	2.2
3	A	189	VAL	2.2
3	A	236	ALA	2.2
4	B	9	VAL	2.2
1	D	125	ASP	2.1
2	F	87	LEU	2.1
3	A	259	CYS	2.1
1	D	187	ALA	2.1
1	D	147[A]	GLN	2.1
4	B	73	THR	2.1
3	A	199	ALA	2.1
2	F	165	VAL	2.1
2	F	184	ASP	2.0
1	D	192	ILE	2.0
1	D	153	VAL	2.0
3	A	247	VAL	2.0
3	A	206	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
7	ASP	F	301	9/9	0.18	0.16	162,167,168,170	0
6	NAG	D	304	14/15	0.40	0.14	116,121,127,128	0
6	NAG	D	302	14/15	0.55	0.13	89,104,110,111	0
8	SO4	A	302	5/5	0.55	0.11	135,141,149,174	0
6	NAG	D	301	14/15	0.68	0.14	85,98,108,110	0
8	SO4	F	304	5/5	0.73	0.08	108,113,130,131	0
6	NAG	D	303	14/15	0.74	0.12	84,91,95,99	0
8	SO4	F	303	5/5	0.75	0.12	106,108,129,142	0
6	NAG	F	302	14/15	0.77	0.12	72,77,81,82	0
9	EDO	A	301	4/4	0.83	0.16	61,64,66,68	0

6.5 Other polymers [i](#)

There are no such residues in this entry.