



Full wwPDB NMR Structure Validation Report ⓘ

Mar 8, 2026 – 12:54 PM UTC

PDB ID : 2N3A / pdb_00002n3a
BMRB ID : 25639
Title : Solution structure of LEDGF/p75 IBD in complex with POGZ peptide (1389-1404)
Authors : Tesina, P.; Cermakova, K.; Horejsi, M.; Prochazkova, K.; Fabry, M.; Sharma, S.; Christ, F.; Demeulemeester, J.; Debyser, Z.; De Rijck, J.; Veverka, V.; Rezacova, P.
Deposited on : 2015-05-26

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 86%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	16	100%
2	B	79	92%

2 Ensemble composition and analysis

This entry contains 29 models. Model 28 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	B:348-B:423 (76)	0.69	28

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 13, 14, 15, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28
2	16, 29
Single-model clusters	11

3 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 1564 atoms, of which 786 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Pogo transposable element with ZNF domain.

Mol	Chain	Residues	Atoms					Trace
1	A	16	Total	C	H	N	O	0
			226	78	97	16	35	

- Molecule 2 is a protein called PC4 and SFRS1-interacting protein.


Mol	Chain	Residues	Atoms						Trace
2	B	79	Total	C	H	N	O	S	0
			1338	404	689	118	120	7	

4 Residue-property plots


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Pogo transposable element with ZNF domain



- Molecule 2: PC4 and SFRS1-interacting protein




4.2 Scores per residue for each member of the ensemble


Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

- Molecule 1: Pogo transposable element with ZNF domain




- Molecule 2: PC4 and SFRS1-interacting protein




4.2.2 Score per residue for model 2

- Molecule 1: Pogo transposable element with ZNF domain




- Molecule 2: PC4 and SFRS1-interacting protein




4.2.3 Score per residue for model 3

- Molecule 1: Pogo transposable element with ZNF domain




- Molecule 2: PC4 and SFRS1-interacting protein




4.2.4 Score per residue for model 4

- Molecule 1: Pogo transposable element with ZNF domain




- Molecule 2: PC4 and SFRS1-interacting protein




4.2.5 Score per residue for model 5

- Molecule 1: Pogo transposable element with ZNF domain




- Molecule 2: PC4 and SFRS1-interacting protein




4.2.6 Score per residue for model 6

- Molecule 1: Pogo transposable element with ZNF domain




- Molecule 2: PC4 and SFRS1-interacting protein




4.2.7 Score per residue for model 7

- Molecule 1: Pogo transposable element with ZNF domain




- Molecule 2: PC4 and SFRS1-interacting protein




4.2.8 Score per residue for model 8

- Molecule 1: Pogo transposable element with ZNF domain




- Molecule 2: PC4 and SFRS1-interacting protein




4.2.9 Score per residue for model 9

- Molecule 1: Pogo transposable element with ZNF domain




- Molecule 2: PC4 and SFRS1-interacting protein




4.2.10 Score per residue for model 10

- Molecule 1: Pogo transposable element with ZNF domain




- Molecule 2: PC4 and SFRS1-interacting protein




4.2.11 Score per residue for model 11

- Molecule 1: Pogo transposable element with ZNF domain




- Molecule 2: PC4 and SFRS1-interacting protein




4.2.12 Score per residue for model 12

- Molecule 1: Pogo transposable element with ZNF domain




- Molecule 2: PC4 and SFRS1-interacting protein




4.2.13 Score per residue for model 13

- Molecule 1: Pogo transposable element with ZNF domain




- Molecule 2: PC4 and SFRS1-interacting protein




4.2.14 Score per residue for model 14

- Molecule 1: Pogo transposable element with ZNF domain

Chain A:  100%

E1389
G1390
E1391
S1392
E1393
T1394
E1395
S1396
F1397
Y1398
G1399
F1400
E1401
E1402
A1403
D1404


- Molecule 2: PC4 and SFRS1-interacting protein

Chain B:  86%  10% .

K348
L368
R372
L380
T394
I403
R404
V408
F423
K424
N425
M426


4.2.15 Score per residue for model 15

- Molecule 1: Pogo transposable element with ZNF domain

Chain A:  100%

E1389
G1390
E1391
S1392
E1393
T1394
E1395
S1396
F1397
Y1398
G1399
F1400
E1401
E1402
A1403
D1404


- Molecule 2: PC4 and SFRS1-interacting protein

Chain B:  87%  9% .

K348
D349
S350
I365
R372
D378
E379
L380
Q391
I403
K424
N425
M426


4.2.16 Score per residue for model 16

- Molecule 1: Pogo transposable element with ZNF domain

Chain A:  100%

E1389
G1390
E1391
S1392
E1393
T1394
E1395
S1396
F1397
Y1398
G1399
F1400
E1401
E1402
A1403
D1404


- Molecule 2: PC4 and SFRS1-interacting protein


Chain B:  91%  5% .

K348
E379
L380
V408
K424
N425
M426

4.2.17 Score per residue for model 17

- Molecule 1: Pogo transposable element with ZNF domain

Chain A:  100%



- Molecule 2: PC4 and SFRS1-interacting protein


Chain B:  90%  5% • •



4.2.18 Score per residue for model 18

- Molecule 1: Pogo transposable element with ZNF domain

Chain A:  100%



- Molecule 2: PC4 and SFRS1-interacting protein


Chain B:  91%  5% • •




4.2.19 Score per residue for model 19

- Molecule 1: Pogo transposable element with ZNF domain

Chain A:  100%




- Molecule 2: PC4 and SFRS1-interacting protein


Chain B:  86%  10% • •



4.2.20 Score per residue for model 20

- Molecule 1: Pogo transposable element with ZNF domain

Chain A:  100%



- Molecule 2: PC4 and SFRS1-interacting protein


Chain B:  94%



4.2.21 Score per residue for model 21

- Molecule 1: Pogo transposable element with ZNF domain

Chain A:  100%



- Molecule 2: PC4 and SFRS1-interacting protein


Chain B:  91%



4.2.22 Score per residue for model 22

- Molecule 1: Pogo transposable element with ZNF domain

Chain A:  100%




- Molecule 2: PC4 and SFRS1-interacting protein

Chain B:  89%




4.2.23 Score per residue for model 23

- Molecule 1: Pogo transposable element with ZNF domain




- Molecule 2: PC4 and SFRS1-interacting protein




4.2.24 Score per residue for model 24

- Molecule 1: Pogo transposable element with ZNF domain




- Molecule 2: PC4 and SFRS1-interacting protein




4.2.25 Score per residue for model 25

- Molecule 1: Pogo transposable element with ZNF domain





- Molecule 2: PC4 and SFRS1-interacting protein



4.2.26 Score per residue for model 26

- Molecule 1: Pogo transposable element with ZNF domain

Chain A:  100%




- Molecule 2: PC4 and SFRS1-interacting protein


Chain B:  95%



4.2.27 Score per residue for model 27

- Molecule 1: Pogo transposable element with ZNF domain

Chain A:  100%




- Molecule 2: PC4 and SFRS1-interacting protein


Chain B:  89%



4.2.28 Score per residue for model 28 (medoid)

- Molecule 1: Pogo transposable element with ZNF domain

Chain A:  100%




- Molecule 2: PC4 and SFRS1-interacting protein


Chain B:  90%




4.2.29 Score per residue for model 29

- Molecule 1: Pogo transposable element with ZNF domain

Chain A:  100%



- Molecule 2: PC4 and SFRS1-interacting protein

Chain B:  86%  10% .



5 Refinement protocol and experimental data overview

The models were refined using the following method: *torsion angle dynamics, molecular dynamics*.

Of the 100 calculated structures, 29 were deposited, based on the following criterion: *structures with the least restraint violations*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	
YASARA	refinement	
CYANA	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1036
Number of shifts mapped to atoms	1036
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	86%

6 Model quality [i](#)

6.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
2	B	0.76±0.04	0±0/628 (0.0± 0.0%)	0.82±0.02	0±0/839 (0.0± 0.0%)
All	All	0.76	0/18212 (0.0%)	0.82	3/24331 (0.0%)

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
2	B	365	ILE	N-CA-C	5.73	116.46	110.62	7	2
2	B	365	ILE	CB-CA-C	-5.25	105.05	112.14	7	1

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	0	0	0	0±0
2	B	623	660	658	0±0
All	All	18067	19140	19082	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
2:B:417:THR:HA	2:B:420:TYR:CD2	0.47	2.45	29	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	0	-	-	-	-	
2	B	75/79 (95%)	72±1 (97±1%)	2±1 (3±1%)	0±0 (0±1%)	44	80
All	All	2175/2755 (79%)	2102 (97%)	68 (3%)	5 (0%)	44	80

All 2 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	B	367	ASN	3
2	B	406	PHE	2

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	0	-	-	-	
2	B	72/75 (96%)	67±2 (94±2%)	5±2 (6±2%)	18	67
All	All	2088/2552 (82%)	1955 (94%)	133 (6%)	18	67

All 31 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	B	368	LEU	23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
2	B	380	LEU	20
2	B	403	ILE	12
2	B	365	ILE	9
2	B	408	VAL	9
2	B	348	MET	7
2	B	350	SER	5
2	B	414	GLU	4
2	B	405	ARG	4
2	B	396	MET	4
2	B	394	THR	4
2	B	404	ARG	3
2	B	379	GLU	3
2	B	378	ASP	3
2	B	364	LYS	2
2	B	395	GLU	2
2	B	372	ARG	2
2	B	423	PHE	2
2	B	391	GLN	2
2	B	387	MET	2
2	B	351	ARG	1
2	B	409	SER	1
2	B	389	GLN	1
2	B	418	MET	1
2	B	361	ASN	1
2	B	349	ASP	1
2	B	367	ASN	1
2	B	366	ASP	1
2	B	400	LEU	1
2	B	388	GLN	1
2	B	401	LYS	1

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 86% for the well-defined parts and 77% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1036
Number of shifts mapped to atoms	1036
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	79	-0.68 ± 0.14	Should be checked
$^{13}\text{C}_\beta$	79	0.36 ± 0.07	None needed (< 0.5 ppm)
$^{13}\text{C}'$	74	2.46 ± 0.13	Should be applied
^{15}N	75	0.07 ± 0.32	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 86%, i.e. 965 atoms were assigned a chemical shift out of a possible 1120. 0 out of 12 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	367/380 (97%)	147/152 (97%)	148/152 (97%)	72/76 (95%)
Sidechain	585/697 (84%)	398/451 (88%)	182/213 (85%)	5/33 (15%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	13/43 (30%)	11/22 (50%)	2/19 (11%)	0/2 (0%)
Overall	965/1120 (86%)	556/625 (89%)	332/384 (86%)	77/111 (69%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 77%, i.e. 1034 atoms were assigned a chemical shift out of a possible 1347. 0 out of 12 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	391/477 (82%)	163/192 (85%)	153/190 (81%)	75/95 (79%)
Sidechain	623/798 (78%)	427/513 (83%)	190/250 (76%)	6/35 (17%)
Aromatic	20/72 (28%)	18/36 (50%)	2/34 (6%)	0/2 (0%)
Overall	1034/1347 (77%)	608/741 (82%)	345/474 (73%)	81/132 (61%)


7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:



Random coil index (RCI) for chain B:

