



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 03:35 AM UTC

PDB ID : 6N3L / pdb_00006n3l
Title : Identification of novel, potent and selective GCN2 inhibitors as first-in-class anti-tumor agents
Authors : Hoffman, I.D.; Fujimoto, J.; Kurasawa, O.; Takagi, T.; Klein, M.G.; Kefala, G.; Ding, S.C.; Cary, D.R.; Mizojiri, R.
Deposited on : 2018-11-15
Resolution : 2.61 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.61 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4951 (2.64-2.60)
Clashscore	190562	5303 (2.64-2.60)
Ramachandran outliers	187476	5217 (2.64-2.60)
Sidechain outliers	187428	5217 (2.64-2.60)
RSRZ outliers	180081	4950 (2.64-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	330	2% 56% 13% . 30%
1	B	330	% 52% 15% .. 31%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3926 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called eIF-2-alpha kinase GCN2,eIF-2-alpha kinase GCN2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	231	Total	C	N	O	S	0	0	0
			1916	1240	331	338	7			
1	B	227	Total	C	N	O	S	0	1	0
			1894	1227	328	332	7			

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	561	MET	-	initiating methionine	UNP Q9P2K8
A	562	HIS	-	expression tag	UNP Q9P2K8
A	563	HIS	-	expression tag	UNP Q9P2K8
A	564	HIS	-	expression tag	UNP Q9P2K8
A	565	HIS	-	expression tag	UNP Q9P2K8
A	566	HIS	-	expression tag	UNP Q9P2K8
A	567	HIS	-	expression tag	UNP Q9P2K8
A	568	GLU	-	expression tag	UNP Q9P2K8
A	569	ASN	-	expression tag	UNP Q9P2K8
A	570	LEU	-	expression tag	UNP Q9P2K8
A	571	TYR	-	expression tag	UNP Q9P2K8
A	572	PHE	-	expression tag	UNP Q9P2K8
A	573	GLN	-	expression tag	UNP Q9P2K8
A	574	GLY	-	expression tag	UNP Q9P2K8
A	575	GLY	-	expression tag	UNP Q9P2K8
A	576	SER	-	expression tag	UNP Q9P2K8
A	658	ASN	-	linker	UNP Q9P2K8
A	807	ALA	LYS	engineered mutation	UNP Q9P2K8
A	848	ASN	ASP	engineered mutation	UNP Q9P2K8
A	899	ALA	THR	engineered mutation	UNP Q9P2K8
A	904	ALA	THR	engineered mutation	UNP Q9P2K8
B	561	MET	-	initiating methionine	UNP Q9P2K8
B	562	HIS	-	expression tag	UNP Q9P2K8
B	563	HIS	-	expression tag	UNP Q9P2K8
B	564	HIS	-	expression tag	UNP Q9P2K8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	565	HIS	-	expression tag	UNP Q9P2K8
B	566	HIS	-	expression tag	UNP Q9P2K8
B	567	HIS	-	expression tag	UNP Q9P2K8
B	568	GLU	-	expression tag	UNP Q9P2K8
B	569	ASN	-	expression tag	UNP Q9P2K8
B	570	LEU	-	expression tag	UNP Q9P2K8
B	571	TYR	-	expression tag	UNP Q9P2K8
B	572	PHE	-	expression tag	UNP Q9P2K8
B	573	GLN	-	expression tag	UNP Q9P2K8
B	574	GLY	-	expression tag	UNP Q9P2K8
B	575	GLY	-	expression tag	UNP Q9P2K8
B	576	SER	-	expression tag	UNP Q9P2K8
B	781	ASN	-	linker	UNP Q9P2K8
B	807	ALA	LYS	engineered mutation	UNP Q9P2K8
B	848	ASN	ASP	engineered mutation	UNP Q9P2K8
B	899	ALA	THR	engineered mutation	UNP Q9P2K8
B	904	ALA	THR	engineered mutation	UNP Q9P2K8

- Molecule 2 is N-{6-[(1-methyl-2-[[4-(trifluoromethyl)phenyl]amino]-1H-benzimidazol-5-yl)oxy]pyrimidin-4-yl}cyclopropanecarboxamide (CCD ID: KAV) (formula: C₂₃H₁₉F₃N₆O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	0	0
			34	23	3	6	2		
2	B	1	Total	C	F	N	O	0	0
			34	23	3	6	2		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	27	Total 27	O 27	0	0
3	B	21	Total 21	O 21	0	0

- Molecule 1: eIF-2-alpha kinase GCN2, eIF-2-alpha kinase GCN2



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	83.89Å 122.23Å 120.67Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 2.61 25.00 – 2.61	Depositor EDS
% Data completeness (in resolution range)	97.6 (25.00-2.61) 97.6 (25.00-2.61)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.06 (at 2.60Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.223 , 0.290 0.225 , 0.290	Depositor DCC
R_{free} test set	960 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	53.5	Xtriage
Anisotropy	0.312	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 30.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3926	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: KAV

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.04	0/1960	1.44	5/2644 (0.2%)
1	B	1.04	0/1939	1.42	5/2611 (0.2%)
All	All	1.04	0/3899	1.43	10/5255 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
All	All	0	4

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	921	ASN	CA-CB-CG	7.33	119.93	112.60
1	B	592	GLU	CB-CA-C	6.21	119.78	109.53
1	B	867	PHE	CA-CB-CG	5.80	119.60	113.80
1	A	796	HIS	CA-C-O	-5.79	114.41	120.55
1	A	867	PHE	CA-CB-CG	5.66	119.46	113.80
1	A	824	ARG	CA-C-N	5.60	127.78	120.28
1	A	824	ARG	C-N-CA	5.60	127.78	120.28
1	B	824	ARG	CA-C-N	5.52	127.68	120.28
1	B	824	ARG	C-N-CA	5.52	127.68	120.28
1	B	591	GLU	CB-CG-CD	5.11	121.29	112.60

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	920	TYR	Peptide
1	A	956	ASP	Peptide
1	B	920	TYR	Peptide
1	B	956	ASP	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1916	0	1907	16	0
1	B	1894	0	1891	30	0
2	A	34	0	0	0	0
2	B	34	0	0	1	0
3	A	27	0	0	0	0
3	B	21	0	0	0	0
All	All	3926	0	3798	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:639:LEU:HD21	1:B:844:MET:HE3	1.53	0.89
1:A:639:LEU:HD21	1:A:844:MET:HE3	1.55	0.88
1:B:650[B]:ARG:HG2	1:B:650[B]:ARG:HH11	1.45	0.80
1:B:650[B]:ARG:HG2	1:B:650[B]:ARG:NH1	2.14	0.62
1:A:584:SER:OG	1:A:585:ARG:N	2.22	0.61
1:B:962:PHE:CE2	1:B:975:LYS:HG3	2.37	0.59
1:B:935:GLU:OE2	1:B:947:ARG:NH2	2.35	0.58
1:B:956:ASP:CG	1:B:957:PRO:HD2	2.30	0.57
1:B:980:TRP:CD1	1:B:990:PRO:HD3	2.41	0.55
1:A:618:VAL:HA	1:A:800:ILE:O	2.08	0.54
1:B:910:PRO:O	1:B:911:GLU:C	2.51	0.54
1:B:909:SER:O	1:B:910:PRO:O	2.27	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:980:TRP:CD1	1:A:990:PRO:HD3	2.44	0.53
1:A:956:ASP:CG	1:A:957:PRO:HD2	2.34	0.52
1:A:623:ILE:HD11	1:A:630:PHE:HD1	1.75	0.52
1:B:623:ILE:HD11	1:B:630:PHE:HD1	1.76	0.51
1:B:618:VAL:HA	1:B:800:ILE:O	2.11	0.50
1:B:911:GLU:OE2	1:B:989:ARG:NH1	2.44	0.49
1:A:825:LEU:HD11	1:A:937:SER:HA	1.94	0.49
1:A:592:GLU:HG2	1:A:605:ILE:HG21	1.96	0.48
1:B:830:ARG:HH21	1:B:1001:LEU:HB2	1.79	0.48
1:A:624:ASN:O	1:A:630:PHE:HB2	2.16	0.46
1:B:949:PHE:CD1	1:B:949:PHE:C	2.94	0.46
1:A:991:THR:OG1	1:A:994:GLU:HB2	2.16	0.45
1:A:584:SER:HG	1:A:585:ARG:H	1.58	0.45
1:B:991:THR:OG1	1:B:994:GLU:HB2	2.17	0.45
1:B:640:LEU:HD21	2:B:2501:KAV:C6	2.47	0.45
1:B:649:VAL:HG12	1:B:802:MET:HE3	1.99	0.44
1:A:650:ARG:HB2	1:A:803:GLU:HB3	2.00	0.44
1:B:846:HIS:O	1:B:847:ARG:HB2	2.17	0.43
1:B:911:GLU:OE2	1:B:989:ARG:NH2	2.50	0.43
1:B:593:LEU:HD11	1:B:608:GLN:HB2	2.01	0.43
1:B:631:ARG:HD3	1:B:631:ARG:HA	1.78	0.43
1:B:962:PHE:CZ	1:B:975:LYS:HG3	2.54	0.43
1:A:620:ARG:HD3	1:A:797:TYR:CE1	2.54	0.43
1:B:825:LEU:HD11	1:B:937:SER:HA	2.00	0.42
1:B:629:GLN:CD	1:B:629:GLN:H	2.27	0.42
1:B:954:LEU:HB3	1:B:982:LEU:HD13	2.01	0.41
1:B:656:ILE:HA	1:B:797:TYR:O	2.20	0.41
1:B:590:PHE:HD2	1:B:607:VAL:HG13	1.85	0.41
1:A:656:ILE:HA	1:A:797:TYR:O	2.21	0.41
1:A:846:HIS:O	1:A:847:ARG:HB2	2.21	0.41
1:B:623:ILE:HD11	1:B:630:PHE:CD1	2.55	0.41
1:A:995:LEU:O	1:A:998:SER:HB3	2.21	0.40
1:B:965:ASP:N	1:B:965:ASP:OD1	2.55	0.40
1:B:591:GLU:OE2	1:B:591:GLU:HA	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	219/330 (66%)	208 (95%)	9 (4%)	2 (1%)	14	28
1	B	214/330 (65%)	206 (96%)	6 (3%)	2 (1%)	14	28
All	All	433/660 (66%)	414 (96%)	15 (4%)	4 (1%)	14	28

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	910	PRO
1	B	956	ASP
1	A	625	PRO
1	A	956	ASP

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	208/291 (72%)	191 (92%)	17 (8%)	10	22
1	B	205/291 (70%)	187 (91%)	18 (9%)	9	19
All	All	413/582 (71%)	378 (92%)	35 (8%)	10	21

All (35) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	584	SER
1	A	591	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	625	PRO
1	A	629	GLN
1	A	646	GLU
1	A	653	ASN
1	A	808	SER
1	A	822	THR
1	A	909	SER
1	A	921	ASN
1	A	926	LEU
1	A	956	ASP
1	A	966	PHE
1	A	973	LYS
1	A	974	GLN
1	A	975	LYS
1	A	983	ASN
1	B	584	SER
1	B	591	GLU
1	B	598	LYS
1	B	607	VAL
1	B	629	GLN
1	B	633	ILE
1	B	637	VAL
1	B	646	GLU
1	B	653	ASN
1	B	847	ARG
1	B	850	LYS
1	B	909	SER
1	B	911	GLU
1	B	926	LEU
1	B	956	ASP
1	B	965	ASP
1	B	966	PHE
1	B	983	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	644	HIS
1	A	647	ASN
1	A	939	HIS
1	A	953	GLN
1	A	974	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	644	HIS
1	B	647	ASN
1	B	816	GLN
1	B	953	GLN
1	B	983	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	KAV	A	2501	-	37,38,38	1.31	3 (8%)	52,56,56	2.36	16 (30%)
2	KAV	B	2501	-	37,38,38	1.15	2 (5%)	52,56,56	2.07	11 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	KAV	A	2501	-	-	2/22/24/24	0/5/5/5
2	KAV	B	2501	-	-	0/22/24/24	0/5/5/5

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	2501	KAV	C3-N4	3.80	1.41	1.36
2	A	2501	KAV	C16-N15	-3.10	1.33	1.39
2	B	2501	KAV	C3-N4	3.07	1.40	1.36
2	A	2501	KAV	O19-C20	-2.58	1.33	1.36
2	B	2501	KAV	O19-C20	-2.01	1.34	1.36

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	2501	KAV	C30-N31-C20	9.58	121.80	114.50
2	A	2501	KAV	N31-C30-N29	-6.39	118.91	128.58
2	B	2501	KAV	C30-N31-C20	6.10	119.14	114.50
2	B	2501	KAV	N31-C30-N29	-5.39	120.43	128.58
2	B	2501	KAV	F13-C11-C8	-4.82	102.57	112.90
2	A	2501	KAV	F13-C11-C8	-4.39	103.50	112.90
2	B	2501	KAV	F14-C11-C8	4.30	122.11	112.90
2	B	2501	KAV	C21-C22-N29	-4.23	117.38	122.92
2	A	2501	KAV	C16-N15-C3	4.17	110.56	104.06
2	A	2501	KAV	C6-C7-C8	-4.00	115.65	121.17
2	A	2501	KAV	C9-C8-C7	3.74	123.60	118.03
2	A	2501	KAV	C7-C6-C5	3.27	124.06	120.30
2	B	2501	KAV	C16-N15-C3	3.01	108.74	104.06
2	B	2501	KAV	C9-C8-C7	2.94	122.41	118.03
2	A	2501	KAV	C26-C24-N23	2.84	120.64	115.16
2	A	2501	KAV	F14-C11-F13	2.82	115.95	105.77
2	B	2501	KAV	C26-C24-N23	2.79	120.55	115.16
2	A	2501	KAV	C18-O19-C20	2.69	126.26	119.06
2	B	2501	KAV	C28-C26-C24	2.67	119.85	117.24
2	A	2501	KAV	O25-C24-C26	-2.54	118.27	122.19
2	B	2501	KAV	C30-N29-C22	2.50	123.43	115.25
2	B	2501	KAV	C6-C7-C8	-2.44	117.79	121.17
2	A	2501	KAV	C21-C22-N29	-2.40	119.77	122.92
2	A	2501	KAV	C27-C26-C24	-2.27	115.02	117.24
2	A	2501	KAV	C34-C16-N15	-2.23	107.35	110.19
2	A	2501	KAV	C21-C20-N31	-2.12	121.47	124.47
2	A	2501	KAV	C30-N29-C22	2.12	122.17	115.25

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	2501	KAV	O25-C24-C26-C27
2	A	2501	KAV	N23-C24-C26-C27


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	2501	KAV	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	231/330 (70%)	-0.02	6 (2%) 57 52	36, 62, 94, 111	0
1	B	227/330 (68%)	-0.05	4 (1%) 67 63	27, 59, 97, 163	1 (0%)
All	All	458/660 (69%)	-0.03	10 (2%) 62 57	27, 60, 96, 163	1 (0%)

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1002	PRO	2.9
1	A	1002	PRO	2.9
1	A	957	PRO	2.6
1	A	904	ALA	2.5
1	A	920	TYR	2.3
1	B	966	PHE	2.3
1	A	962	PHE	2.2
1	B	920	TYR	2.1
1	A	626	ALA	2.0
1	B	626	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)



In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	KAV	A	2501	34/34	0.95	0.07	39,46,51,53	0
2	KAV	B	2501	34/34	0.95	0.07	36,45,50,54	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around KAV A 2501:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.