



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 05:15 PM UTC

PDB ID : 7N3H / pdb_00007n3h
Title : Crystal structure of an anti-SARS-CoV-2 human neutralizing antibody Fab fragment C099
Authors : Flyak, A.I.; Bjorkman, P.J.; Barnes, C.O.
Deposited on : 2021-06-01
Resolution : 1.26 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.26 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1693 (1.28-1.24)
Clashscore	190562	1730 (1.28-1.24)
Ramachandran outliers	187476	1695 (1.28-1.24)
Sidechain outliers	187428	1694 (1.28-1.24)
RSRZ outliers	180081	1693 (1.28-1.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	230	6% 83% 10% 7%
2	L	215	5% 91% 8%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7139 atoms, of which 3241 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called C099 Fab Heavy Chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	H	214	Total	C	H	N	O	S	0	11	0
			3225	1018	1610	266	323	8			

- Molecule 2 is a protein called C099 Fab Light Chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	L	213	Total	C	H	N	O	S	0	8	0
			3282	1037	1631	277	333	4			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	H	310	Total	O	0	0
			310	310		
3	L	322	Total	O	0	0
			322	322		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: C099 Fab Heavy Chain



- Molecule 2: C099 Fab Light Chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	44.10Å 92.58Å 115.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.95 – 1.26 42.95 – 1.26	Depositor EDS
% Data completeness (in resolution range)	90.5 (42.95-1.26) 90.4 (42.95-1.26)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.23 (at 1.26Å)	Xtriage
Refinement program	PHENIX 1.18.2_3874	Depositor
R, R_{free}	0.180 , 0.202 0.181 , 0.203	Depositor DCC
R_{free} test set	5834 reflections (4.56%)	wwPDB-VP
Wilson B-factor (Å ²)	13.2	Xtriage
Anisotropy	0.039	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 36.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7139	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	H	0.43	0/1682	0.63	0/2287
2	L	0.41	0/1710	0.59	0/2320
All	All	0.42	0/3392	0.61	0/4607

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1615	1610	1612	25	0
2	L	1651	1631	1635	18	1
3	H	310	0	0	11	2
3	L	322	0	0	13	2
All	All	3898	3241	3247	43	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:101:GLN:NE2	3:L:302:HOH:O	2.07	0.86

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:156:GLN:OE1	3:L:301:HOH:O	1.98	0.81
1:H:204:ASN:OD1	3:H:301:HOH:O	1.98	0.80
3:H:533:HOH:O	2:L:124:GLU:HG2	1.81	0.80
1:H:187:SER:CB	3:H:302:HOH:O	2.31	0.79
1:H:3:GLN:NE2	3:H:304:HOH:O	2.20	0.74
2:L:170:LYS:HG3	3:L:557:HOH:O	1.86	0.74
2:L:124:GLU:OE1	3:L:303:HOH:O	2.07	0.71
2:L:170:LYS:CG	3:L:557:HOH:O	2.41	0.69
1:H:13:GLN:NE2	3:H:305:HOH:O	2.25	0.68
1:H:57[B]:THR:HG21	1:H:69:ILE:HG22	1.75	0.67
2:L:82:GLU:OE1	3:L:304:HOH:O	2.13	0.64
1:H:126:PRO:O	3:H:303:HOH:O	2.15	0.64
1:H:57[B]:THR:HG21	1:H:59:TYR:CZ	2.37	0.59
2:L:144:GLU:HG3	3:L:307:HOH:O	2.01	0.59
1:H:57[B]:THR:HG21	1:H:59:TYR:OH	2.05	0.57
2:L:18:ARG:NE	3:L:311:HOH:O	2.37	0.57
1:H:57[B]:THR:CG2	1:H:69:ILE:HG22	2.35	0.56
2:L:18:ARG:NH2	3:L:311:HOH:O	2.39	0.55
2:L:150:LYS:NZ	3:L:310:HOH:O	2.33	0.54
2:L:18:ARG:CZ	3:L:311:HOH:O	2.57	0.53
1:H:51:MET:HE1	1:H:70:SER:HA	1.93	0.51
1:H:12:ILE:HG21	1:H:82(C):LEU:HD13	1.94	0.50
2:L:124:GLU:CD	3:L:303:HOH:O	2.54	0.50
1:H:178:LEU:C	1:H:178:LEU:HD12	2.37	0.50
1:H:214:LYS:NZ	3:H:307:HOH:O	2.45	0.50
1:H:126:PRO:HD3	1:H:138:LEU:HD23	1.95	0.49
1:H:29:VAL:HG13	1:H:34:MET:HG3	1.94	0.48
1:H:57[B]:THR:CG2	1:H:69:ILE:CG2	2.92	0.48
1:H:127:SER:O	1:H:128:SER:C	2.58	0.46
1:H:51:MET:HE3	1:H:69:ILE:HG23	1.98	0.46
1:H:57[B]:THR:CG2	1:H:59:TYR:CZ	2.99	0.46
1:H:2:VAL:N	3:H:309:HOH:O	2.48	0.46
1:H:57[B]:THR:HG23	1:H:69:ILE:CG2	2.47	0.45
1:H:94:ARG:C	1:H:94:ARG:HD2	2.44	0.43
1:H:213:PRO:HA	3:H:303:HOH:O	2.17	0.43
1:H:57[B]:THR:HG22	1:H:59:TYR:CE2	2.54	0.43
1:H:187:SER:HB3	3:H:302:HOH:O	2.09	0.42
2:L:122:SER:HB2	2:L:124:GLU:OE1	2.20	0.41
3:H:533:HOH:O	2:L:124:GLU:CG	2.55	0.41
2:L:123:ASP:HB2	3:L:303:HOH:O	2.20	0.41
2:L:119:PHE:HA	2:L:120:PRO:HD3	1.96	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:164:VAL:HG22	2:L:176:LEU:HD12	2.04	0.40

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:412:HOH:O	3:L:554:HOH:O[4_455]	1.81	0.39
3:H:504:HOH:O	3:H:570:HOH:O[3_555]	1.98	0.22
3:H:485:HOH:O	3:L:337:HOH:O[3_555]	2.00	0.20
2:L:5:THR:OG1	2:L:153[B]:ASN:OD1[4_455]	2.08	0.12

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	221/230 (96%)	216 (98%)	4 (2%)	1 (0%)	24	6
2	L	219/215 (102%)	215 (98%)	4 (2%)	0	100	100
All	All	440/445 (99%)	431 (98%)	8 (2%)	1 (0%)	43	16

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	100	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	188/193 (97%)	188 (100%)	0	100	100
2	L	189/183 (103%)	187 (99%)	2 (1%)	65	30
All	All	377/376 (100%)	375 (100%)	2 (0%)	78	56

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	L	34	LEU
2	L	204	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	H	39	GLN
1	H	171	GLN
2	L	39	GLN
2	L	161	GLN
2	L	200	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	H	214/230 (93%)	0.24	14 (6%)	25 26	7, 17, 39, 56	11 (5%)
2	L	213/215 (99%)	0.29	11 (5%)	33 34	10, 18, 36, 56	8 (3%)
All	All	427/445 (95%)	0.26	25 (5%)	28 29	7, 17, 37, 56	19 (4%)

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	L	213	GLY	5.6
1	H	41	PRO	4.8
1	H	2	VAL	4.4
1	H	214	LYS	3.6
1	H	42	GLY	3.5
1	H	213	PRO	3.2
1	H	127	SER	3.2
1	H	138	LEU	3.2
1	H	133	GLY	3.2
2	L	203	SER	3.2
2	L	126	LEU	3.1
2	L	128	SER	3.1
2	L	11[A]	LEU	3.0
1	H	128	SER	2.8
1	H	43	LYS	2.7
2	L	1	GLU	2.6
1	H	40	ALA	2.6
2	L	124	GLU	2.4
2	L	204	SER	2.3
2	L	189	LYS	2.2
1	H	212	GLU	2.2
1	H	125	ALA	2.1
2	L	108	LYS	2.1
1	H	13	GLN	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	L	127	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.