



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 10:48 PM UTC

PDB ID : 9N3N / pdb_00009n3n
Title : Crystal structure of PRMT5:MEP50 in complex with MTA and oxamide compound 3
Authors : Whittington, D.A.
Deposited on : 2025-01-31
Resolution : 2.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1009 (2.76-2.76)
Clashscore	190562	1044 (2.76-2.76)
Ramachandran outliers	187476	1024 (2.76-2.76)
Sidechain outliers	187428	1024 (2.76-2.76)
RSRZ outliers	180081	1009 (2.76-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	645	87% 10% ..
2	B	350	3% 77% 9% 14%

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 7636 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein arginine N-methyltransferase 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	625	Total	C	N	O	S	0	1	0
			5071	3243	873	931	24			

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	initiating methionine	UNP O14744
A	-6	ASP	-	expression tag	UNP O14744
A	-5	TYR	-	expression tag	UNP O14744
A	-4	LYS	-	expression tag	UNP O14744
A	-3	ASP	-	expression tag	UNP O14744
A	-2	ASP	-	expression tag	UNP O14744
A	-1	ASP	-	expression tag	UNP O14744
A	0	ASP	-	expression tag	UNP O14744
A	1	LYS	-	expression tag	UNP O14744

- Molecule 2 is a protein called Methylosome protein 50.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	302	Total	C	N	O	S	0	0	0
			2286	1434	391	448	13			

There are 9 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	-7	MET	-	initiating methionine	UNP Q9BQA1
B	-6	HIS	-	expression tag	UNP Q9BQA1
B	-5	HIS	-	expression tag	UNP Q9BQA1
B	-4	HIS	-	expression tag	UNP Q9BQA1
B	-3	HIS	-	expression tag	UNP Q9BQA1
B	-2	HIS	-	expression tag	UNP Q9BQA1
B	-1	HIS	-	expression tag	UNP Q9BQA1

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	0	HIS	-	expression tag	UNP Q9BQA1
B	1	HIS	-	expression tag	UNP Q9BQA1

- Molecule 3 is 5'-DEOXY-5'-METHYLTHIOADENOSINE (CCD ID: MTA) (formula: $C_{11}H_{15}N_5O_3S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			20	11	5	3	1		

- Molecule 4 is N-(6-amino-5-cyclopropylpyridin-3-yl)-2-[(2R,5S)-2-(1,3-benzothiazol-5-yl)-5-methylpiperidin-1-yl]-2-oxoacetamide (CCD ID: A1BV5) (formula: $C_{23}H_{25}N_5O_2S$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			31	23	5	2	1		

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	4	Total Cl 4 4	0	0

- Molecule 6 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $\text{C}_2\text{H}_6\text{O}_2$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0

- Molecule 7 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			7	4	3		
7	A	1	Total	C	O	0	0
			7	4	3		
7	B	1	Total	C	O	0	0
			7	4	3		

- Molecule 8 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: $C_6H_{14}O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			10	6	4		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	109	Total 109	O 109	0	0
9	B	28	Total 28	O 28	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Protein arginine N-methyltransferase 5



- Molecule 2: Methylosome protein 50



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	102.49Å 136.41Å 178.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.32 – 2.75 46.32 – 2.75	Depositor EDS
% Data completeness (in resolution range)	100.0 (46.32-2.75) 99.9 (46.32-2.75)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.69 (at 2.77Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487, REFMAC 5.8.0267	Depositor
R, R_{free}	0.218 , 0.264 (Not available) , 0.244	Depositor DCC
R_{free} test set	1577 reflections (4.81%)	wwPDB-VP
Wilson B-factor (Å ²)	56.6	Xtriage
Anisotropy	0.146	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 35.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7636	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, CL, A1BV5, MTA, PGE, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.11	0/5213	0.33	0/7092
2	B	0.11	0/2340	0.37	0/3195
All	All	0.11	0/7553	0.34	0/10287

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5071	0	4964	38	0
2	B	2286	0	2203	13	0
3	A	20	0	15	0	0
4	A	31	0	0	0	0
5	A	4	0	0	0	0
6	A	48	0	72	6	0
6	B	8	0	12	1	0
7	A	14	0	20	1	0
7	B	7	0	10	0	0
8	A	10	0	14	1	0
9	A	109	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
9	B	28	0	0	1	0
All	All	7636	0	7310	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:360:VAL:HG13	1:A:430:ASP:H	1.51	0.74
2:B:223:GLN:HE22	2:B:269:PRO:HD3	1.56	0.71
2:B:35:ARG:HA	2:B:304:THR:HG21	1.74	0.69
2:B:35:ARG:NH1	2:B:86:LEU:O	2.32	0.62
1:A:318:ASN:OD1	1:A:393:LYS:NZ	2.33	0.61
2:B:223:GLN:NE2	2:B:267:SER:O	2.34	0.60
2:B:210:ALA:HB3	2:B:211:PRO:HD3	1.82	0.59
2:B:27:MET:HE1	2:B:68:PRO:HB2	1.87	0.56
1:A:360:VAL:HG13	1:A:430:ASP:N	2.21	0.55
1:A:465:PRO:HA	1:A:559:LEU:HA	1.87	0.55
1:A:140:ASN:O	1:A:144:THR:HG23	2.08	0.54
1:A:405:GLN:HA	1:A:409:TRP:HB2	1.91	0.52
1:A:592:VAL:HG21	1:A:598:ILE:HD11	1.90	0.52
1:A:200:LYS:H	8:A:721:PGE:H62	1.75	0.52
1:A:545:ASN:OD1	1:A:594:GLU:N	2.41	0.51
1:A:47:HIS:HB3	1:A:50:PHE:HB2	1.93	0.51
2:B:102:ALA:HB2	2:B:122:TYR:CD1	2.46	0.51
1:A:510:HIS:HA	6:A:711:EDO:H22	1.94	0.51
1:A:361:LEU:HD21	1:A:375:SER:HB3	1.92	0.50
1:A:122:PHE:HB2	6:A:715:EDO:H12	1.94	0.50
1:A:493:ASP:HB3	1:A:496:ALA:HB2	1.93	0.50
2:B:264:SER:OG	2:B:270:PHE:N	2.40	0.49
1:A:554:TYR:HB3	1:A:567:ILE:HG13	1.93	0.49
1:A:315:LEU:HD11	1:A:421:ARG:NH1	2.28	0.49
1:A:51:LYS:O	1:A:62:ARG:NH2	2.43	0.49
1:A:599:CYS:HB3	1:A:619:ALA:HB3	1.95	0.48
1:A:170:ASN:ND2	2:B:205:GLN:O	2.46	0.48
1:A:185:THR:O	1:A:188:TRP:HB2	2.15	0.46
1:A:70:ASP:OD1	1:A:70:ASP:N	2.51	0.44
1:A:114:GLY:C	6:A:715:EDO:H22	2.42	0.44
1:A:220:ARG:HA	6:A:716:EDO:H12	1.99	0.44
1:A:273:SER:OG	6:B:401:EDO:H22	2.17	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:362:MET:HG2	1:A:389:TYR:HB2	1.99	0.44
1:A:68:ARG:HG2	2:B:51:GLY:O	2.18	0.44
2:B:145:LYS:NZ	9:B:502:HOH:O	2.49	0.44
2:B:134:LEU:HD23	2:B:175:ALA:HB1	2.00	0.43
1:A:51:LYS:HG3	1:A:89:TRP:CE3	2.54	0.43
1:A:228:ALA:HA	1:A:263:GLN:O	2.18	0.43
1:A:250:HIS:O	1:A:254:ILE:HG12	2.18	0.43
1:A:419:ASP:OD2	1:A:421:ARG:NH1	2.52	0.43
1:A:546:THR:OG1	6:A:716:EDO:O2	2.30	0.42
2:B:171:THR:HG21	2:B:216:THR:HA	2.02	0.41
1:A:82:ILE:O	1:A:120:PRO:HD2	2.21	0.41
1:A:13:ARG:HB2	1:A:285:GLU:OE1	2.21	0.41
1:A:73:LEU:HB2	1:A:78:TRP:CE2	2.56	0.41
1:A:143:HIS:HB3	6:A:714:EDO:H21	2.03	0.40
1:A:479:LYS:HD3	7:A:719:PEG:H11	2.03	0.40
1:A:556:GLU:OE2	1:A:568:ARG:NH1	2.45	0.40
1:A:419:ASP:OD1	1:A:421:ARG:HD3	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	624/645 (97%)	598 (96%)	25 (4%)	1 (0%)	43	64
2	B	300/350 (86%)	281 (94%)	16 (5%)	3 (1%)	12	24
All	All	924/995 (93%)	879 (95%)	41 (4%)	4 (0%)	30	50

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	147	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	210	ALA
2	B	224	GLN
1	A	441	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	441/570 (77%)	430 (98%)	11 (2%)	42	64
2	B	239/298 (80%)	230 (96%)	9 (4%)	29	52
All	All	680/868 (78%)	660 (97%)	20 (3%)	37	61

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	169	GLU
1	A	215	ASN
1	A	234	SER
1	A	273	SER
1	A	277	PHE
1	A	285	GLU
1	A	358	VAL
1	A	360	VAL
1	A	467	GLU
1	A	479	LYS
1	A	629	THR
2	B	44	LEU
2	B	80	GLU
2	B	141	VAL
2	B	206	ILE
2	B	227	VAL
2	B	240	VAL
2	B	247	CYS
2	B	274	LEU
2	B	294	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	147	HIS
1	A	289	GLN
2	B	156	GLN
2	B	169	GLN
2	B	223	GLN
2	B	294	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 24 ligands modelled in this entry, 4 are monoatomic - leaving 20 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	EDO	A	707	-	3,3,3	0.43	0	2,2,2	0.38	0
6	EDO	A	716	-	3,3,3	0.35	0	2,2,2	0.42	0
6	EDO	B	401	-	3,3,3	0.40	0	2,2,2	0.34	0
6	EDO	A	712	-	3,3,3	0.42	0	2,2,2	0.38	0
6	EDO	A	709	-	3,3,3	0.43	0	2,2,2	0.38	0
6	EDO	A	715	-	3,3,3	0.41	0	2,2,2	0.32	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	EDO	B	402	-	3,3,3	0.41	0	2,2,2	0.36	0
8	PGE	A	721	-	9,9,9	0.32	0	8,8,8	0.29	0
4	A1BV5	A	702	-	35,35,35	1.11	3 (8%)	41,51,51	3.15	10 (24%)
6	EDO	A	710	-	3,3,3	0.43	0	2,2,2	0.35	0
6	EDO	A	708	-	3,3,3	0.43	0	2,2,2	0.34	0
6	EDO	A	718	-	3,3,3	0.43	0	2,2,2	0.38	0
6	EDO	A	713	-	3,3,3	0.41	0	2,2,2	0.38	0
7	PEG	B	403	-	6,6,6	0.09	0	5,5,5	0.12	0
7	PEG	A	719	-	6,6,6	0.11	0	5,5,5	0.09	0
6	EDO	A	711	-	3,3,3	0.43	0	2,2,2	0.37	0
6	EDO	A	714	-	3,3,3	0.43	0	2,2,2	0.31	0
6	EDO	A	717	-	3,3,3	0.41	0	2,2,2	0.38	0
7	PEG	A	720	-	6,6,6	0.11	0	5,5,5	0.10	0
3	MTA	A	701	-	22,22,22	0.54	0	32,32,32	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	A	707	-	-	1/1/1/1	-
6	EDO	A	716	-	-	0/1/1/1	-
6	EDO	B	401	-	-	1/1/1/1	-
6	EDO	A	712	-	-	1/1/1/1	-
6	EDO	A	709	-	-	1/1/1/1	-
6	EDO	A	715	-	-	1/1/1/1	-
6	EDO	B	402	-	-	0/1/1/1	-
8	PGE	A	721	-	-	3/7/7/7	-
4	A1BV5	A	702	-	-	2/20/35/35	0/5/5/5
6	EDO	A	710	-	-	0/1/1/1	-
6	EDO	A	708	-	-	0/1/1/1	-
6	EDO	A	718	-	-	0/1/1/1	-
6	EDO	A	713	-	-	0/1/1/1	-
7	PEG	B	403	-	-	3/4/4/4	-
7	PEG	A	719	-	-	3/4/4/4	-
6	EDO	A	711	-	-	0/1/1/1	-
6	EDO	A	714	-	-	1/1/1/1	-
6	EDO	A	717	-	-	0/1/1/1	-
7	PEG	A	720	-	-	3/4/4/4	-
3	MTA	A	701	-	-	0/7/23/23	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	702	A1BV5	C7-C8	-3.32	1.49	1.53
4	A	702	A1BV5	C22-N5	-2.47	1.35	1.39
4	A	702	A1BV5	C11-N4	2.13	1.39	1.34

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	702	A1BV5	C22-N5-C21	15.70	117.75	109.83
4	A	702	A1BV5	S-C21-N5	-8.56	111.64	117.01
4	A	702	A1BV5	C7-C8-N2	4.20	119.66	112.28
4	A	702	A1BV5	C10-N3-C11	3.30	121.83	118.64
4	A	702	A1BV5	O1-C7-C8	3.14	119.78	116.72
4	A	702	A1BV5	C9-C10-N3	-2.67	121.86	124.07
4	A	702	A1BV5	O2-C8-C7	-2.55	117.23	120.45
4	A	702	A1BV5	C15-C14-C12	-2.53	117.03	121.75
4	A	702	A1BV5	C19-C20-S	2.51	131.76	127.36
4	A	702	A1BV5	C16-C14-C12	-2.26	117.53	121.75

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	A	721	PGE	O3-C5-C6-O4
8	A	721	PGE	O2-C3-C4-O3
7	B	403	PEG	O1-C1-C2-O2
7	A	719	PEG	O1-C1-C2-O2
7	A	720	PEG	O1-C1-C2-O2
4	A	702	A1BV5	O1-C7-C8-N2
7	A	719	PEG	O2-C3-C4-O4
7	B	403	PEG	O2-C3-C4-O4
6	A	707	EDO	O1-C1-C2-O2
6	A	709	EDO	O1-C1-C2-O2
6	A	715	EDO	O1-C1-C2-O2
7	A	720	PEG	O2-C3-C4-O4
8	A	721	PGE	C1-C2-O2-C3
7	B	403	PEG	C1-C2-O2-C3
7	A	720	PEG	C1-C2-O2-C3
6	A	712	EDO	O1-C1-C2-O2
6	A	714	EDO	O1-C1-C2-O2
7	A	719	PEG	C4-C3-O2-C2
6	B	401	EDO	O1-C1-C2-O2

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	A	702	A1BV5	C23-C17-C5-C4

There are no ring outliers.

7 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	716	EDO	2	0
6	B	401	EDO	1	0
6	A	715	EDO	2	0
8	A	721	PGE	1	0
7	A	719	PEG	1	0
6	A	711	EDO	1	0
6	A	714	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	625/645 (96%)	-0.01	9 (1%) 73 73	27, 51, 95, 125	1 (0%)
2	B	302/350 (86%)	0.53	11 (3%) 46 44	50, 71, 100, 144	0
All	All	927/995 (93%)	0.16	20 (2%) 62 60	27, 60, 98, 144	1 (0%)

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	208	CYS	5.0
2	B	210	ALA	3.2
2	B	86	LEU	2.9
2	B	328	PRO	2.8
2	B	247	CYS	2.8
1	A	294	PRO	2.8
2	B	246	SER	2.7
2	B	211	PRO	2.7
2	B	248	VAL	2.6
2	B	326	VAL	2.5
2	B	268	VAL	2.5
1	A	277	PHE	2.4
1	A	529	MET	2.4
1	A	292	PRO	2.2
2	B	209	SER	2.2
1	A	174	THR	2.1
1	A	293	PRO	2.1
1	A	299	LEU	2.1
1	A	297	TYR	2.1
1	A	146	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.


6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	EDO	B	401	4/4	0.64	0.16	80,82,83,83	0
6	EDO	A	712	4/4	0.67	0.19	74,79,84,85	0
7	PEG	B	403	7/7	0.78	0.17	59,78,89,89	0
8	PGE	A	721	10/10	0.79	0.20	72,77,82,84	0
6	EDO	A	718	4/4	0.80	0.14	61,61,66,66	0
6	EDO	A	708	4/4	0.80	0.15	70,70,79,84	0
7	PEG	A	720	7/7	0.81	0.14	71,75,78,81	0
6	EDO	A	716	4/4	0.81	0.17	47,53,56,62	0
6	EDO	B	402	4/4	0.81	0.13	61,66,67,73	0
6	EDO	A	714	4/4	0.82	0.14	59,62,65,69	0
6	EDO	A	715	4/4	0.83	0.18	48,54,58,66	0
6	EDO	A	717	4/4	0.83	0.22	53,53,53,68	0
6	EDO	A	710	4/4	0.86	0.15	49,57,59,61	0
6	EDO	A	709	4/4	0.88	0.13	57,61,64,64	0
7	PEG	A	719	7/7	0.89	0.12	54,57,66,69	0
6	EDO	A	711	4/4	0.92	0.10	53,55,59,65	0
6	EDO	A	707	4/4	0.93	0.10	50,53,53,59	0
5	CL	A	705	1/1	0.94	0.08	67,67,67,67	0
5	CL	A	703	1/1	0.94	0.10	66,66,66,66	0
5	CL	A	706	1/1	0.95	0.08	57,57,57,57	0
5	CL	A	704	1/1	0.96	0.06	64,64,64,64	0
4	A1BV5	A	702	31/31	0.96	0.07	35,42,50,52	0
6	EDO	A	713	4/4	0.97	0.08	50,54,55,60	0
3	MTA	A	701	20/20	0.97	0.06	33,40,48,48	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different

orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.