



wwPDB NMR Structure Validation Summary Report ⓘ

Mar 12, 2026 – 02:06 PM UTC

PDB ID : 2N4U / pdb_00002n4u
BMRB ID : 25681
Title : NMR structure of Fbp28 WW domain E454Y mutant
Authors : Medina, J.; Macias, M.; Martin-Malpartida, P.; Scheraga, H.A.
Deposited on : 2015-07-01

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
BMRB Restraints Analysis : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 45%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	37	35% 30% 5% • 27%

2 Ensemble composition and analysis

This entry contains 20 models. Model 20 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:434-A:460 (27)	0.30	20

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters. No single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 4, 5, 6, 7, 8, 11, 12, 13, 14, 15, 17, 20
2	9, 10, 19
3	16, 18

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 603 atoms, of which 291 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Transcription elongation regulator 1.

Mol	Chain	Residues	Atoms					Trace
1	A	37	Total	C	H	N	O	0
			603	199	291	49	64	

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	454	TYR	GLU	engineered mutation	UNP O14776

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: Transcription elongation regulator 1



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 20. Colouring as in section 4.1 above.

- Molecule 1: Transcription elongation regulator 1



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 300 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNSSOLVE	structure solution	
CNSSOLVE	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	228
Number of shifts mapped to atoms	228
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	45%

6 Model quality [i](#)

6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	242	217	217	14±2
All	All	4840	4340	4340	282

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 31.

5 of 39 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:449:ASN:HB3	1:A:454:TYR:HD1	0.98	1.14	13	20
1:A:449:ASN:HB3	1:A:454:TYR:CD1	0.94	1.97	2	20
1:A:449:ASN:O	1:A:453:LEU:HA	0.89	1.67	9	20
1:A:439:LYS:HE2	1:A:443:GLY:HA2	0.80	1.51	10	1
1:A:448:TYR:HA	1:A:454:TYR:O	0.80	1.77	1	20

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	27/37 (73%)	25±1 (94±2%)	1±1 (3±3%)	1±0 (3±1%)	5	35
All	All	540/740 (73%)	506 (94%)	16 (3%)	18 (3%)	5	35

All 1 unique Ramachandran outliers are listed below.

Mol	Chain	Res	Type	Models (Total)
1	A	453	LEU	18

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	25/32 (78%)	21±1 (84±3%)	4±1 (16±3%)	4	40
All	All	500/640 (78%)	419 (84%)	81 (16%)	4	40

5 of 10 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	436	THR	20
1	A	449	ASN	14
1	A	453	LEU	14
1	A	437	GLU	11
1	A	439	LYS	8

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 45% for the well-defined parts and 44% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	228
Number of shifts mapped to atoms	228
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	56

7.1.2 Chemical shift referencing [i](#)

No chemical shift referencing corrections were calculated (not enough data).

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 45%, i.e. 170 atoms were assigned a chemical shift out of a possible 379. 0 out of 1 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	53/134 (40%)	44/54 (81%)	7/54 (13%)	2/26 (8%)
Sidechain	96/176 (55%)	91/111 (82%)	5/57 (9%)	0/8 (0%)
Aromatic	21/69 (30%)	19/32 (59%)	2/35 (6%)	0/2 (0%)
Overall	170/379 (45%)	154/197 (78%)	14/146 (10%)	2/36 (6%)

7.1.4 Statistically unusual chemical shifts ⓘ

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	455	SER	C	3.42	166.15 – 183.14	-100.8
1	A	456	THR	C	9.92	166.08 – 183.07	-96.9
1	A	453	LEU	C	1.55	167.56 – 186.66	-91.9
1	A	461	GLN	C	4.19	166.94 – 185.80	-91.3
1	A	457	TRP	C	2.50	166.32 – 186.14	-87.7
1	A	454	TYR	CD2	3.49	125.28 – 140.14	-87.0
1	A	457	TRP	CE3	1.50	111.58 – 129.41	-66.7
1	A	457	TRP	CD2	1.58	118.40 – 137.65	-65.7
1	A	457	TRP	CG	4.17	101.31 – 120.62	-55.3
1	A	459	LYS	CE	-0.13	37.57 – 46.21	-48.6
1	A	460	PRO	CA	1.81	55.85 – 70.84	-41.0
1	A	461	GLN	N	8.88	102.61 – 137.42	-31.9
1	A	456	THR	HG21	7.85	0.08 – 2.19	31.8
1	A	456	THR	HG22	7.85	0.08 – 2.19	31.8
1	A	456	THR	HG23	7.85	0.08 – 2.19	31.8
1	A	461	GLN	CG	2.10	28.36 – 39.21	-29.2
1	A	457	TRP	N	6.71	101.51 – 141.60	-28.6
1	A	462	GLU	CA	1.06	47.03 – 67.62	-27.3
1	A	458	GLU	CA	1.16	47.03 – 67.62	-27.3
1	A	459	LYS	CD	0.83	23.50 – 34.42	-25.8
1	A	461	GLN	CA	3.93	46.17 – 66.97	-25.3
1	A	460	PRO	N	3.57	108.67 – 162.11	-24.7
1	A	456	THR	N	2.97	91.89 – 138.78	-24.0
1	A	463	LEU	CG	1.58	21.37 – 32.19	-23.3
1	A	457	TRP	CA	7.11	45.21 – 70.26	-20.2
1	A	463	LEU	CB	7.44	33.11 – 51.34	-19.1
1	A	460	PRO	HG2	7.39	0.41 – 3.45	18.0
1	A	455	SER	HB2	8.29	2.61 – 5.13	17.5
1	A	457	TRP	HD1	1.88	5.46 – 8.81	-15.7
1	A	461	GLN	CB	1.84	20.34 – 37.98	-15.5
1	A	454	TYR	HB2	8.32	1.09 – 4.72	14.9
1	A	463	LEU	CD1	1.33	16.71 – 32.55	-14.7
1	A	463	LEU	CD2	1.36	15.73 – 32.47	-13.6
1	A	457	TRP	HB3	7.97	1.31 – 4.93	13.4
1	A	453	LEU	CD2	2.06	15.73 – 32.47	-13.2
1	A	456	THR	CG2	7.18	16.06 – 27.03	-13.1
1	A	462	GLU	H	1.14	5.45 – 11.20	-12.5
1	A	453	LEU	HD11	4.13	-0.61 – 2.12	12.4

Continued on next page...


Continued from previous page...

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	453	LEU	HD12	4.13	-0.61 – 2.12	12.4
1	A	453	LEU	HD13	4.13	-0.61 – 2.12	12.4
1	A	458	GLU	H	1.24	5.45 – 11.20	-12.3
1	A	453	LEU	CD1	7.02	16.71 – 32.55	-11.1
1	A	454	TYR	HE1	9.18	5.59 – 7.82	11.1
1	A	459	LYS	HG2	3.71	0.13 – 2.61	9.4
1	A	460	PRO	HD2	6.72	1.93 – 5.38	8.9
1	A	449	ASN	HB3	0.00	1.12 – 4.38	-8.4
1	A	460	PRO	HA	1.73	2.78 – 6.00	-8.2
1	A	459	LYS	HD2	-0.06	0.58 – 2.64	-8.1
1	A	454	TYR	HD1	4.62	5.49 – 8.39	-8.0
1	A	458	GLU	HB3	0.35	0.95 – 3.05	-7.9
1	A	454	TYR	HB3	5.83	0.93 – 4.76	7.8
1	A	462	GLU	HB3	0.65	0.95 – 3.05	-6.4
1	A	463	LEU	HB3	3.75	-0.26 – 3.31	6.2
1	A	458	GLU	HB2	0.82	1.00 – 3.05	-5.8
1	A	449	ASN	HD22	4.34	4.69 – 9.61	-5.7
1	A	459	LYS	HE2	1.89	1.95 – 3.88	-5.3

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:



8 NMR restraints analysis [i](#)

8.1 Conformationally restricting restraints [i](#)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	580
Intra-residue ($ i-j =0$)	0
Sequential ($ i-j =1$)	205
Medium range ($ i-j >1$ and $ i-j <5$)	117
Long range ($ i-j \geq 5$)	258
Inter-chain	0
Hydrogen bond restraints	0
Disulfide bond restraints	0
Total dihedral-angle restraints	0
Number of unmapped restraints	0
Number of restraints per residue	15.7
Number of long range restraints per residue ¹	7.0

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations [i](#)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model [i](#)

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	None	None
0.2-0.5 (Medium)	None	None
>0.5 (Large)	None	None

8.2.2 Average number of dihedral-angle violations per model ⓘ

Dihedral-angle violations less than 1° are not included in the calculation. There are no dihedral-angle violations

9 Distance violation analysis [i](#)


9.1 Summary of distance violations [i](#)

The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Restrains type	Count	% ¹	Violated ³			Consistently Violated ⁴		
			Count	% ²	% ¹	Count	% ²	% ¹
Intra-residue (i-j =0)	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sequential (i-j =1)	205	35.3	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	55	9.5	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	131	22.6	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	19	3.3	0	0.0	0.0	0	0.0	0.0
Medium range (i-j >1 & i-j <5)	117	20.2	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	15	2.6	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	53	9.1	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	49	8.4	0	0.0	0.0	0	0.0	0.0
Long range (i-j ≥5)	258	44.5	1	0.4	0.2	0	0.0	0.0
Backbone-Backbone	19	3.3	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	73	12.6	1	1.4	0.2	0	0.0	0.0
Sidechain-Sidechain	166	28.6	0	0.0	0.0	0	0.0	0.0
Inter-chain	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	580	100.0	1	0.2	0.2	0	0.0	0.0
Backbone-Backbone	89	15.3	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	257	44.3	1	0.4	0.2	0	0.0	0.0
Sidechain-Sidechain	234	40.3	0	0.0	0.0	0	0.0	0.0

¹ percentage calculated with respect to the total number of distance restraints, ² percentage calculated with respect to the number of restraints in a particular restraint category, ³ violated in at least one model, ⁴ violated in all the models

9.1.1 Bar chart : Distribution of distance restraints and violations [i](#)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model [i](#)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

Model ID	Number of violations						Mean (Å)	Max (Å)	SD ⁶ (Å)	Median (Å)
	IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total				
1	0	0	0	0	0	0	0.0	0.0	0.0	0.0
2	0	0	0	0	0	0	0.0	0.0	0.0	0.0
3	0	0	0	0	0	0	0.0	0.0	0.0	0.0
4	0	0	0	0	0	0	0.0	0.0	0.0	0.0
5	0	0	0	0	0	0	0.0	0.0	0.0	0.0
6	0	0	0	0	0	0	0.0	0.0	0.0	0.0
7	0	0	0	0	0	0	0.0	0.0	0.0	0.0
8	0	0	0	0	0	0	0.0	0.0	0.0	0.0
9	0	0	0	0	0	0	0.0	0.0	0.0	0.0
10	0	0	0	0	0	0	0.0	0.0	0.0	0.0

Continued on next page...


Continued from previous page...

Model ID	Number of violations						Mean (Å)	Max (Å)	SD ⁶ (Å)	Median (Å)
	IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total				
11	0	0	0	0	0	0	0.0	0.0	0.0	0.0
12	0	0	0	0	0	0	0.0	0.0	0.0	0.0
13	0	0	0	1	0	1	0.1	0.1	0.0	0.1
14	0	0	0	0	0	0	0.0	0.0	0.0	0.0
15	0	0	0	0	0	0	0.0	0.0	0.0	0.0
16	0	0	0	0	0	0	0.0	0.0	0.0	0.0
17	0	0	0	0	0	0	0.0	0.0	0.0	0.0
18	0	0	0	0	0	0	0.0	0.0	0.0	0.0
19	0	0	0	0	0	0	0.0	0.0	0.0	0.0
20	0	0	0	0	0	0	0.0	0.0	0.0	0.0

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints,

⁵Inter-chain restraints, ⁶Standard deviation

9.2.1 Bar graph : Distance Violation statistics for each model ⓘ



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right

9.3 Distance violation statistics for the ensemble


Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 579(IR:0, SQ:205, MR:117, LR:257, IC:0) restraints are not violated in the ensemble.

Number of violated restraints						Fraction of the ensemble	
IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total	Count ⁶	%
0	0	0	1	0	1	1	5.0
0	0	0	0	0	0	2	10.0
0	0	0	0	0	0	3	15.0
0	0	0	0	0	0	4	20.0
0	0	0	0	0	0	5	25.0
0	0	0	0	0	0	6	30.0
0	0	0	0	0	0	7	35.0
0	0	0	0	0	0	8	40.0
0	0	0	0	0	0	9	45.0
0	0	0	0	0	0	10	50.0
0	0	0	0	0	0	11	55.0
0	0	0	0	0	0	12	60.0
0	0	0	0	0	0	13	65.0
0	0	0	0	0	0	14	70.0
0	0	0	0	0	0	15	75.0
0	0	0	0	0	0	16	80.0
0	0	0	0	0	0	17	85.0
0	0	0	0	0	0	18	90.0
0	0	0	0	0	0	19	95.0
0	0	0	0	0	0	20	100.0

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints,

⁵Inter-chain restraints, ⁶ Number of models with violations

9.3.1 Bar graph : Distance violation statistics for the ensemble [i](#)



9.4 Most violated distance restraints in the ensemble [i](#)

No violations found

9.5 All violated distance restraints [i](#)

9.5.1 Histogram : Distribution of distance violations [i](#)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.

Data insufficient to plot histogram

9.5.2 Table : All distance violations [i](#)

The following table lists the absolute value of the violation for each restraint in the ensemble sorted by its value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,436)	1:449:A:ASN:HA	1:454:A:TYR:HE1	13	0.1

Continued on next page...

Continued from previous page...

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,436)	1:449:A:ASN:HA	1:454:A:TYR:HE2	13	0.1

10 Dihedral-angle violation analysis ⓘ

No dihedral-angle restraints found