



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 09:22 PM UTC

PDB ID : 3N42 / pdb_00003n42
Title : Crystal structures of the mature envelope glycoprotein complex (furin cleavage) of Chikungunya virus.
Authors : Voss, J.; Vaney, M.C.; Duquerroy, S.; Rey, F.A.
Deposited on : 2010-05-21
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	65	23% 63% 14% • 20%
2	B	360	6% 84% 9% • 6%
3	F	473	3% 72% 9% • 16%
4	C	2	50% 50%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6205 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called E3 envelope glycoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	52	Total	C	N	O	S	0	0	0
			406	253	64	80	9			

- Molecule 2 is a protein called E2 envelope glycoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	338	Total	C	N	O	S	0	0	0
			2669	1666	483	500	20			

- Molecule 3 is a protein called E1 envelope glycoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	F	395	Total	C	N	O	S	0	0	0
			3000	1896	503	577	24			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	-1	GLY	-	SEE REMARK 999	UNP Q1H8W5
F	0	GLY	-	SEE REMARK 999	UNP Q1H8W5

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	C	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	26	Total	O	0	0
			26	26		
6	F	62	Total	O	0	0
			62	62		



3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: E3 envelope glycoprotein



- Molecule 2: E2 envelope glycoprotein



- Molecule 3: E1 envelope glycoprotein



- Molecule 4: 2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	58.39Å 90.68Å 178.41Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.60 – 3.00 44.60 – 3.00	Depositor EDS
% Data completeness (in resolution range)	96.2 (44.60-3.00) 96.1 (44.60-3.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.96 (at 3.01Å)	Xtriage
Refinement program	BUSTER 2.9.3	Depositor
R, R_{free}	0.247 , 0.278 0.258 , 0.289	Depositor DCC
R_{free} test set	980 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	53.9	Xtriage
Anisotropy	0.475	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 47.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	6205	wwPDB-VP
Average B, all atoms (Å ²)	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, NDG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.78	0/415	1.31	0/565
2	B	0.77	0/2741	1.18	4/3731 (0.1%)
3	F	0.90	9/3077 (0.3%)	1.09	5/4197 (0.1%)
All	All	0.84	9/6233 (0.1%)	1.15	9/8493 (0.1%)

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	207	THR	CA-CB	-7.22	1.45	1.53
3	F	194	ALA	CA-CB	-6.78	1.44	1.53
3	F	207	THR	CA-C	-6.64	1.46	1.52
3	F	207	THR	C-O	-6.63	1.18	1.24
3	F	194	ALA	C-O	-5.77	1.19	1.24
3	F	191	PRO	CA-C	-5.48	1.44	1.52
3	F	213	VAL	CA-CB	-5.18	1.47	1.55
3	F	214	TYR	CA-C	-5.07	1.46	1.52
3	F	243	TRP	C-O	-5.03	1.17	1.24

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	30	SER	N-CA-C	6.16	117.57	109.93
2	B	209	GLY	CA-C-N	5.86	131.44	123.11
2	B	209	GLY	C-N-CA	5.86	131.44	123.11
2	B	100	PHE	N-CA-C	5.74	118.90	109.72
3	F	193	GLY	N-CA-C	-5.47	107.03	116.01
3	F	255	ALA	N-CA-C	5.40	116.50	109.64
3	F	266	VAL	N-CA-CB	5.31	116.84	110.31
3	F	190	PRO	O-C-N	5.30	127.71	121.46
3	F	245	LYS	N-CA-C	-5.06	106.20	112.93

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	406	0	386	5	0
2	B	2669	0	2588	10	0
3	F	3000	0	2903	31	0
4	C	28	0	24	1	0
5	B	14	0	13	0	0
6	B	26	0	0	0	0
6	F	62	0	0	0	0
All	All	6205	0	5914	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:F:245:LYS:HE2	3:F:246:GLU:OE2	1.79	0.82
3:F:243:TRP:O	3:F:247:ARG:HG2	1.84	0.77
1:A:27:TYR:OH	2:B:234:LYS:NZ	2.26	0.65
3:F:42:LEU:HD11	3:F:266:VAL:HG22	1.80	0.64
2:B:14:PRO:HG3	2:B:68:ARG:HG2	1.78	0.63
3:F:212:ASP:C	3:F:213:VAL:CG1	2.72	0.63
3:F:190:PRO:C	3:F:191:PRO:O	2.35	0.62
3:F:212:ASP:C	3:F:213:VAL:HG13	2.25	0.61
2:B:245:ASN:HB3	2:B:248:LEU:HD22	1.83	0.60
2:B:10:LYS:O	2:B:234:LYS:NZ	2.34	0.59
3:F:209:GLU:C	3:F:210:SER:O	2.39	0.59
3:F:263:THR:HG22	3:F:264:ASN:N	2.19	0.57
3:F:245:LYS:CE	3:F:246:GLU:OE2	2.53	0.57
3:F:326:GLY:O	3:F:346:VAL:HG22	2.06	0.56
3:F:207:THR:OG1	3:F:210:SER:HB2	2.07	0.55
3:F:242:TYR:CE1	3:F:246:GLU:HG3	2.42	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:F:190:PRO:O	3:F:191:PRO:O	2.27	0.52
2:B:196:THR:O	2:B:230:THR:HG23	2.09	0.52
3:F:207:THR:OG1	3:F:210:SER:N	2.45	0.50
3:F:60:VAL:HG22	3:F:102:GLN:HG3	1.94	0.50
2:B:230:THR:O	2:B:230:THR:OG1	2.30	0.49
3:F:209:GLU:O	3:F:210:SER:O	2.30	0.49
3:F:38:LEU:HB2	3:F:268:ALA:HB3	1.94	0.49
3:F:263:THR:CG2	3:F:264:ASN:H	2.26	0.48
2:B:52:LEU:HD22	2:B:67:LEU:HD21	1.96	0.48
3:F:207:THR:OG1	3:F:210:SER:CB	2.63	0.47
3:F:263:THR:CG2	3:F:264:ASN:N	2.77	0.47
1:A:16:PRO:HG2	1:A:19:GLN:HB2	1.96	0.46
3:F:263:THR:HG22	3:F:264:ASN:H	1.80	0.46
2:B:42:THR:HB	2:B:136:ILE:HD11	1.98	0.45
3:F:19:VAL:HB	3:F:27:MET:HB3	1.97	0.45
3:F:50:GLU:OE1	3:F:241:LYS:HE2	2.17	0.45
3:F:243:TRP:CE3	3:F:244:LEU:HD12	2.51	0.45
1:A:42:VAL:HG11	2:B:253:GLY:HA2	1.99	0.45
3:F:28:VAL:HG23	3:F:329:ALA:HB1	1.98	0.45
3:F:209:GLU:O	3:F:210:SER:C	2.62	0.42
3:F:59:TYR:HB3	3:F:103:LEU:HB3	2.01	0.41
3:F:141:ASN:HD22	4:C:1:NAG:H83	1.84	0.41
1:A:55:LEU:HD13	2:B:8:VAL:HA	2.03	0.41
3:F:296:LEU:HD12	3:F:369:VAL:O	2.21	0.41
3:F:332:SER:HB2	3:F:339:ILE:HG12	2.03	0.41
1:A:10:LEU:HB2	1:A:15:PHE:CE1	2.56	0.41
3:F:7:ILE:HG12	3:F:278:ILE:HD12	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	50/65 (77%)	46 (92%)	2 (4%)	2 (4%)	2	14
2	B	336/360 (93%)	310 (92%)	22 (6%)	4 (1%)	10	40
3	F	393/473 (83%)	378 (96%)	13 (3%)	2 (0%)	24	60
All	All	779/898 (87%)	734 (94%)	37 (5%)	8 (1%)	12	45

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	6	PHE
1	A	11	ALA
3	F	210	SER
2	B	73	HIS
2	B	164	THR
2	B	186	GLY
1	A	14	THR
3	F	191	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	49/61 (80%)	45 (92%)	4 (8%)	10	37
2	B	300/319 (94%)	284 (95%)	16 (5%)	20	54
3	F	329/378 (87%)	312 (95%)	17 (5%)	21	55
All	All	678/758 (89%)	641 (94%)	37 (6%)	19	53

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	LEU
1	A	10	LEU
1	A	14	THR
1	A	25	CYS
2	B	40	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	54	ILE
2	B	73	HIS
2	B	189	LYS
2	B	192	VAL
2	B	212	THR
2	B	215	LYS
2	B	230	THR
2	B	231	ASN
2	B	233	LYS
2	B	235	TRP
2	B	248	LEU
2	B	263	ASN
2	B	265	THR
2	B	280	LYS
2	B	313	HIS
3	F	55	ILE
3	F	120	SER
3	F	125	HIS
3	F	136	LEU
3	F	181	LYS
3	F	187	MET
3	F	209	GLU
3	F	213	VAL
3	F	221	LEU
3	F	241	LYS
3	F	244	LEU
3	F	247	ARG
3	F	327	LYS
3	F	345	GLU
3	F	347	GLU
3	F	364	GLU
3	F	370	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	12	ASN
1	A	49	GLN
2	B	73	HIS
2	B	183	GLN
2	B	184	GLN
2	B	193	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	224	GLN
2	B	236	GLN
2	B	245	ASN
2	B	282	GLN
2	B	307	GLN
3	F	102	GLN
3	F	138	GLN
3	F	152	HIS
3	F	199	GLN
3	F	331	HIS
3	F	351	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	C	1	3,4	14,14,15	1.21	1 (7%)	17,19,21	1.13	2 (11%)
4	NDG	C	2	4	14,14,15	1.54	1 (7%)	17,19,21	1.30	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	C	1	3,4	-	2/6/23/26	0/1/1/1
4	NDG	C	2	4	-	0/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	2	NDG	C1-C2	4.23	1.58	1.52
4	C	1	NAG	C1-C2	2.89	1.56	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	2	NDG	C1-C2-N2	3.60	116.10	110.43
4	C	1	NAG	C1-O5-C5	2.67	115.76	112.19
4	C	1	NAG	C2-N2-C7	2.64	126.43	122.90
4	C	2	NDG	C4-C3-C2	2.22	114.27	111.02

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	C	1	NAG	C8-C7-N2-C2
4	C	1	NAG	O7-C7-N2-C2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	B	1001	2	14,14,15	1.54	1 (7%)	17,19,21	1.46	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	B	1001	2	-	1/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	1001	NAG	C1-C2	4.42	1.58	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	1001	NAG	C1-O5-C5	3.78	117.25	112.19
5	B	1001	NAG	O5-C1-C2	2.35	114.93	111.29

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	1001	NAG	C1-C2-N2-C7

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	52/65 (80%)	1.51	15 (28%) 1 1	84, 101, 119, 126	0
2	B	338/360 (93%)	0.76	22 (6%) 25 13	33, 62, 83, 97	0
3	F	395/473 (83%)	0.50	12 (3%) 52 30	26, 46, 80, 106	0
All	All	785/898 (87%)	0.68	49 (6%) 26 14	26, 55, 95, 126	0

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	250	ASP	3.8
1	A	58	SER	3.6
1	A	9	LEU	3.6
2	B	187	ASN	3.5
1	A	8	CYS	3.4
3	F	393	SER	3.4
2	B	162	ALA	3.3
3	F	212	ASP	3.3
1	A	12	ASN	3.2
2	B	266	CYS	3.1
2	B	147	HIS	3.1
1	A	13	THR	3.0
2	B	212	THR	3.0
2	B	186	GLY	3.0
2	B	184	GLN	2.9
3	F	0	GLY	2.9
2	B	249	GLY	2.8
3	F	126	THR	2.8
2	B	205	GLY	2.7
2	B	342	LEU	2.7
1	A	10	LEU	2.6
1	A	7	MET	2.6
1	A	57	CYS	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	F	89	TRP	2.5
2	B	206	SER	2.4
2	B	201	CYS	2.4
3	F	68	CYS	2.4
3	F	376	CYS	2.4
2	B	160	THR	2.4
2	B	210	LEU	2.4
3	F	151	ASP	2.3
2	B	183	GLN	2.2
2	B	247	GLU	2.2
2	B	146	GLN	2.2
1	A	55	LEU	2.2
3	F	302	GLU	2.2
2	B	288	TYR	2.1
3	F	62	CYS	2.1
2	B	211	THR	2.1
1	A	17	CYS	2.1
2	B	223	ASP	2.1
1	A	21	PRO	2.1
1	A	11	ALA	2.0
1	A	25	CYS	2.0
3	F	96	CYS	2.0
3	F	114	CYS	2.0
1	A	20	PRO	2.0
1	A	22	CYS	2.0
2	B	216	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	C	1	14/15	-	-	51,54,57,57	0
4	NDG	C	2	14/15	-	-	57,61,64,64	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	NAG	B	1001	14/15	0.57	0.18	71,75,77,77	0

6.5 Other polymers [i](#)

There are no such residues in this entry.