



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 12:59 PM UTC

PDB ID : 3N43 / pdb_00003n43
Title : Crystal structures of the mature envelope glycoprotein complex (trypsin cleavage) of Chikungunya virus.
Authors : Voss, J.; Vaney, M.C.; Duquerroy, S.; Rey, F.A.
Deposited on : 2010-05-21
Resolution : 2.58 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.58 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4770 (2.60-2.56)
Clashscore	190562	5124 (2.60-2.56)
Ramachandran outliers	187476	5046 (2.60-2.56)
Sidechain outliers	187428	5046 (2.60-2.56)
RSRZ outliers	180081	4770 (2.60-2.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	65	74% 77% 22%
2	B	360	24% 88% 5% 7%
3	F	473	6% 79% 17%
4	C	2	100%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6238 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called E3 envelope glycoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	51	Total	C	N	O	S	0	0	0
			400	250	63	78	9			

- Molecule 2 is a protein called E2 envelope glycoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	336	Total	C	N	O	S	0	1	0
			2650	1653	480	497	20			

- Molecule 3 is a protein called E1 envelope glycoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	F	393	Total	C	N	O	S	0	0	0
			2989	1890	501	574	24			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
F	-1	GLY	-	SEE REMARK 999	UNP Q1H8W5
F	0	GLY	-	SEE REMARK 999	UNP Q1H8W5

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-alpha-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	C	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is ACETATE ION (CCD ID: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	F	1	Total	C	O	0	0
			4	2	2		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	46	Total 46	O 46	0	0
7	F	107	Total 107	O 107	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: E3 envelope glycoprotein



- Molecule 2: E2 envelope glycoprotein



- Molecule 3: E1 envelope glycoprotein



- Molecule 4: 2-acetamido-2-deoxy- α -D-glucopyranose-(1-4)-2-acetamido-2-deoxy- β -D-glucopyranose

Chain C:

100%

MAG1
MDG2

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	58.41Å 90.81Å 179.45Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.00 – 2.58 44.00 – 2.58	Depositor EDS
% Data completeness (in resolution range)	97.8 (44.00-2.58) 97.7 (44.00-2.58)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.13	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.91 (at 2.58Å)	Xtriage
Refinement program	BUSTER 2.9.3	Depositor
R, R_{free}	0.245 , 0.257 0.260 , 0.272	Depositor DCC
R_{free} test set	1532 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	34.6	Xtriage
Anisotropy	0.692	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 35.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	6238	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, NDG, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.73	0/409	1.22	0/557
2	B	0.74	0/2712	1.14	2/3691 (0.1%)
3	F	0.72	0/3066	1.07	1/4182 (0.0%)
All	All	0.73	0/6187	1.11	3/8430 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	88	SER	N-CA-C	-5.49	107.23	114.31
2	B	100	PHE	N-CA-C	5.41	118.55	109.95
3	F	382	PRO	N-CA-C	5.31	115.57	110.47

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	400	0	381	0	0
2	B	2650	0	2572	4	0
3	F	2989	0	2893	7	0
4	C	28	0	24	0	0
5	B	14	0	13	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	F	4	0	3	0	0
7	B	46	0	0	0	0
7	F	107	0	0	0	0
All	All	6238	0	5886	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:F:60:VAL:HG22	3:F:102:GLN:HG3	1.88	0.55
3:F:42:LEU:HD11	3:F:266:VAL:HG22	1.89	0.53
3:F:38:LEU:HB2	3:F:268:ALA:HB3	1.90	0.52
2:B:38:ARG:HB2	2:B:47:LYS:HB3	1.91	0.51
2:B:248:LEU:HD13	2:B:251:ARG:HB2	1.94	0.50
3:F:28:VAL:HG23	3:F:329:ALA:HB1	1.94	0.48
3:F:263:THR:HG22	3:F:266:VAL:HG12	2.00	0.43
3:F:41:THR:HB	3:F:125:HIS:HB2	2.00	0.42
2:B:13:ARG:HE	2:B:236:GLN:HE21	1.68	0.42
3:F:19:VAL:HB	3:F:27:MET:HB3	2.00	0.42
2:B:172:PRO:HG3	2:B:236:GLN:HB2	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	49/65 (75%)	46 (94%)	2 (4%)	1 (2%)	6	11
2	B	334/360 (93%)	317 (95%)	16 (5%)	1 (0%)	36	56

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	F	391/473 (83%)	383 (98%)	8 (2%)	0	100	100
All	All	774/898 (86%)	746 (96%)	26 (3%)	2 (0%)	36	56

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	11	ALA
2	B	72	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	48/61 (79%)	48 (100%)	0	100	100
2	B	297/319 (93%)	289 (97%)	8 (3%)	39	65
3	F	328/378 (87%)	323 (98%)	5 (2%)	57	78
All	All	673/758 (89%)	660 (98%)	13 (2%)	50	73

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	54	ILE
2	B	160	THR
2	B	192	VAL
2	B	206	SER
2	B	208	GLU
2	B	220	CYS
2	B	235	TRP
2	B	295	LEU
3	F	187	MET
3	F	209	GLU
3	F	346	VAL
3	F	352	LEU
3	F	370	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	49	GLN
2	B	29	HIS
2	B	183	GLN
2	B	224	GLN
2	B	236	GLN
2	B	281	ASN
2	B	282	GLN
2	B	299	ASN
3	F	102	GLN
3	F	331	HIS
3	F	351	GLN
3	F	373	GLN
3	F	381	HIS
3	F	389	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	C	1	4,3	14,14,15	1.24	1 (7%)	17,19,21	0.89	0
4	NDG	C	2	4	14,14,15	1.53	1 (7%)	17,19,21	1.41	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	C	1	4,3	-	0/6/23/26	0/1/1/1
4	NDG	C	2	4	-	1/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	2	NDG	C1-C2	4.23	1.58	1.52
4	C	1	NAG	C1-C2	3.29	1.56	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	2	NDG	C1-C2-N2	3.94	116.64	110.43

There are no chirality outliers.


All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	C	2	NDG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	B	2001	2	14,14,15	1.30	1 (7%)	17,19,21	0.91	1 (5%)
6	ACT	F	452	-	3,3,3	1.09	0	3,3,3	0.87	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	B	2001	2	-	1/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	2001	NAG	C1-C2	3.25	1.56	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	2001	NAG	C1-C2-N2	2.10	113.74	110.43

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	2001	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	51/65 (78%)	3.41	48 (94%) 0 0	78, 92, 110, 116	0
2	B	336/360 (93%)	1.46	85 (25%) 1 1	27, 49, 71, 95	0
3	F	393/473 (83%)	0.66	29 (7%) 20 17	19, 32, 55, 71	0
All	All	780/898 (86%)	1.18	162 (20%) 2 2	19, 41, 83, 116	0

All (162) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	9	LEU	8.7
1	A	10	LEU	5.8
1	A	31	PRO	5.1
2	B	206	SER	4.8
2	B	342	LEU	4.7
1	A	13	THR	4.6
1	A	7	MET	4.5
2	B	250	ASP	4.5
2	B	248	LEU	4.4
1	A	14	THR	4.3
2	B	93	ILE	4.3
1	A	15	PHE	4.3
1	A	11	ALA	4.3
2	B	231	ASN	4.3
1	A	23	THR	4.2
1	A	19	GLN	4.2
3	F	212	ASP	4.2
1	A	12	ASN	4.1
2	B	205	GLY	4.1
1	A	24	PRO	4.0
1	A	18	SER	4.0
2	B	247	GLU	4.0
1	A	16	PRO	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	43	MET	3.9
1	A	22	CYS	3.9
1	A	57	CYS	3.9
2	B	208	GLU	3.9
1	A	21	PRO	3.9
1	A	27	TYR	3.9
2	B	163	THR	3.9
2	B	73	HIS	3.8
1	A	29	LYS	3.8
2	B	185	SER	3.7
2	B	252	LYS	3.7
1	A	20	PRO	3.7
3	F	297	THR	3.6
3	F	-1	GLY	3.5
1	A	17	CYS	3.5
1	A	53	ALA	3.5
2	B	207	ASN	3.4
2	B	9	TYR	3.4
2	B	107	LYS	3.4
1	A	8	CYS	3.4
1	A	55	LEU	3.4
2	B	120	LYS	3.3
2	B	269	PRO	3.3
1	A	25	CYS	3.3
2	B	160	THR	3.3
2	B	23	GLY	3.3
1	A	56	THR	3.2
2	B	67	LEU	3.2
3	F	387	ILE	3.2
1	A	51	LEU	3.2
1	A	38	LEU	3.2
1	A	35	LEU	3.1
3	F	291	VAL	3.1
1	A	36	ARG	3.1
1	A	44	ARG	3.1
2	B	119	ARG	3.1
3	F	0	GLY	3.1
2	B	280	LYS	3.1
1	A	47	TYR	3.0
1	A	39	GLU	3.0
2	B	162	ALA	3.0
2	B	142	HIS	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	288	TYR	3.0
3	F	89	TRP	3.0
3	F	379	GLU	3.0
3	F	245	LYS	3.0
2	B	249	GLY	2.9
1	A	28	GLU	2.9
1	A	37	MET	2.9
2	B	219	ASN	2.8
2	B	223	ASP	2.8
2	B	144	ARG	2.8
1	A	26	CYS	2.8
3	F	375	HIS	2.8
2	B	255	ILE	2.8
3	F	108	VAL	2.7
2	B	184	GLN	2.7
1	A	42	VAL	2.7
2	B	113	VAL	2.7
2	B	92	THR	2.7
3	F	126	THR	2.7
2	B	147	HIS	2.7
1	A	54	SER	2.6
2	B	24	GLU	2.6
2	B	224	GLN	2.6
3	F	290	VAL	2.6
2	B	7	ASN	2.5
3	F	213	VAL	2.5
2	B	161	ALA	2.5
3	F	302	GLU	2.5
2	B	121	ILE	2.5
2	B	164	THR	2.5
1	A	45	PRO	2.4
2	B	276	VAL	2.4
2	B	204	GLY	2.4
2	B	56	ILE	2.4
2	B	75	PRO	2.4
2	B	270	LYS	2.4
2	B	272	ARG	2.4
2	B	84	PHE	2.4
2	B	215	LYS	2.4
3	F	209	GLU	2.4
3	F	325	LYS	2.4
3	F	196	ARG	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	203	CYS	2.4
2	B	100	PHE	2.4
3	F	390	TYR	2.4
1	A	40	ASP	2.3
2	B	212	THR	2.3
1	A	52	GLN	2.3
2	B	313	HIS	2.3
2	B	58	THR	2.3
2	B	65	THR	2.3
2	B	30	SER	2.3
2	B	122	SER	2.3
1	A	50	LEU	2.3
2	B	266	CYS	2.3
1	A	34	THR	2.3
2	B	25	GLY	2.3
2	B	158	GLN	2.2
2	B	151	LEU	2.2
2	B	210	LEU	2.2
2	B	220	CYS	2.2
3	F	288	THR	2.2
2	B	118	SER	2.2
3	F	309	SER	2.2
3	F	391	PRO	2.2
3	F	110	LYS	2.2
2	B	282	GLN	2.2
2	B	82	GLY	2.2
2	B	64	TRP	2.2
2	B	123	HIS	2.1
2	B	97	MET	2.1
1	A	30	GLU	2.1
3	F	362	SER	2.1
1	A	33	GLU	2.1
2	B	27	SER	2.1
2	B	159	SER	2.1
2	B	183	GLN	2.1
2	B	78	ALA	2.1
1	A	41	ASN	2.1
2	B	253	GLY	2.1
2	B	278	TYR	2.1
2	B	157	VAL	2.1
2	B	246	ALA	2.1
2	B	263	ASN	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	F	70	ASP	2.1
2	B	285	MET	2.1
3	F	336	ALA	2.0
2	B	83	LEU	2.0
2	B	202	ASN	2.0
3	F	360	LEU	2.0
2	B	233	LYS	2.0
3	F	123	ARG	2.0
1	A	46	GLY	2.0
2	B	200	LYS	2.0
3	F	175	ASN	2.0
2	B	323	THR	2.0
2	B	62	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	C	1	14/15	-	-	45,49,51,51	0
4	NDG	C	2	14/15	-	-	52,56,58,58	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	NAG	B	2001	14/15	0.54	0.23	73,77,80,80	0
6	ACT	F	452	4/4	0.91	0.12	41,41,41,41	0

6.5 Other polymers [i](#)

There are no such residues in this entry.