



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 09:04 AM UTC

PDB ID : 3N4D / pdb_00003n4d
Title : Crystal structure of Cg10062 inactivated by(R)-oxirane-2-carboxylate
Authors : Guo, Y.; Robertson, B.A.; Hackert, M.L.; Whitman, C.P.
Deposited on : 2010-05-21
Resolution : 2.54 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.54 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1091 (2.54-2.54)
Clashscore	190562	1120 (2.54-2.54)
Ramachandran outliers	187476	1106 (2.54-2.54)
Sidechain outliers	187428	1106 (2.54-2.54)
RSRZ outliers	180081	1091 (2.54-2.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	148	5% 55% 34% 7% ...
1	B	148	% 55% 36% ...
1	C	148	% 64% 32% ...
1	D	148	4% 54% 38% 6% ...
1	E	148	% 57% 36% 5% ...

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	148	
1	G	148	
1	H	148	
1	I	148	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
1	PR4	B	1	-	-	X	-
1	PR4	C	1	-	-	X	-
1	PR4	E	1	-	-	X	-
1	PR4	F	1	-	-	X	-
1	PR4	I	1	-	-	X	-

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 11026 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative tautomerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	145	Total	C	N	O	S	0	0	0
			1179	746	204	226	3			
1	B	145	Total	C	N	O	S	0	0	0
			1190	753	204	230	3			
1	C	146	Total	C	N	O	S	0	0	0
			1196	756	205	232	3			
1	D	145	Total	C	N	O	S	0	0	0
			1182	749	204	226	3			
1	E	145	Total	C	N	O	S	0	0	0
			1154	732	199	220	3			
1	F	145	Total	C	N	O	S	0	0	0
			1186	751	204	228	3			
1	G	145	Total	C	N	O	S	0	0	0
			1171	740	203	225	3			
1	H	140	Total	C	N	O	S	0	0	0
			1122	712	197	210	3			
1	I	145	Total	C	N	O	S	0	0	0
			1166	740	203	220	3			

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	53	Total	O	0	0
			53	53		
2	B	48	Total	O	0	0
			48	48		
2	C	50	Total	O	0	0
			50	50		
2	D	43	Total	O	0	0
			43	43		
2	E	55	Total	O	0	0
			55	55		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	F	59	Total 59	O 59	0	0
2	G	49	Total 49	O 49	0	0
2	H	49	Total 49	O 49	0	0
2	I	74	Total 74	O 74	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Putative tautomerase




• Molecule 1: Putative tautomerase



• Molecule 1: Putative tautomerase




• Molecule 1: Putative tautomerase





- Molecule 1: Putative tautomerase



- Molecule 1: Putative tautomerase



- Molecule 1: Putative tautomerase




- Molecule 1: Putative tautomerase



- Molecule 1: Putative tautomerase





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	57.10Å 131.00Å 179.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.00 – 2.54 45.00 – 2.54	Depositor EDS
% Data completeness (in resolution range)	100.0 (45.00-2.54) 99.9 (45.00-2.54)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.32 (at 2.01Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.223 , 0.243 0.224 , 0.244	Depositor DCC
R_{free} test set	4473 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	34.8	Xtriage
Anisotropy	0.013	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 53.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	11026	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 63.44 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 9.6669e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PR4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.79	1/1191 (0.1%)	0.97	3/1616 (0.2%)
1	B	0.67	1/1202 (0.1%)	0.90	3/1630 (0.2%)
1	C	0.52	0/1208	0.91	4/1638 (0.2%)
1	D	0.61	1/1194 (0.1%)	0.91	1/1620 (0.1%)
1	E	0.69	2/1166 (0.2%)	0.96	2/1587 (0.1%)
1	F	0.59	3/1198 (0.3%)	0.90	2/1625 (0.1%)
1	G	0.57	0/1183	0.95	3/1607 (0.2%)
1	H	0.58	0/1134	0.91	1/1543 (0.1%)
1	I	0.75	2/1178 (0.2%)	0.98	7/1601 (0.4%)
All	All	0.65	10/10654 (0.1%)	0.93	26/14467 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	C	0	1
1	E	0	1
1	G	0	1
1	I	0	1
All	All	0	5

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	37	TYR	C-O	-6.00	1.16	1.24
1	A	10	ILE	C-O	-5.59	1.17	1.24
1	B	32	ALA	CA-CB	-5.50	1.44	1.53
1	I	50	SER	C-O	-5.40	1.17	1.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	I	51	TYR	C-O	-5.31	1.17	1.24

The worst 5 of 26 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	125	GLU	N-CA-C	9.24	121.36	111.28
1	F	119	LEU	N-CA-C	-8.30	98.79	110.50
1	C	47	GLU	CA-C-N	6.78	127.34	119.47
1	C	47	GLU	C-N-CA	6.78	127.34	119.47
1	C	48	PRO	N-CA-C	6.66	123.46	113.81

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	1	PR4	Mainchain
1	C	1	PR4	Mainchain
1	E	1	PR4	Mainchain
1	G	1	PR4	Mainchain
1	I	1	PR4	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1179	0	1150	51	0
1	B	1190	0	1167	57	0
1	C	1196	0	1172	55	0
1	D	1182	0	1159	54	0
1	E	1154	0	1107	53	0
1	F	1186	0	1163	52	0
1	G	1171	0	1127	55	0
1	H	1122	0	1083	54	0
1	I	1166	0	1136	56	0
2	A	53	0	0	10	0
2	B	48	0	0	10	0
2	C	50	0	0	11	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	43	0	0	11	0
2	E	55	0	0	7	0
2	F	59	0	0	8	0
2	G	49	0	0	10	0
2	H	49	0	0	9	0
2	I	74	0	0	12	0
All	All	11026	0	10264	442	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

The worst 5 of 442 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:4:THR:HB	2:H:504:HOH:O	1.32	1.29
1:I:1:PR4:HABA	1:I:112:MET:HE1	1.33	1.05
1:F:1:PR4:HABA	1:F:112:MET:HE1	1.39	1.03
1:A:113:THR:HG22	2:C:161:HOH:O	1.58	1.03
1:C:1:PR4:HDA	1:C:1:PR4:CAD	1.81	1.03

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	143/148 (97%)	132 (92%)	5 (4%)	6 (4%)	2 1
1	B	143/148 (97%)	136 (95%)	4 (3%)	3 (2%)	5 5
1	C	144/148 (97%)	137 (95%)	6 (4%)	1 (1%)	18 25
1	D	143/148 (97%)	137 (96%)	5 (4%)	1 (1%)	18 25
1	E	143/148 (97%)	138 (96%)	3 (2%)	2 (1%)	9 11

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	F	143/148 (97%)	138 (96%)	3 (2%)	2 (1%)	9	11
1	G	143/148 (97%)	136 (95%)	5 (4%)	2 (1%)	9	11
1	H	138/148 (93%)	126 (91%)	10 (7%)	2 (1%)	9	11
1	I	143/148 (97%)	135 (94%)	5 (4%)	3 (2%)	5	5
All	All	1283/1332 (96%)	1215 (95%)	46 (4%)	22 (2%)	7	8

5 of 22 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	124	GLU
1	B	10	ILE
1	B	32	ALA
1	D	10	ILE
1	E	10	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	123/129 (95%)	115 (94%)	8 (6%)	15	22
1	B	126/129 (98%)	117 (93%)	9 (7%)	13	19
1	C	127/129 (98%)	121 (95%)	6 (5%)	23	36
1	D	124/129 (96%)	116 (94%)	8 (6%)	15	22
1	E	117/129 (91%)	108 (92%)	9 (8%)	12	17
1	F	125/129 (97%)	120 (96%)	5 (4%)	28	42
1	G	120/129 (93%)	115 (96%)	5 (4%)	26	40
1	H	114/129 (88%)	110 (96%)	4 (4%)	32	48
1	I	120/129 (93%)	116 (97%)	4 (3%)	33	50
All	All	1096/1161 (94%)	1038 (95%)	58 (5%)	20	31

5 of 58 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	113	THR
1	I	73	ARG
1	E	31	LEU
1	I	38	LEU
1	G	143	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 40 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	111	ASN
1	I	18	GLN
1	H	8	GLN
1	H	77	GLN
1	I	62	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

9 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	PR4	H	1	1	12,13,14	0.88	0	16,17,19	2.05	5 (31%)
1	PR4	G	1	1	12,13,14	1.15	1 (8%)	16,17,19	2.28	4 (25%)
1	PR4	B	1	1	12,13,14	1.29	1 (8%)	16,17,19	1.83	3 (18%)
1	PR4	D	1	1	12,13,14	0.87	0	16,17,19	2.64	7 (43%)
1	PR4	F	1	1	12,13,14	1.09	0	16,17,19	1.70	4 (25%)
1	PR4	A	1	1	12,13,14	1.21	1 (8%)	16,17,19	2.73	6 (37%)
1	PR4	I	1	1	12,13,14	1.21	1 (8%)	16,17,19	2.00	5 (31%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	PR4	E	1	1	12,13,14	1.34	1 (8%)	16,17,19	2.25	2 (12%)
1	PR4	C	1	1	12,13,14	0.76	0	16,17,19	1.77	6 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PR4	H	1	1	-	3/8/20/22	0/1/1/1
1	PR4	G	1	1	-	5/8/20/22	0/1/1/1
1	PR4	B	1	1	-	7/8/20/22	0/1/1/1
1	PR4	D	1	1	-	5/8/20/22	0/1/1/1
1	PR4	F	1	1	-	5/8/20/22	0/1/1/1
1	PR4	A	1	1	-	6/8/20/22	0/1/1/1
1	PR4	I	1	1	-	5/8/20/22	0/1/1/1
1	PR4	E	1	1	-	3/8/20/22	0/1/1/1
1	PR4	C	1	1	-	6/8/20/22	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	1	PR4	CAC-CAD	3.48	1.57	1.52
1	G	1	PR4	CAC-CAD	3.17	1.56	1.52
1	B	1	PR4	CAC-CAD	3.14	1.56	1.52
1	A	1	PR4	OAE-CAD	-2.14	1.23	1.30
1	I	1	PR4	OAE-CAD	-2.06	1.24	1.30

The worst 5 of 42 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	1	PR4	C-CA-N	-7.24	105.66	112.28
1	D	1	PR4	C-CA-N	-6.74	106.11	112.28
1	A	1	PR4	OAE-CAD-CAC	6.67	126.83	112.74
1	G	1	PR4	C-CA-N	-5.77	107.00	112.28
1	I	1	PR4	C-CA-N	-5.45	107.29	112.28

There are no chirality outliers.

5 of 45 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	1	PR4	CAC-CAB-N-CD
1	A	1	PR4	N-CAB-CAC-OAG
1	B	1	PR4	CAC-CAB-N-CD
1	B	1	PR4	N-CAB-CAC-OAG
1	C	1	PR4	CAC-CAB-N-CD

There are no ring outliers.

9 monomers are involved in 51 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	H	1	PR4	1	0
1	G	1	PR4	3	0
1	B	1	PR4	7	0
1	D	1	PR4	3	0
1	F	1	PR4	8	0
1	A	1	PR4	1	0
1	I	1	PR4	6	0
1	E	1	PR4	7	0
1	C	1	PR4	15	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	144/148 (97%)	0.19	7 (4%) 35 34	13, 31, 77, 79	0
1	B	144/148 (97%)	0.17	2 (1%) 73 74	18, 37, 66, 83	0
1	C	145/148 (97%)	0.10	2 (1%) 73 74	13, 35, 59, 65	0
1	D	144/148 (97%)	0.21	6 (4%) 40 41	11, 31, 87, 92	0
1	E	144/148 (97%)	0.04	1 (0%) 84 86	16, 34, 48, 56	0
1	F	144/148 (97%)	0.02	1 (0%) 84 86	16, 33, 53, 61	0
1	G	144/148 (97%)	0.10	4 (2%) 55 56	14, 30, 67, 70	0
1	H	139/148 (93%)	0.05	2 (1%) 73 74	15, 34, 58, 62	0
1	I	144/148 (97%)	-0.10	1 (0%) 84 86	10, 28, 56, 62	0
All	All	1292/1332 (96%)	0.09	26 (2%) 65 65	10, 33, 67, 92	0

The worst 5 of 26 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	145	GLY	3.7
1	D	145	GLY	3.6
1	H	140	LEU	3.5
1	A	120	MET	3.4
1	G	145	GLY	3.4

6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
1	PR4	C	1	13/14	0.73	0.17	28,30,40,41	0
1	PR4	F	1	13/14	0.85	0.14	27,28,34,38	0
1	PR4	A	1	13/14	0.88	0.12	27,29,40,43	0
1	PR4	D	1	13/14	0.91	0.10	30,31,36,36	0
1	PR4	B	1	13/14	0.91	0.10	27,29,34,38	0
1	PR4	I	1	13/14	0.91	0.10	24,25,28,30	0
1	PR4	G	1	13/14	0.92	0.11	29,30,36,36	0
1	PR4	E	1	13/14	0.93	0.09	23,25,33,36	0
1	PR4	H	1	13/14	0.94	0.09	26,28,33,33	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.