



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 10:51 AM UTC

PDB ID : 4N4U / pdb_00004n4u
Title : Crystal structure of ABC transporter solute binding protein BB0719 from *Bordetella bronchiseptica* RB50, TARGET EFI-510049
Authors : Patskovsky, Y.; Vetting, M.W.; Toro, R.; Bhosle, R.; Al Obaidi, N.; Stead, M.; Washington, E.; Scott Glenn, A.; Chowdhury, S.; Evans, B.; Hammonds, J.; Hillerich, B.; Love, J.; Seidel, R.D.; Imker, H.J.; Jacobson, M.P.; Gerlt, J.A.; Almo, S.C.; Enzyme Function Initiative (EFI)
Deposited on : 2013-10-08
Resolution : 1.57 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.57 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1094 (1.58-1.58)
Clashscore	190562	1105 (1.58-1.58)
Ramachandran outliers	187476	1082 (1.58-1.58)
Sidechain outliers	187428	1081 (1.58-1.58)
RSRZ outliers	180081	1094 (1.58-1.58)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	326	3% 89% 5% 7%
1	B	326	16% 84% 8% 7%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5226 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative ABC transporter periplasmic solute-binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	304	Total	C	N	O	S	0	11	0
			2371	1504	413	444	10			
1	B	302	Total	C	N	O	S	0	9	0
			2356	1495	412	439	10			

There are 46 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP Q7WPG5
A	2	HIS	-	expression tag	UNP Q7WPG5
A	3	HIS	-	expression tag	UNP Q7WPG5
A	4	HIS	-	expression tag	UNP Q7WPG5
A	5	HIS	-	expression tag	UNP Q7WPG5
A	6	HIS	-	expression tag	UNP Q7WPG5
A	7	HIS	-	expression tag	UNP Q7WPG5
A	8	SER	-	expression tag	UNP Q7WPG5
A	9	SER	-	expression tag	UNP Q7WPG5
A	10	GLY	-	expression tag	UNP Q7WPG5
A	11	VAL	-	expression tag	UNP Q7WPG5
A	12	ASP	-	expression tag	UNP Q7WPG5
A	13	LEU	-	expression tag	UNP Q7WPG5
A	14	GLY	-	expression tag	UNP Q7WPG5
A	15	THR	-	expression tag	UNP Q7WPG5
A	16	GLU	-	expression tag	UNP Q7WPG5
A	17	ASN	-	expression tag	UNP Q7WPG5
A	18	LEU	-	expression tag	UNP Q7WPG5
A	19	TYR	-	expression tag	UNP Q7WPG5
A	20	PHE	-	expression tag	UNP Q7WPG5
A	21	GLN	-	expression tag	UNP Q7WPG5
A	22	SER	-	expression tag	UNP Q7WPG5
A	23	MET	-	expression tag	UNP Q7WPG5
B	1	MET	-	expression tag	UNP Q7WPG5
B	2	HIS	-	initiating methionine	UNP Q7WPG5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	3	HIS	-	expression tag	UNP Q7WPG5
B	4	HIS	-	expression tag	UNP Q7WPG5
B	5	HIS	-	expression tag	UNP Q7WPG5
B	6	HIS	-	expression tag	UNP Q7WPG5
B	7	HIS	-	expression tag	UNP Q7WPG5
B	8	SER	-	expression tag	UNP Q7WPG5
B	9	SER	-	expression tag	UNP Q7WPG5
B	10	GLY	-	expression tag	UNP Q7WPG5
B	11	VAL	-	expression tag	UNP Q7WPG5
B	12	ASP	-	expression tag	UNP Q7WPG5
B	13	LEU	-	expression tag	UNP Q7WPG5
B	14	GLY	-	expression tag	UNP Q7WPG5
B	15	THR	-	expression tag	UNP Q7WPG5
B	16	GLU	-	expression tag	UNP Q7WPG5
B	17	ASN	-	expression tag	UNP Q7WPG5
B	18	LEU	-	expression tag	UNP Q7WPG5
B	19	TYR	-	expression tag	UNP Q7WPG5
B	20	PHE	-	expression tag	UNP Q7WPG5
B	21	GLN	-	expression tag	UNP Q7WPG5
B	22	SER	-	expression tag	UNP Q7WPG5
B	23	MET	-	expression tag	UNP Q7WPG5

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	331	Total 334	O 334	0	3
3	B	158	Total 159	O 159	0	1


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative ABC transporter periplasmic solute-binding protein



- Molecule 1: Putative ABC transporter periplasmic solute-binding protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	59.57Å 68.53Å 153.01Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.57 50.00 – 1.57	Depositor EDS
% Data completeness (in resolution range)	99.8 (50.00-1.57) 99.8 (50.00-1.57)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.07 (at 1.57Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.176 , 0.212 0.187 , 0.221	Depositor DCC
R_{free} test set	2661 reflections (3.02%)	wwPDB-VP
Wilson B-factor (Å ²)	22.1	Xtriage
Anisotropy	0.501	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 45.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5226	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.74	0/2448	1.01	6/3303 (0.2%)
1	B	0.65	2/2427 (0.1%)	0.92	5/3273 (0.2%)
All	All	0.70	2/4875 (0.0%)	0.96	11/6576 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	307[A]	ARG	CA-C	8.96	1.64	1.52
1	B	307[B]	ARG	CA-C	8.96	1.64	1.52

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	177[A]	ARG	CA-C-O	16.10	138.29	120.10
1	A	177[B]	ARG	CA-C-O	16.10	138.29	120.10
1	A	177[A]	ARG	N-CA-C	11.68	125.14	111.71
1	A	177[B]	ARG	N-CA-C	11.68	125.14	111.71
1	B	307[A]	ARG	CA-C-O	9.87	131.01	120.55
1	B	307[B]	ARG	CA-C-O	9.87	131.01	120.55
1	B	307[A]	ARG	N-CA-C	8.61	120.67	111.28
1	B	307[B]	ARG	N-CA-C	8.61	120.67	111.28
1	B	306	ALA	O-C-N	5.84	128.32	122.12
1	A	142	PHE	CA-C-N	-5.00	113.58	119.84
1	A	142	PHE	C-N-CA	-5.00	113.58	119.84

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2371	0	2400	12	0
1	B	2356	0	2393	14	0
2	A	6	0	8	0	0
3	A	334	0	0	5	0
3	B	159	0	0	3	0
All	All	5226	0	4801	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:77[A]:GLN:HG3	3:A:530:HOH:O	1.82	0.78
1:B:109:ARG:NH2	1:B:295:THR:OG1	2.29	0.66
1:A:216:MET:HG2	3:B:445:HOH:O	1.96	0.65
1:B:149:SER:HB2	1:B:153:LEU:HD22	1.81	0.61
1:B:149:SER:CB	1:B:153:LEU:HD22	2.33	0.59
1:B:34[B]:SER:HB3	1:B:36:ASN:OD1	2.10	0.52
1:A:165:ARG:C	1:A:165:ARG:HD3	2.37	0.50
1:A:152:LYS:HE2	3:A:561:HOH:O	2.12	0.49
1:A:51:GLU:HG2	1:A:59:LYS:HA	1.94	0.48
1:A:152:LYS:CE	3:A:561:HOH:O	2.63	0.46
1:B:246:MET:HA	1:B:246:MET:HE2	1.96	0.46
1:A:143:PRO:HD3	1:A:265:ILE:HG22	1.98	0.46
1:B:25:THR:CG2	1:B:61:ARG:HG2	2.45	0.46
1:B:114:PHE:CZ	1:B:139:HIS:CE1	3.05	0.45
1:B:43:MET:HE1	1:B:87:ALA:HA	1.98	0.44
1:B:127:ARG:O	1:B:131[B]:LYS:HG2	2.18	0.43
1:B:25:THR:HG21	1:B:61:ARG:HG2	2.01	0.43
1:A:85:ARG:NH2	3:A:740:HOH:O	2.52	0.42
1:A:279:LYS:HE2	3:B:521:HOH:O	2.20	0.42
1:A:85:ARG:CZ	3:A:740:HOH:O	2.68	0.41
1:A:27:MET:HE2	1:A:27:MET:HA	2.02	0.41
1:B:163[A]:LYS:NZ	3:B:487:HOH:O	2.46	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:27:MET:HG3	1:B:84:VAL:HG12	2.01	0.41
1:B:148:PHE:HB3	1:B:217:LEU:HD13	2.03	0.41
1:A:290:MET:HB2	1:A:290:MET:HE3	1.91	0.41
1:B:151:ARG:O	1:B:152:LYS:C	2.65	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	313/326 (96%)	308 (98%)	5 (2%)	0	100	100
1	B	309/326 (95%)	302 (98%)	7 (2%)	0	100	100
All	All	622/652 (95%)	610 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	245/256 (96%)	244 (100%)	1 (0%)	84	75
1	B	244/256 (95%)	239 (98%)	5 (2%)	48	20
All	All	489/512 (96%)	483 (99%)	6 (1%)	61	40

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	39	ASN
1	B	151	ARG
1	B	155	SER
1	B	211	LYS
1	B	279	LYS
1	B	308	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	71	GLN
1	A	111	GLN
1	A	257	GLN
1	A	315	GLN
1	B	176	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	A	401	-	5,5,5	0.19	0	5,5,5	0.36	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	401	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	304/326 (93%)	0.12	10 (3%) 49 62	13, 22, 40, 61	11 (3%)
1	B	302/326 (92%)	0.91	53 (17%) 4 6	16, 37, 71, 92	9 (2%)
All	All	606/652 (92%)	0.52	63 (10%) 11 19	13, 28, 62, 92	20 (3%)

All (63) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	185	LEU	5.6
1	B	293	ALA	5.3
1	B	288	PRO	5.3
1	A	24	ALA	5.2
1	B	183	VAL	4.6
1	B	289	ALA	4.6
1	A	72	ALA	4.2
1	B	297	LEU	4.1
1	B	156	LEU	3.9
1	A	186	GLY	3.9
1	B	284	VAL	3.9
1	B	212	ILE	3.9
1	B	157	ALA	3.9
1	B	198	THR	3.6
1	B	199	ILE	3.6
1	B	219	TYR	3.3
1	B	282	VAL	3.3
1	B	190	VAL	3.2
1	B	196	ALA	3.1
1	B	193	ALA	3.1
1	B	195	SER	3.0
1	B	155	SER	3.0
1	A	23	MET	2.9
1	A	184	THR	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	287	THR	2.9
1	A	36	ASN	2.9
1	B	159	ILE	2.9
1	B	281	MET	2.9
1	B	276	LEU	2.9
1	B	187	GLY	2.9
1	B	294	ALA	2.9
1	B	283	ILE	2.9
1	B	290	MET	2.9
1	B	150	ALA	2.8
1	B	188	ALA	2.8
1	B	36	ASN	2.8
1	B	25	THR	2.8
1	B	191	PRO	2.8
1	B	217	LEU	2.7
1	B	280	GLY	2.7
1	A	189	GLU	2.7
1	B	181	ILE	2.7
1	A	25	THR	2.6
1	B	278	GLN	2.6
1	B	218	LYS	2.6
1	B	194	LEU	2.6
1	A	187	GLY	2.6
1	B	153	LEU	2.5
1	B	285	PRO	2.4
1	B	221	LEU	2.4
1	B	180	GLY	2.2
1	B	152	LYS	2.2
1	B	185	LEU	2.2
1	B	216	MET	2.2
1	B	72	ALA	2.2
1	B	215	ASP	2.2
1	B	279	LYS	2.1
1	B	160	LYS	2.1
1	B	182	PRO	2.1
1	B	207	ALA	2.1
1	B	275	ALA	2.0
1	B	292	GLN	2.0
1	B	298	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GOL	A	401	6/6	0.72	0.16	33,35,42,49	0

6.5 Other polymers [i](#)

There are no such residues in this entry.