



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 02:51 PM EDT

PDB ID : 6N4N / pdb_00006n4n
Title : Crystal structure of the designed protein DNCR2/danoprevir/NS3a complex
Authors : Wang, Z.; Foight, G.W.; Baker, D.; Maly, D.J.
Deposited on : 2018-11-19
Resolution : 2.29 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.29 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	197	3% 96% ..
1	B	197	3% 96% ..
2	C	229	9% 77% 20%
2	F	229	12% 79% 17%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6161 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NS3 protease.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	195	Total	C	N	O	S	0	0	0
			1448	894	266	282	6			
1	B	195	Total	C	N	O	S	0	0	0
			1448	894	266	282	6			

There are 58 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	986	MET	-	expression tag	UNP A0A0B4WYC6
A	987	LYS	-	expression tag	UNP A0A0B4WYC6
A	988	LYS	-	expression tag	UNP A0A0B4WYC6
A	989	LYS	-	expression tag	UNP A0A0B4WYC6
A	990	GLY	-	expression tag	UNP A0A0B4WYC6
A	991	SER	-	expression tag	UNP A0A0B4WYC6
A	992	VAL	-	expression tag	UNP A0A0B4WYC6
A	993	VAL	-	expression tag	UNP A0A0B4WYC6
A	994	ILE	-	expression tag	UNP A0A0B4WYC6
A	995	VAL	-	expression tag	UNP A0A0B4WYC6
A	996	GLY	-	expression tag	UNP A0A0B4WYC6
A	997	ARG	-	expression tag	UNP A0A0B4WYC6
A	998	ILE	-	expression tag	UNP A0A0B4WYC6
A	999	ASN	-	expression tag	UNP A0A0B4WYC6
A	1000	LEU	-	expression tag	UNP A0A0B4WYC6
A	1001	SER	-	expression tag	UNP A0A0B4WYC6
A	1002	GLY	-	expression tag	UNP A0A0B4WYC6
A	1003	ASP	-	expression tag	UNP A0A0B4WYC6
A	1013	GLU	LEU	conflict	UNP A0A0B4WYC6
A	1014	GLU	LEU	conflict	UNP A0A0B4WYC6
A	1017	GLN	ILE	conflict	UNP A0A0B4WYC6
A	1018	GLU	ILE	conflict	UNP A0A0B4WYC6
A	1021	GLN	LEU	conflict	UNP A0A0B4WYC6
A	1047	SER	CYS	conflict	UNP A0A0B4WYC6
A	1052	LEU	CYS	conflict	UNP A0A0B4WYC6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	1072	THR	ILE	conflict	UNP A0A0B4WYC6
A	1086	GLN	PRO	conflict	UNP A0A0B4WYC6
A	1139	ALA	SER	conflict	UNP A0A0B4WYC6
A	1159	SER	CYS	conflict	UNP A0A0B4WYC6
B	986	MET	-	expression tag	UNP A0A0B4WYC6
B	987	LYS	-	expression tag	UNP A0A0B4WYC6
B	988	LYS	-	expression tag	UNP A0A0B4WYC6
B	989	LYS	-	expression tag	UNP A0A0B4WYC6
B	990	GLY	-	expression tag	UNP A0A0B4WYC6
B	991	SER	-	expression tag	UNP A0A0B4WYC6
B	992	VAL	-	expression tag	UNP A0A0B4WYC6
B	993	VAL	-	expression tag	UNP A0A0B4WYC6
B	994	ILE	-	expression tag	UNP A0A0B4WYC6
B	995	VAL	-	expression tag	UNP A0A0B4WYC6
B	996	GLY	-	expression tag	UNP A0A0B4WYC6
B	997	ARG	-	expression tag	UNP A0A0B4WYC6
B	998	ILE	-	expression tag	UNP A0A0B4WYC6
B	999	ASN	-	expression tag	UNP A0A0B4WYC6
B	1000	LEU	-	expression tag	UNP A0A0B4WYC6
B	1001	SER	-	expression tag	UNP A0A0B4WYC6
B	1002	GLY	-	expression tag	UNP A0A0B4WYC6
B	1003	ASP	-	expression tag	UNP A0A0B4WYC6
B	1013	GLU	LEU	conflict	UNP A0A0B4WYC6
B	1014	GLU	LEU	conflict	UNP A0A0B4WYC6
B	1017	GLN	ILE	conflict	UNP A0A0B4WYC6
B	1018	GLU	ILE	conflict	UNP A0A0B4WYC6
B	1021	GLN	LEU	conflict	UNP A0A0B4WYC6
B	1047	SER	CYS	conflict	UNP A0A0B4WYC6
B	1052	LEU	CYS	conflict	UNP A0A0B4WYC6
B	1072	THR	ILE	conflict	UNP A0A0B4WYC6
B	1086	GLN	PRO	conflict	UNP A0A0B4WYC6
B	1139	ALA	SER	conflict	UNP A0A0B4WYC6
B	1159	SER	CYS	conflict	UNP A0A0B4WYC6


- Molecule 2 is a protein called Rosetta-designed danoprevir/NS3a complex reader 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	184	Total	C	N	O	S	0	0	0
			1432	872	260	299	1			
2	F	191	Total	C	N	O	S	0	0	0
			1490	904	275	310	1			

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		
3	B	1	Total	Zn	0	0
			1	1		

- Molecule 4 is (2R,6S,12Z,13aS,14aR,16aS)-6-[(tert-butoxycarbonyl)amino]-14a-[(cyclopropylsulfonyl)carbamoyl]-5,16-dioxo-1,2,3,5,6,7,8,9,10,11,13a,14,14a,15,16,16a-hexadecahydrocyclopropa[e]pyrrolo[1,2-a][1,4]diazacyclopentadecin-2-yl 4-fluoro-2H-isoindole-2-carboxylate (CCD ID: TSV) (formula: C₃₅H₄₄FN₅O₉S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total	C	F	N	O	S	0	0
			51	35	1	5	9	1		
4	B	1	Total	C	F	N	O	S	0	0
			51	35	1	5	9	1		

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	C	1	Total	O	S	0	0
			5	4	1		
5	F	1	Total	O	S	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	65	Total	O	0	0
			65	65		
6	B	63	Total	O	0	0
			63	63		
6	C	37	Total	O	0	0
			37	37		
6	F	44	Total	O	0	0
			44	44		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: NS3 protease





- Molecule 1: NS3 protease



- Molecule 2: Rosetta-designed danoprevir/NS3a complex reader 2



- Molecule 2: Rosetta-designed danoprevir/NS3a complex reader 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	70.84Å 69.26Å 99.34Å 90.00° 108.59° 90.00°	Depositor
Resolution (Å)	94.16 – 2.29 94.16 – 2.29	Depositor EDS
% Data completeness (in resolution range)	98.3 (94.16-2.29) 98.3 (94.16-2.29)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.08 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.203 , 0.241 0.204 , 0.241	Depositor DCC
R_{free} test set	1979 reflections (4.80%)	wwPDB-VP
Wilson B-factor (Å ²)	33.4	Xtriage
Anisotropy	0.040	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 32.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.016 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6161	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TSV, ZN, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.73	0/1473	0.90	1/2000 (0.1%)
1	B	0.73	0/1473	0.89	1/2000 (0.1%)
2	C	0.65	0/1440	1.04	0/1945
2	F	0.67	0/1500	1.07	0/2027
All	All	0.70	0/5886	0.98	2/7972 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1005	ALA	N-CA-C	5.55	118.06	110.24
1	A	1005	ALA	N-CA-C	5.53	118.04	110.24

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1448	0	1452	4	0
1	B	1448	0	1452	2	0
2	C	1432	0	1418	1	0
2	F	1490	0	1476	4	0
3	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	1	0	0	0	0
4	A	51	0	44	1	0
4	B	51	0	44	0	0
5	A	10	0	0	0	0
5	B	10	0	0	0	0
5	C	5	0	0	0	0
5	F	5	0	0	0	0
6	A	65	0	0	0	0
6	B	63	0	0	0	0
6	C	37	0	0	0	0
6	F	44	0	0	0	0
All	All	6161	0	5886	9	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (9) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:27:THR:HG21	2:F:32:VAL:HG11	1.85	0.58
1:A:1161:ARG:HD3	2:F:82:ARG:NH2	2.31	0.44
2:F:65:LEU:HG	2:F:119:LEU:HD23	2.00	0.44
1:A:1104:LEU:HD22	1:A:1151:VAL:HG21	2.01	0.42
4:A:1202:TSV:H13A	4:A:1202:TSV:H3	1.93	0.42
1:B:1104:LEU:HD22	1:B:1151:VAL:HG21	2.02	0.41
1:A:1072:THR:HG21	1:B:1061:THR:O	2.20	0.41
1:A:1161:ARG:HD3	2:F:82:ARG:CZ	2.51	0.40
2:C:65:LEU:HG	2:C:119:LEU:HD23	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	193/197 (98%)	190 (98%)	3 (2%)	0	100	100
1	B	193/197 (98%)	190 (98%)	3 (2%)	0	100	100
2	C	180/229 (79%)	176 (98%)	4 (2%)	0	100	100
2	F	189/229 (82%)	184 (97%)	5 (3%)	0	100	100
All	All	755/852 (89%)	740 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	158/160 (99%)	157 (99%)	1 (1%)	78	89
1	B	158/160 (99%)	157 (99%)	1 (1%)	78	89
2	C	145/184 (79%)	140 (97%)	5 (3%)	32	49
2	F	152/184 (83%)	148 (97%)	4 (3%)	40	59
All	All	613/688 (89%)	602 (98%)	11 (2%)	51	70

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1001	SER
1	B	1001	SER
2	C	11	GLU
2	C	22	GLU
2	C	59	VAL
2	C	63	LEU
2	C	157	GLU
2	F	11	GLU
2	F	22	GLU
2	F	59	VAL
2	F	87	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	1027	ASN
1	B	1017	GLN
1	B	1027	ASN
1	B	1049	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	SO4	C	301	-	4,4,4	0.43	0	6,6,6	0.10	0
4	TSV	B	1202	-	49,56,56	2.19	10 (20%)	69,85,85	2.33	15 (21%)
4	TSV	A	1202	-	49,56,56	2.16	9 (18%)	69,85,85	2.28	15 (21%)
5	SO4	F	301	-	4,4,4	0.45	0	6,6,6	0.15	0
5	SO4	B	1204	-	4,4,4	0.44	0	6,6,6	0.07	0
5	SO4	A	1203	-	4,4,4	0.40	0	6,6,6	0.31	0
5	SO4	A	1204	-	4,4,4	0.46	0	6,6,6	0.12	0
5	SO4	B	1203	-	4,4,4	0.38	0	6,6,6	0.24	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	TSV	B	1202	-	-	5/59/83/83	0/5/6/6
4	TSV	A	1202	-	-	7/59/83/83	0/5/6/6

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	1202	TSV	C12-N4	6.68	1.48	1.38
4	A	1202	TSV	C12-N4	6.64	1.48	1.38
4	B	1202	TSV	C31-N4	6.49	1.48	1.38
4	A	1202	TSV	C31-N4	6.16	1.47	1.38
4	B	1202	TSV	O1-C20	6.10	1.46	1.34
4	A	1202	TSV	O1-C20	6.02	1.46	1.34
4	A	1202	TSV	O8-C35	5.45	1.45	1.34
4	B	1202	TSV	O8-C35	5.26	1.44	1.34
4	B	1202	TSV	O4-S1	3.99	1.48	1.43
4	A	1202	TSV	C17-C21	3.72	1.41	1.36
4	A	1202	TSV	C35-N4	-3.65	1.34	1.40
4	A	1202	TSV	O4-S1	3.54	1.47	1.43
4	B	1202	TSV	C35-N4	-3.50	1.34	1.40
4	B	1202	TSV	C17-C21	3.30	1.40	1.36
4	B	1202	TSV	O6-S1	3.23	1.47	1.43
4	A	1202	TSV	O6-S1	2.68	1.46	1.43
4	A	1202	TSV	O1-C9	-2.61	1.43	1.48
4	B	1202	TSV	O1-C9	-2.38	1.44	1.48
4	B	1202	TSV	C3-C5	2.04	1.52	1.49

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	1202	TSV	C4-C2-N2	8.12	124.77	116.10
4	A	1202	TSV	C4-C2-N2	7.88	124.52	116.10
4	B	1202	TSV	O6-S1-O4	-6.74	109.04	118.88
4	B	1202	TSV	C3-C2-N2	-5.92	106.21	117.25
4	B	1202	TSV	C17-C21-C26	-5.77	118.48	123.59
4	A	1202	TSV	C17-C21-C26	-5.68	118.57	123.59
4	A	1202	TSV	C3-C2-N2	-5.59	106.82	117.25
4	A	1202	TSV	C12-N4-C31	5.44	114.80	108.92
4	A	1202	TSV	O6-S1-O4	-5.44	110.94	118.88

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	1202	TSV	C12-N4-C31	4.88	114.19	108.92
4	B	1202	TSV	O1-C20-N1	4.73	117.78	110.03
4	A	1202	TSV	O1-C20-N1	4.71	117.74	110.03
4	A	1202	TSV	C26-C31-N4	-4.43	102.23	108.84
4	A	1202	TSV	C24-C12-N4	-4.38	102.76	108.44
4	B	1202	TSV	C24-C12-N4	-4.18	103.03	108.44
4	B	1202	TSV	C26-C31-N4	-4.17	102.62	108.84
4	A	1202	TSV	O1-C20-O2	-4.08	118.39	125.64
4	B	1202	TSV	C9-O1-C20	3.95	126.96	120.97
4	B	1202	TSV	O1-C20-O2	-3.93	118.65	125.64
4	A	1202	TSV	C9-O1-C20	3.87	126.84	120.97
4	A	1202	TSV	C1-C2-N2	-3.81	112.74	117.77
4	B	1202	TSV	C1-C2-N2	-3.29	113.43	117.77
4	B	1202	TSV	O8-C35-O5	-3.08	118.56	124.92
4	A	1202	TSV	C27-C15-C11	3.01	117.11	111.36
4	A	1202	TSV	O8-C35-O5	-2.96	118.80	124.92
4	A	1202	TSV	O4-S1-C23	2.73	111.62	107.98
4	B	1202	TSV	C27-C15-C11	2.34	115.85	111.36
4	B	1202	TSV	O4-S1-C23	2.20	110.91	107.98
4	A	1202	TSV	C14-C16-C24	-2.13	117.26	120.16
4	B	1202	TSV	O5-C35-N4	2.05	125.36	122.59

There are no chirality outliers.

All (12) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	B	1202	TSV	C4-N3-S1-C23
4	A	1202	TSV	C18-C19-C22-C25
4	B	1202	TSV	C18-C19-C22-C25
4	A	1202	TSV	C4-N3-S1-O6
4	A	1202	TSV	C1-C3-C5-C10
4	B	1202	TSV	C1-C3-C5-C10
4	A	1202	TSV	C4-N3-S1-C23
4	A	1202	TSV	C2-C3-C5-C10
4	B	1202	TSV	C2-C3-C5-C10
4	A	1202	TSV	N2-C2-C4-O9
4	B	1202	TSV	N2-C11-C15-N5
4	A	1202	TSV	C13-C18-C19-C22


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1202	TSV	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	195/197 (98%)	0.05	6 (3%) 51 53	18, 30, 56, 81	0
1	B	195/197 (98%)	0.05	5 (2%) 57 59	20, 33, 59, 85	0
2	C	184/229 (80%)	0.74	20 (10%) 10 12	21, 50, 116, 141	0
2	F	191/229 (83%)	0.70	28 (14%) 6 6	20, 47, 117, 147	0
All	All	765/852 (89%)	0.38	59 (7%) 19 21	18, 37, 95, 147	0

All (59) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1004	THR	5.9
2	C	59	VAL	5.5
2	C	162	VAL	5.3
1	A	1004	THR	5.3
2	F	59	VAL	4.9
1	B	1001	SER	4.2
1	A	1001	SER	4.2
2	F	55	SER	3.8
2	C	54	PRO	3.8
1	A	1003	ASP	3.7
1	B	1003	ASP	3.7
2	F	54	PRO	3.7
2	F	27	THR	3.6
2	F	58	ASP	3.4
2	F	57	SER	3.2
1	A	1002	GLY	3.1
2	F	6	ALA	3.1
2	F	114	ASP	3.0
2	F	28	GLY	2.9
2	F	56	SER	2.9
2	F	191	GLU	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	C	62	ALA	2.8
1	B	1002	GLY	2.7
2	C	191	GLU	2.7
2	F	51	LYS	2.6
2	F	9	LEU	2.6
2	C	58	ASP	2.6
2	C	7	ARG	2.5
2	C	55	SER	2.5
2	F	1	SER	2.5
2	C	60	ASN	2.5
2	C	57	SER	2.5
2	F	108	ARG	2.5
2	F	47	ALA	2.4
1	B	1017	GLN	2.4
2	C	3	GLU	2.4
2	F	169	ARG	2.4
2	C	170	ASN	2.3
2	C	171	VAL	2.3
2	F	2	ASP	2.3
2	C	117	GLU	2.2
2	F	61	GLU	2.2
2	F	117	GLU	2.2
1	A	988	LYS	2.2
2	C	107	GLN	2.2
2	C	65	LEU	2.2
1	A	1014	GLU	2.2
2	F	10	ILE	2.1
2	F	14	LYS	2.1
2	F	15	GLU	2.1
2	C	26	ARG	2.1
2	F	11	GLU	2.1
2	F	3	GLU	2.1
2	F	170	ASN	2.0
2	C	31	ARG	2.0
2	F	50	VAL	2.0
2	C	161	GLU	2.0
2	C	30	PRO	2.0
2	F	190	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	SO4	C	301	5/5	0.77	0.14	100,101,103,104	0
5	SO4	A	1204	5/5	0.78	0.17	94,96,97,100	0
5	SO4	F	301	5/5	0.81	0.12	75,79,80,81	0
5	SO4	B	1204	5/5	0.83	0.16	92,93,94,95	0
5	SO4	A	1203	5/5	0.95	0.12	48,49,51,51	0
4	TSV	B	1202	51/51	0.96	0.07	18,21,26,29	0
5	SO4	B	1203	5/5	0.97	0.08	49,49,50,51	0
4	TSV	A	1202	51/51	0.97	0.06	17,20,25,30	0
3	ZN	B	1201	1/1	0.98	0.04	41,41,41,41	0
3	ZN	A	1201	1/1	0.99	0.02	40,40,40,40	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around TSV B 1202:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around TSV A 1202:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.