



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 09:11 PM UTC

PDB ID : 1N54 / pdb_00001n54
Title : Cap Binding Complex m7GpppG free
Authors : Calero, G.; Wilson, K.; Ly, T.; Rios-Steiner, J.; Clardy, J.; Cerione, R.
Deposited on : 2002-11-04
Resolution : 2.72 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.72 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4348 (2.74-2.70)
Clashscore	190562	4665 (2.74-2.70)
Ramachandran outliers	187476	4584 (2.74-2.70)
Sidechain outliers	187428	4585 (2.74-2.70)
RSRZ outliers	180081	4348 (2.74-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	790	3% 44% 42% 7% 7%
2	B	156	3% 33% 38% • • 22%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	1556	-	X	-	-
3	GOL	A	1557	-	X	-	-
3	GOL	A	1558	-	X	-	-
3	GOL	A	1560	-	X	-	-
3	GOL	B	1555	-	X	-	-
3	GOL	B	1559	-	X	-	-
3	GOL	B	1561	-	X	-	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7292 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 80 kDa nuclear cap binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	733	5997	3865	1010	1084	38	0	0	0

- Molecule 2 is a protein called 20 kDa nuclear cap binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	121	998	628	171	193	6	0	0	0

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	A	1	6	3	3	0	0
3	A	1	6	3	3	0	0


Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	208	Total	O	0	0
			208	208		
4	B	47	Total	O	0	0
			47	47		



● Molecule 2: 20 kDa nuclear cap binding protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	111.77Å 111.77Å 175.73Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	24.41 – 2.72 24.41 – 2.72	Depositor EDS
% Data completeness (in resolution range)	72.9 (24.41-2.72) 89.6 (24.41-2.72)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.47 (at 2.58Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.263 , 0.282 0.258 , 0.281	Depositor DCC
R_{free} test set	2294 reflections (6.77%)	wwPDB-VP
Wilson B-factor (Å ²)	64.5	Xtriage
Anisotropy	0.264	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 55.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.045 for -h,-k,l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	7292	wwPDB-VP
Average B, all atoms (Å ²)	72.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.65	1/6149 (0.0%)	1.08	28/8343 (0.3%)
2	B	0.68	0/1015	1.08	7/1356 (0.5%)
All	All	0.65	1/7164 (0.0%)	1.08	35/9699 (0.4%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	303	MET	SD-CE	5.91	1.94	1.79

The worst 5 of 35 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	193	ASP	N-CA-C	-11.47	100.00	114.56
1	A	186	GLU	N-CA-C	-10.48	100.51	113.20
2	B	25	PHE	N-CA-C	9.41	124.76	108.02
2	B	17	LEU	N-CA-C	8.23	121.02	110.65
1	A	62	PRO	N-CA-C	-8.18	101.77	113.47

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5997	0	5978	380	1
2	B	998	0	970	69	0
3	A	24	0	19	3	0
3	B	18	0	13	3	0
4	A	208	0	0	50	1
4	B	47	0	0	5	0
All	All	7292	0	6980	445	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 32.

The worst 5 of 445 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:1561:GOL:C1	3:B:1561:GOL:O1	1.63	1.44
3:A:1556:GOL:O1	3:A:1556:GOL:C1	1.63	1.42
1:A:187:LYS:HE3	1:A:188:LYS:N	1.45	1.31
1:A:187:LYS:CE	1:A:188:LYS:H	1.53	1.22
1:A:513:THR:HG21	4:A:1743:HOH:O	1.37	1.20

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:187:LYS:CD	1:A:187:LYS:CD[5_675]	1.70	0.50
4:A:1599:HOH:O	4:A:1599:HOH:O[6_555]	1.77	0.43

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	727/790 (92%)	622 (86%)	87 (12%)	18 (2%)	4 10

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	119/156 (76%)	98 (82%)	17 (14%)	4 (3%)	3	6
All	All	846/946 (89%)	720 (85%)	104 (12%)	22 (3%)	4	10

5 of 22 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	66	SER
1	A	83	LEU
1	A	174	LEU
1	A	341	GLY
1	A	363	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	672/724 (93%)	630 (94%)	42 (6%)	16	37
2	B	106/130 (82%)	94 (89%)	12 (11%)	5	14
All	All	778/854 (91%)	724 (93%)	54 (7%)	14	33

5 of 54 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	449	VAL
1	A	635	LEU
2	B	31	GLU
1	A	455	LYS
1	A	503	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 28 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	425	GLN
2	B	46	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	561	HIS
1	A	748	GLN
1	A	463	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	B	1559	-	5,5,5	4.67	4 (80%)	5,5,5	5.75	3 (60%)
3	GOL	A	1556	-	5,5,5	4.68	5 (100%)	5,5,5	5.81	3 (60%)
3	GOL	B	1561	-	5,5,5	4.51	4 (80%)	5,5,5	6.23	3 (60%)
3	GOL	A	1557	-	5,5,5	4.71	5 (100%)	5,5,5	6.18	3 (60%)
3	GOL	A	1560	-	5,5,5	4.75	5 (100%)	5,5,5	6.29	3 (60%)
3	GOL	B	1555	-	5,5,5	4.81	4 (80%)	5,5,5	6.23	3 (60%)
3	GOL	A	1558	-	5,5,5	4.77	5 (100%)	5,5,5	6.48	3 (60%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	1559	-	-	3/4/4/4	-
3	GOL	A	1556	-	-	2/4/4/4	-
3	GOL	B	1561	-	-	2/4/4/4	-
3	GOL	A	1557	-	-	2/4/4/4	-
3	GOL	A	1560	-	-	2/4/4/4	-
3	GOL	B	1555	-	-	2/4/4/4	-
3	GOL	A	1558	-	-	2/4/4/4	-

The worst 5 of 32 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1560	GOL	C3-C2	-8.23	1.20	1.51
3	B	1555	GOL	C3-C2	-8.22	1.20	1.51
3	A	1557	GOL	C3-C2	-7.90	1.21	1.51
3	A	1558	GOL	C3-C2	-7.71	1.22	1.51
3	A	1556	GOL	C3-C2	-7.64	1.22	1.51

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1558	GOL	O3-C3-C2	12.07	164.70	110.38
3	A	1560	GOL	O3-C3-C2	11.34	161.42	110.38
3	A	1557	GOL	O3-C3-C2	11.20	160.81	110.38
3	B	1561	GOL	O3-C3-C2	11.02	159.99	110.38
3	B	1555	GOL	O3-C3-C2	10.87	159.31	110.38

There are no chirality outliers.

5 of 15 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1556	GOL	C1-C2-C3-O3
3	A	1557	GOL	O1-C1-C2-C3
3	A	1557	GOL	C1-C2-C3-O3
3	A	1558	GOL	C1-C2-C3-O3
3	A	1560	GOL	O1-C1-C2-C3

There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1556	GOL	3	0
3	B	1561	GOL	2	0
3	B	1555	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	733/790 (92%)	0.26	20 (2%) 56 53	20, 73, 102, 129	0
2	B	121/156 (77%)	0.25	5 (4%) 41 38	31, 74, 101, 114	0
All	All	854/946 (90%)	0.26	25 (2%) 53 50	20, 73, 102, 129	0

The worst 5 of 25 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	180	VAL	3.4
1	A	184	LEU	2.9
2	B	17	LEU	2.8
1	A	662	ALA	2.7
1	A	540	ASN	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GOL	A	1557	6/6	0.62	0.13	78,80,81,83	0
3	GOL	B	1561	6/6	0.66	0.11	58,59,59,61	0
3	GOL	A	1556	6/6	0.77	0.11	75,78,79,81	0
3	GOL	B	1559	6/6	0.79	0.10	74,76,76,77	0
3	GOL	A	1558	6/6	0.85	0.12	52,56,56,57	0
3	GOL	A	1560	6/6	0.87	0.18	66,68,70,72	0
3	GOL	B	1555	6/6	0.93	0.09	56,57,58,61	0

6.5 Other polymers [i](#)

There are no such residues in this entry.