



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 11:06 AM UTC

PDB ID : 1N67 / pdb_00001n67
Title : Clumping Factor A from Staphylococcus aureus
Authors : Deivanayagam, C.C.S.; Wann, E.R.; Chen, W.; Carson, M.; Rajashankar, K.R.; Hook, M.; Narayana, S.V.L.
Deposited on : 2002-11-08
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	359	5% 74% 17% 8%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2926 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Clumping Factor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	332	Total	C	N	O	S	0	0	0
			2547	1600	414	530	3			

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	202	MET	-	expression tag	UNP Q53653
A	203	ARG	-	expression tag	UNP Q53653
A	204	GLY	-	expression tag	UNP Q53653
A	205	SER	-	expression tag	UNP Q53653
A	206	HIS	-	expression tag	UNP Q53653
A	207	HIS	-	expression tag	UNP Q53653
A	208	HIS	-	expression tag	UNP Q53653
A	209	HIS	-	expression tag	UNP Q53653
A	210	HIS	-	expression tag	UNP Q53653
A	211	HIS	-	expression tag	UNP Q53653
A	212	GLY	-	expression tag	UNP Q53653
A	213	SER	-	expression tag	UNP Q53653
A	560	LYS	-	cloning artifact	UNP Q53653

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total	Mg	0	0
			3	3		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	376	Total	O	0	0
			376	376		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	38.70Å 80.54Å 113.89Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.28 – 1.90 24.28 – 1.90	Depositor EDS
% Data completeness (in resolution range)	96.8 (24.28-1.90) 98.8 (24.28-1.90)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.49 (at 1.90Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.206 , 0.256 0.212 , 0.260	Depositor DCC
R_{free} test set	2833 reflections (9.93%)	wwPDB-VP
Wilson B-factor (Å ²)	22.1	Xtriage
Anisotropy	0.518	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 36.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2926	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	0/2600	0.98	16/3559 (0.4%)

There are no bond length outliers.

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	235	GLN	OE1-CD-NE2	-9.79	112.81	122.60
1	A	253	GLN	OE1-CD-NE2	-9.79	112.81	122.60
1	A	302	GLN	OE1-CD-NE2	-9.72	112.88	122.60
1	A	416	LEU	N-CA-C	-7.42	99.50	110.48
1	A	541	LYS	N-CA-C	7.25	114.52	108.07
1	A	466	PHE	CA-C-N	7.24	126.94	119.56
1	A	466	PHE	C-N-CA	7.24	126.94	119.56
1	A	345	LYS	N-CA-C	6.81	121.44	113.20
1	A	253	GLN	CG-CD-NE2	6.66	126.39	116.40
1	A	235	GLN	CG-CD-NE2	6.64	126.36	116.40
1	A	302	GLN	CG-CD-NE2	6.60	126.30	116.40
1	A	546	GLU	N-CA-C	-5.72	101.57	110.10
1	A	384	ILE	N-CA-C	-5.68	97.52	109.34
1	A	513	ASN	N-CA-C	-5.35	101.51	109.96
1	A	376	TYR	CB-CA-C	-5.02	110.40	117.23
1	A	344	VAL	N-CA-C	-5.01	98.75	106.72

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2547	0	2467	54	0
2	A	3	0	0	0	0
3	A	376	0	0	5	0
All	All	2926	0	2467	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:495:ILE:HD11	1:A:503:LEU:HD13	1.53	0.89
1:A:389:LYS:HG3	1:A:531:ASN:HB2	1.61	0.80
1:A:437:LYS:HE2	3:A:1080:HOH:O	1.90	0.70
1:A:387:ILE:HD12	1:A:389:LYS:HE3	1.77	0.67
1:A:389:LYS:HE2	1:A:529:PHE:HA	1.77	0.66
1:A:535:SER:HB3	3:A:1075:HOH:O	1.96	0.64
1:A:389:LYS:HZ3	1:A:531:ASN:N	1.97	0.63
1:A:531:ASN:OD1	3:A:1205:HOH:O	2.15	0.62
1:A:389:LYS:HE2	1:A:529:PHE:CA	2.29	0.62
1:A:389:LYS:HE2	1:A:529:PHE:C	2.25	0.62
1:A:558:PRO:HG3	3:A:942:HOH:O	2.03	0.59
1:A:536:GLY:O	1:A:537:ASP:HB2	2.04	0.58
1:A:389:LYS:HZ3	1:A:531:ASN:HB2	1.69	0.58
1:A:381:LYS:HG3	1:A:448:TYR:CZ	2.39	0.57
1:A:389:LYS:HZ3	1:A:531:ASN:CA	2.16	0.57
1:A:421:ASP:O	1:A:501:GLY:HA3	2.03	0.57
1:A:495:ILE:CD1	1:A:503:LEU:HD13	2.32	0.56
1:A:541:LYS:HB2	1:A:542:PRO:HD2	1.86	0.56
1:A:516:ILE:HG23	1:A:516:ILE:O	2.06	0.55
1:A:559:GLU:O	1:A:560:LYS:HB3	2.06	0.54
1:A:414:GLY:HA3	1:A:472:TYR:CE2	2.42	0.54
1:A:530:ASN:H	1:A:530:ASN:HD22	1.55	0.54
1:A:526:GLU:HB3	1:A:539:ILE:HB	1.90	0.54
1:A:423:ASN:OD1	1:A:495:ILE:HD12	2.07	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:383:THR:O	1:A:396:GLN:HG3	2.08	0.53
1:A:345:LYS:HD3	3:A:1239:HOH:O	2.11	0.51
1:A:420:THR:HG22	1:A:421:ASP:N	2.28	0.49
1:A:423:ASN:CG	1:A:495:ILE:HG23	2.38	0.48
1:A:389:LYS:HZ3	1:A:531:ASN:CB	2.26	0.47
1:A:521:MET:HE3	1:A:522:SER:O	2.14	0.47
1:A:526:GLU:OE2	1:A:539:ILE:HD12	2.15	0.47
1:A:322:TYR:CZ	1:A:326:LYS:HG3	2.50	0.46
1:A:406:ASN:OD1	1:A:484:THR:HG22	2.15	0.46
1:A:546:GLU:HG3	1:A:560:LYS:HB3	1.97	0.45
1:A:550:GLU:HB2	1:A:553:GLU:HG3	1.98	0.45
1:A:279:VAL:HG21	1:A:285:LEU:HD11	1.96	0.45
1:A:381:LYS:HG3	1:A:448:TYR:OH	2.16	0.45
1:A:426:ILE:O	1:A:470:ASN:HA	2.16	0.45
1:A:550:GLU:CB	1:A:553:GLU:HG3	2.47	0.45
1:A:387:ILE:HD12	1:A:389:LYS:CE	2.45	0.45
1:A:286:ASN:HA	1:A:336:PRO:HG2	1.98	0.44
1:A:373:GLY:O	1:A:379:SER:HA	2.17	0.44
1:A:439:ASP:OD2	1:A:454:ASN:HA	2.17	0.44
1:A:503:LEU:O	1:A:526:GLU:HA	2.18	0.43
1:A:530:ASN:C	1:A:532:GLY:H	2.26	0.43
1:A:530:ASN:O	1:A:531:ASN:HB3	2.19	0.43
1:A:245:SER:O	1:A:363:LYS:NZ	2.52	0.42
1:A:389:LYS:NZ	1:A:531:ASN:N	2.66	0.42
1:A:322:TYR:CE2	1:A:326:LYS:HG3	2.55	0.42
1:A:416:LEU:CD2	1:A:424:ALA:HB3	2.49	0.42
1:A:515:ASN:O	1:A:516:ILE:HG22	2.19	0.42
1:A:424:ALA:HA	1:A:495:ILE:HD13	2.02	0.41
1:A:235:GLN:O	1:A:266:PRO:CG	2.69	0.41
1:A:455:PHE:CD1	1:A:455:PHE:C	2.99	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	330/359 (92%)	316 (96%)	13 (4%)	1 (0%)	36	29

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	286	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	292/311 (94%)	288 (99%)	4 (1%)	59	59

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	405	ASP
1	A	481	ASP
1	A	530	ASN
1	A	541	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	253	GLN
1	A	284	ASN
1	A	419	ASN
1	A	460	ASN
1	A	468	ASN
1	A	471	GLN
1	A	477	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	332/359 (92%)	0.38	17 (5%) 33 36	17, 24, 37, 60	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	532	GLY	7.3
1	A	535	SER	4.3
1	A	530	ASN	4.2
1	A	536	GLY	4.1
1	A	531	ASN	3.7
1	A	534	GLY	3.6
1	A	533	SER	3.6
1	A	529	PHE	3.5
1	A	419	ASN	3.0
1	A	516	ILE	3.0
1	A	453	GLU	2.9
1	A	229	GLY	2.5
1	A	500	LYS	2.5
1	A	560	LYS	2.4
1	A	495	ILE	2.3
1	A	346	LYS	2.2
1	A	389	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MG	A	901	1/1	0.95	0.29	32,32,32,32	0
2	MG	A	902	1/1	0.96	0.08	36,36,36,36	0
2	MG	A	900	1/1	0.99	0.02	29,29,29,29	0

6.5 Other polymers [i](#)

There are no such residues in this entry.