



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 02:06 AM UTC

PDB ID : 4N6G / pdb_00004n6g
Title : Tailoring Small Molecules for an Allosteric Site on Procaspase-6
Authors : Murray, J.M.; Steffek, M.
Deposited on : 2013-10-11
Resolution : 2.14 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.14 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3689 (2.16-2.12)
Clashscore	190562	3812 (2.16-2.12)
Ramachandran outliers	187476	3773 (2.16-2.12)
Sidechain outliers	187428	3772 (2.16-2.12)
RSRZ outliers	180081	3691 (2.16-2.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	282	
1	B	282	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PO4	B	403	-	X	-	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4224 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Caspase-6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	257	Total	C	N	O	S	0	0	0
			2042	1304	355	370	13			
1	B	243	Total	C	N	O	S	0	0	0
			1942	1241	336	352	13			

There are 26 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	21	MET	-	initiating methionine	UNP P55212
A	22	GLY	-	expression tag	UNP P55212
A	23	SER	-	expression tag	UNP P55212
A	163	ALA	CYS	engineered mutation	UNP P55212
A	294	GLY	-	expression tag	UNP P55212
A	295	ASN	-	expression tag	UNP P55212
A	296	SER	-	expression tag	UNP P55212
A	297	HIS	-	expression tag	UNP P55212
A	298	HIS	-	expression tag	UNP P55212
A	299	HIS	-	expression tag	UNP P55212
A	300	HIS	-	expression tag	UNP P55212
A	301	HIS	-	expression tag	UNP P55212
A	302	HIS	-	expression tag	UNP P55212
B	21	MET	-	initiating methionine	UNP P55212
B	22	GLY	-	expression tag	UNP P55212
B	23	SER	-	expression tag	UNP P55212
B	163	ALA	CYS	engineered mutation	UNP P55212
B	294	GLY	-	expression tag	UNP P55212
B	295	ASN	-	expression tag	UNP P55212
B	296	SER	-	expression tag	UNP P55212
B	297	HIS	-	expression tag	UNP P55212
B	298	HIS	-	expression tag	UNP P55212
B	299	HIS	-	expression tag	UNP P55212
B	300	HIS	-	expression tag	UNP P55212
B	301	HIS	-	expression tag	UNP P55212

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	302	HIS	-	expression tag	UNP P55212

- Molecule 2 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is 3-(pyrrolidin-1-yl)isoquinolin-1(2H)-one (CCD ID: 2GQ) (formula: $C_{13}H_{14}N_2O$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	N	O	0	0
			12	9	2	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	112	Total	O	0	0
			112	112		
4	B	96	Total	O	0	0
			96	96		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Caspase-6



• Molecule 1: Caspase-6



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	116.08Å 116.08Å 79.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	82.08 – 2.14 82.08 – 2.14	Depositor EDS
% Data completeness (in resolution range)	99.9 (82.08-2.14) 100.0 (82.08-2.14)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.18 (at 2.14Å)	Xtriage
Refinement program	PHENIX 1.8.4_1492	Depositor
R, R_{free}	0.179 , 0.217 0.183 , 0.220	Depositor DCC
R_{free} test set	1570 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å ²)	32.3	Xtriage
Anisotropy	0.131	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 60.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4224	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.93% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 2GQ, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.85	4/2091 (0.2%)	1.03	11/2821 (0.4%)
1	B	0.71	0/1987	0.87	3/2677 (0.1%)
All	All	0.78	4/4078 (0.1%)	0.95	14/5498 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	264	CYS	CA-C	-7.25	1.43	1.52
1	A	29	GLU	CA-C	5.86	1.60	1.52
1	A	265	LYS	CA-C	-5.70	1.45	1.52
1	A	60	THR	CA-CB	5.47	1.61	1.53

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	264	CYS	N-CA-C	-8.80	94.24	108.41
1	A	266	ASP	N-CA-C	8.25	120.71	109.24
1	A	263	PHE	CA-C-N	8.16	133.27	122.42
1	A	263	PHE	C-N-CA	8.16	133.27	122.42
1	A	263	PHE	O-C-N	7.37	129.69	122.03
1	A	267	PRO	CA-N-CD	-6.96	102.25	112.00
1	A	147	LYS	N-CA-C	-6.00	104.86	111.82

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	267	PRO	N-CA-C	5.72	124.25	112.47
1	B	32	ASP	CA-C-N	5.59	125.30	119.82
1	B	32	ASP	C-N-CA	5.59	125.30	119.82
1	A	266	ASP	CA-C-O	5.20	123.17	119.32
1	A	29	GLU	N-CA-C	5.12	121.70	110.80
1	A	266	ASP	O-C-N	5.04	125.99	121.35
1	B	245	PHE	N-CA-C	5.03	116.77	111.28

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	264	CYS	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2042	0	1979	19	0
1	B	1942	0	1902	26	0
2	A	10	0	0	0	0
2	B	10	0	0	1	0
3	B	12	0	6	2	0
4	A	112	0	0	2	0
4	B	96	0	0	2	0
All	All	4224	0	3887	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:172:VAL:O	1:A:174:PRO:HD3	1.88	0.74
1:B:201:PRO:HB2	1:B:281:MET:SD	2.40	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:263:PHE:O	1:A:265:LYS:O	2.19	0.61
1:A:201:PRO:HB2	1:A:281:MET:SD	2.41	0.60
1:A:202:ALA:HB3	1:A:208:MET:HE3	1.82	0.60
1:A:255:LYS:NZ	4:A:545:HOH:O	2.36	0.59
1:B:71:ARG:NH2	4:B:589:HOH:O	2.24	0.58
1:A:200:LEU:HD11	3:B:401:2GQ:H2	1.87	0.57
1:B:118:PHE:HB3	1:B:127:ILE:CD1	2.35	0.56
1:A:158:PHE:CD1	1:A:208:MET:HE1	2.42	0.55
1:A:64:ARG:HA	1:A:220:ARG:NH2	2.22	0.54
1:B:202:ALA:HB3	1:B:208:MET:HE3	1.89	0.54
1:B:60:THR:O	1:B:167:GLN:NE2	2.41	0.53
1:B:135:GLU:HB2	1:B:138:THR:HG23	1.89	0.53
1:A:44:ARG:HD2	1:A:81:LEU:O	2.08	0.53
1:B:158:PHE:CD1	1:B:208:MET:HE1	2.46	0.50
1:B:145:GLY:N	2:B:403:PO4:O1	2.43	0.49
1:B:206:PHE:HD2	1:B:208:MET:CE	2.26	0.49
1:A:239:TYR:CZ	1:B:33:PRO:HG3	2.49	0.48
1:B:227:TRP:CD2	1:B:259:ARG:HD3	2.48	0.48
1:B:268:SER:O	1:B:272:LYS:HE2	2.13	0.48
1:A:46:ILE:HG22	1:A:48:LEU:HD12	1.95	0.48
1:A:199:THR:OG1	1:B:201:PRO:HD3	2.14	0.48
1:B:46:ILE:HG22	1:B:48:LEU:HD12	1.96	0.47
1:B:48:LEU:HD13	1:B:114:PHE:CE1	2.50	0.46
1:B:200:LEU:HD11	3:B:401:2GQ:O	2.15	0.46
1:B:220:ARG:HA	1:B:226:SER:HA	1.98	0.46
1:B:64:ARG:HA	1:B:220:ARG:NH2	2.31	0.45
1:A:206:PHE:HD2	1:A:208:MET:CE	2.29	0.45
1:B:191:GLU:OE1	1:B:220:ARG:NH2	2.51	0.44
1:A:94:GLU:HG3	4:A:547:HOH:O	2.19	0.43
1:A:158:PHE:HB2	1:A:208:MET:SD	2.58	0.43
1:A:135:GLU:HB2	1:A:138:THR:HG23	2.01	0.42
1:B:101:HIS:O	1:B:105:THR:HG23	2.19	0.42
1:B:206:PHE:HB3	1:B:208:MET:HE3	2.01	0.42
1:A:93:ALA:O	1:A:97:LEU:HD23	2.20	0.42
1:B:126:HIS:HB3	1:B:133:LYS:HB2	2.02	0.42
1:B:71:ARG:NE	4:B:589:HOH:O	2.43	0.41
1:B:151:LEU:HD23	1:B:151:LEU:HA	1.93	0.41
1:A:201:PRO:HD3	1:B:199:THR:OG1	2.20	0.41
1:A:206:PHE:HB3	1:A:208:MET:HE3	2.02	0.41
1:B:120:SER:O	1:B:162:ALA:HA	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	253/282 (90%)	237 (94%)	14 (6%)	2 (1%)	16	9
1	B	239/282 (85%)	229 (96%)	9 (4%)	1 (0%)	30	26
All	All	492/564 (87%)	466 (95%)	23 (5%)	3 (1%)	21	15

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	265	LYS
1	B	145	GLY
1	A	267	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	216/247 (87%)	206 (95%)	10 (5%)	24	20
1	B	208/247 (84%)	203 (98%)	5 (2%)	43	45
All	All	424/494 (86%)	409 (96%)	15 (4%)	32	31

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	27	LYS
1	A	28	ARG
1	A	29	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	60	THR
1	A	77	ARG
1	A	96	LEU
1	A	139	LEU
1	A	170	VAL
1	A	251	LEU
1	A	268	SER
1	B	60	THR
1	B	139	LEU
1	B	249	LEU
1	B	251	LEU
1	B	268	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	161	GLN
1	A	166	ASN
1	A	167	GLN
1	A	168	HIS
1	A	230	GLN
1	B	101	HIS
1	B	126	HIS
1	B	161	GLN
1	B	230	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PO4	A	402	-	4,4,4	1.92	2 (50%)	6,6,6	1.96	1 (16%)
2	PO4	B	403	-	4,4,4	2.28	2 (50%)	6,6,6	4.62	6 (100%)
2	PO4	A	401	-	4,4,4	0.62	0	6,6,6	0.89	0
3	2GQ	B	401	-	13,13,18	0.32	0	16,18,25	0.32	0
2	PO4	B	402	-	4,4,4	0.64	0	6,6,6	1.09	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	2GQ	B	401	-	-	-	0/2/2/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	403	PO4	P-O1	3.79	1.59	1.50
2	A	402	PO4	P-O2	2.67	1.62	1.54
2	B	403	PO4	P-O2	2.33	1.61	1.54
2	A	402	PO4	P-O1	2.30	1.56	1.50

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	403	PO4	O4-P-O2	6.48	128.07	107.91
2	B	403	PO4	O2-P-O1	-5.64	91.02	110.95
2	B	403	PO4	O4-P-O3	-4.37	94.32	107.91
2	B	403	PO4	O3-P-O2	4.09	120.65	107.91
2	A	402	PO4	O4-P-O2	3.53	118.89	107.91

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	403	PO4	O3-P-O1	-3.10	100.00	110.95
2	B	403	PO4	O4-P-O1	2.97	121.46	110.95

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	403	PO4	1	0
3	B	401	2GQ	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	257/282 (91%)	0.19	27 (10%) 11 13	21, 35, 74, 88	0
1	B	243/282 (86%)	-0.00	8 (3%) 49 53	20, 37, 65, 78	0
All	All	500/564 (88%)	0.10	35 (7%) 22 26	20, 36, 69, 88	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	30	MET	5.3
1	A	174	PRO	4.8
1	A	264	CYS	4.6
1	B	187	THR	4.3
1	A	25	PHE	4.0
1	A	170	VAL	4.0
1	A	26	TYR	3.6
1	A	29	GLU	3.5
1	A	168	HIS	3.2
1	A	31	PHE	3.1
1	B	30	MET	3.0
1	A	171	PRO	3.0
1	A	262	ASP	2.9
1	B	31	PHE	2.9
1	A	173	ILE	2.8
1	A	189	ILE	2.8
1	A	169	ASP	2.8
1	A	58	HIS	2.7
1	A	28	ARG	2.7
1	A	267	PRO	2.6
1	B	40	ASP	2.5
1	B	263	PHE	2.5
1	A	27	LYS	2.5
1	A	54	ARG	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	265	LYS	2.4
1	B	32	ASP	2.3
1	A	266	ASP	2.2
1	A	292	SER	2.2
1	A	166	ASN	2.2
1	A	40	ASP	2.2
1	A	186	ASP	2.1
1	A	270	ILE	2.1
1	B	34	ALA	2.1
1	A	125	ASN	2.0
1	B	146	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	2GQ	B	401	12/16	0.65	0.18	61,67,73,74	0
2	PO4	B	403	5/5	0.67	0.22	57,76,84,88	0
2	PO4	A	402	5/5	0.90	0.10	41,51,67,116	0
2	PO4	B	402	5/5	0.98	0.05	28,29,30,31	0
2	PO4	A	401	5/5	0.99	0.05	22,28,29,30	0

6.5 Other polymers [i](#)

There are no such residues in this entry.