



## wwPDB EM Validation Summary Report ⓘ

Mar 10, 2026 – 07:03 AM UTC

PDB ID : 5N60 / pdb\_00005n60  
EMDB ID : EMD-3592  
Title : Cryo-EM structure of RNA polymerase I in complex with Rrn3 and Core Factor (Orientation I)  
Authors : Engel, C.; Gubbey, T.; Neyer, S.; Sainsbury, S.; Oberthuer, C.; Baejen, C.; Bernecky, C.; Cramer, P.  
Deposited on : 2017-02-14  
Resolution : 7.70 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev132  
MolProbity : 4-5-2 with Phenix2.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49

# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:  
*ELECTRON MICROSCOPY*

The reported resolution of this entry is 7.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric                | Whole archive<br>(#Entries) | EM structures<br>(#Entries) | Similar EM resolution<br>(#Entries, resolution range(Å)) |
|-----------------------|-----------------------------|-----------------------------|----------------------------------------------------------|
| Clashscore            | 229148                      | 23984                       | -                                                        |
| Ramachandran outliers | 224038                      | 23583                       | -                                                        |
| Sidechain outliers    | 223484                      | 23102                       | -                                                        |
| Q-score               | -                           | 25397                       | 384 ( 7.20 - 8.20 )                                      |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|------------------|
| 1   | A     | 1664   |                  |
| 2   | B     | 1203   |                  |
| 3   | C     | 335    |                  |
| 4   | D     | 137    |                  |

Continued on next page...

Continued from previous page...

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|------------------|
| 5   | E     | 215    |                  |
| 6   | F     | 155    |                  |
| 7   | G     | 326    |                  |
| 8   | H     | 146    |                  |
| 9   | I     | 125    |                  |
| 10  | J     | 70     |                  |
| 11  | K     | 142    |                  |
| 12  | L     | 70     |                  |
| 13  | M     | 415    |                  |
| 14  | N     | 233    |                  |
| 15  | O     | 627    |                  |
| 16  | P     | 894    |                  |
| 17  | Q     | 514    |                  |
| 18  | R     | 507    |                  |

## 2 Entry composition

There are 19 unique types of molecules in this entry. The entry contains 48004 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase I subunit RPA190.

| Mol | Chain | Residues | Atoms |      |      |      |    | AltConf | Trace |
|-----|-------|----------|-------|------|------|------|----|---------|-------|
| 1   | A     | 1467     | Total | C    | N    | O    | S  | 0       | 0     |
|     |       |          | 11598 | 7327 | 2017 | 2193 | 61 |         |       |

- Molecule 2 is a protein called DNA-directed RNA polymerase I subunit RPA135.

| Mol | Chain | Residues | Atoms |      |      |      |    | AltConf | Trace |
|-----|-------|----------|-------|------|------|------|----|---------|-------|
| 2   | B     | 1172     | Total | C    | N    | O    | S  | 0       | 0     |
|     |       |          | 9312  | 5891 | 1633 | 1738 | 50 |         |       |

- Molecule 3 is a protein called DNA-directed RNA polymerases I and III subunit RPAC1.

| Mol | Chain | Residues | Atoms |      |     |     |   | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|---------|-------|
| 3   | C     | 305      | Total | C    | N   | O   | S | 0       | 0     |
|     |       |          | 2423  | 1539 | 416 | 460 | 8 |         |       |

- Molecule 4 is a protein called DNA-directed RNA polymerase I subunit RPA14.

| Mol | Chain | Residues | Atoms |     |    |    | AltConf | Trace |
|-----|-------|----------|-------|-----|----|----|---------|-------|
| 4   | D     | 58       | Total | C   | N  | O  | 0       | 0     |
|     |       |          | 459   | 289 | 78 | 92 |         |       |

- Molecule 5 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC1.

| Mol | Chain | Residues | Atoms |      |     |     |    | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| 5   | E     | 212      | Total | C    | N   | O   | S  | 0       | 0     |
|     |       |          | 1735  | 1102 | 306 | 316 | 11 |         |       |

- Molecule 6 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC2.

| Mol | Chain | Residues | Atoms |     |     |     |   | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 6   | F     | 100      | Total | C   | N   | O   | S | 0       | 0     |
|     |       |          | 823   | 522 | 144 | 154 | 3 |         |       |

- Molecule 7 is a protein called DNA-directed RNA polymerase I subunit RPA43.

| Mol | Chain | Residues | Atoms |     |     |     |   | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 7   | G     | 193      | Total | C   | N   | O   | S | 0       | 0     |
|     |       |          | 1526  | 985 | 262 | 274 | 5 |         |       |

- Molecule 8 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC3.

| Mol | Chain | Residues | Atoms |     |     |     |   | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 8   | H     | 131      | Total | C   | N   | O   | S | 0       | 0     |
|     |       |          | 1052  | 664 | 176 | 208 | 4 |         |       |

- Molecule 9 is a protein called DNA-directed RNA polymerase I subunit RPA12.

| Mol | Chain | Residues | Atoms |     |     |     |   | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 9   | I     | 99       | Total | C   | N   | O   | S | 0       | 0     |
|     |       |          | 755   | 472 | 125 | 149 | 9 |         |       |

- Molecule 10 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC5.

| Mol | Chain | Residues | Atoms |     |     |     |   | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 10  | J     | 69       | Total | C   | N   | O   | S | 0       | 0     |
|     |       |          | 569   | 362 | 101 | 100 | 6 |         |       |

- Molecule 11 is a protein called DNA-directed RNA polymerases I and III subunit RPAC2.

| Mol | Chain | Residues | Atoms |     |     |     |   | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 11  | K     | 101      | Total | C   | N   | O   | S | 0       | 0     |
|     |       |          | 793   | 496 | 130 | 162 | 5 |         |       |

- Molecule 12 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC4.

| Mol | Chain | Residues | Atoms |     |    |    |   | AltConf | Trace |
|-----|-------|----------|-------|-----|----|----|---|---------|-------|
| 12  | L     | 44       | Total | C   | N  | O  | S | 0       | 0     |
|     |       |          | 352   | 217 | 70 | 61 | 4 |         |       |

- Molecule 13 is a protein called DNA-directed RNA polymerase I subunit RPA49.

| Mol | Chain | Residues | Atoms |     |     |     |  | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|--|---------|-------|
| 13  | M     | 108      | Total | C   | N   | O   |  | 0       | 0     |
|     |       |          | 856   | 543 | 142 | 171 |  |         |       |

- Molecule 14 is a protein called DNA-directed RNA polymerase I subunit RPA34.

| Mol | Chain | Residues | Atoms |     |     |     |   | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 14  | N     | 145      | Total | C   | N   | O   | S | 0       | 0     |
|     |       |          | 1151  | 735 | 188 | 224 | 4 |         |       |

- Molecule 15 is a protein called RNA polymerase I-specific transcription initiation factor RRN3.

| Mol | Chain | Residues | Atoms |      |     |     |    | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| 15  | O     | 473      | Total | C    | N   | O   | S  | 0       | 0     |
|     |       |          | 3907  | 2533 | 642 | 711 | 21 |         |       |

- Molecule 16 is a protein called RNA polymerase I-specific transcription initiation factor RRN6.

| Mol | Chain | Residues | Atoms |      |     |     |    | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| 16  | P     | 583      | Total | C    | N   | O   | S  | 0       | 0     |
|     |       |          | 4729  | 3010 | 803 | 905 | 11 |         |       |

- Molecule 17 is a protein called RNA polymerase I-specific transcription initiation factor RRN7.

| Mol | Chain | Residues | Atoms |      |     |     |    | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| 17  | Q     | 410      | Total | C    | N   | O   | S  | 0       | 0     |
|     |       |          | 3421  | 2219 | 579 | 603 | 20 |         |       |

- Molecule 18 is a protein called RNA polymerase I-specific transcription initiation factor RRN11.

| Mol | Chain | Residues | Atoms |      |     |     |    | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| 18  | R     | 303      | Total | C    | N   | O   | S  | 0       | 0     |
|     |       |          | 2535  | 1634 | 456 | 434 | 11 |         |       |

- Molecule 19 is ZINC ION (CCD ID: ZN) (formula: Zn).

| Mol | Chain | Residues | Atoms |    | AltConf |
|-----|-------|----------|-------|----|---------|
| 19  | A     | 2        | Total | Zn | 0       |
|     |       |          | 2     | 2  |         |
| 19  | B     | 1        | Total | Zn | 0       |
|     |       |          | 1     | 1  |         |
| 19  | I     | 2        | Total | Zn | 0       |
|     |       |          | 2     | 2  |         |
| 19  | J     | 1        | Total | Zn | 0       |
|     |       |          | 1     | 1  |         |

*Continued on next page...*


*Continued from previous page...*


| Mol | Chain | Residues | Atoms      |         | AltConf |
|-----|-------|----------|------------|---------|---------|
| 19  | L     | 1        | Total<br>1 | Zn<br>1 | 0       |
| 19  | Q     | 1        | Total<br>1 | Zn<br>1 | 0       |

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA-directed RNA polymerase I subunit RPA190







• Molecule 2: DNA-directed RNA polymerase I subunit RPA135










• Molecule 3: DNA-directed RNA polymerases I and III subunit RPAC1



• Molecule 4: DNA-directed RNA polymerase I subunit RPA14



• Molecule 5: DNA-directed RNA polymerases I, II, and III subunit RPABC1




• Molecule 6: DNA-directed RNA polymerases I, II, and III subunit RPABC2









- Molecule 10: DNA-directed RNA polymerases I, II, and III subunit RPABC5




- Molecule 11: DNA-directed RNA polymerases I and III subunit RPAC2



- Molecule 12: DNA-directed RNA polymerases I, II, and III subunit RPABC4




- Molecule 13: DNA-directed RNA polymerase I subunit RPA49





GLY  
ILE  
PRO  
LYS  
LYS  
THR  
THR  
ALA  
ALA  
SER  
TYR  
LYS  
ASP  
ILE  
ALA  
THR  
MET  
ASP  
LYS  
VAL  
VAL  
PRO  
PHE  
PHE  
LYS  
LEU  
PRO  
GLU  
PHE  
MET  
THR  
THR  
ARG  
ARG  
GLY  
ARG  
GLY  
PRO  
PRO  
ARG

• Molecule 14: DNA-directed RNA polymerase I subunit RPA34




• Molecule 15: RNA polymerase I-specific transcription initiation factor RRN3






• Molecule 16: RNA polymerase I-specific transcription initiation factor RRN6







## 4 Experimental information

| Property                             | Value                                   | Source    |
|--------------------------------------|-----------------------------------------|-----------|
| EM reconstruction method             | SINGLE PARTICLE                         | Depositor |
| Imposed symmetry                     | POINT, Not provided                     |           |
| Number of particles used             | 8317                                    | Depositor |
| Resolution determination method      | FSC 0.143 CUT-OFF                       | Depositor |
| CTF correction method                | PHASE FLIPPING AND AMPLITUDE CORRECTION | Depositor |
| Microscope                           | FEI TITAN KRIOS                         | Depositor |
| Voltage (kV)                         | 300                                     | Depositor |
| Electron dose ( $e^-/\text{\AA}^2$ ) | 40                                      | Depositor |
| Minimum defocus (nm)                 | Not provided                            |           |
| Maximum defocus (nm)                 | Not provided                            |           |
| Magnification                        | Not provided                            |           |
| Image detector                       | GATAN K2 SUMMIT (4k x 4k)               | Depositor |
| Maximum map value                    | 0.081                                   | Depositor |
| Minimum map value                    | -0.047                                  | Depositor |
| Average map value                    | 0.000                                   | Depositor |
| Map value standard deviation         | 0.006                                   | Depositor |
| Recommended contour level            | 0.023                                   | Depositor |
| Map size (Å)                         | 324.0, 324.0, 324.0                     | wwPDB     |
| Map dimensions                       | 240, 240, 240                           | wwPDB     |
| Map angles (°)                       | 90.0, 90.0, 90.0                        | wwPDB     |
| Pixel spacing (Å)                    | 1.35, 1.35, 1.35                        | Depositor |

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths |                | Bond angles |                  |
|-----|-------|--------------|----------------|-------------|------------------|
|     |       | RMSZ         | # $ Z  > 5$    | RMSZ        | # $ Z  > 5$      |
| 1   | A     | 0.64         | 1/11809 (0.0%) | 0.96        | 37/15943 (0.2%)  |
| 2   | B     | 0.62         | 0/9518         | 0.98        | 36/12863 (0.3%)  |
| 3   | C     | 0.53         | 0/2475         | 0.94        | 8/3354 (0.2%)    |
| 4   | D     | 0.51         | 0/465          | 0.81        | 0/630            |
| 5   | E     | 0.49         | 0/1771         | 0.88        | 3/2383 (0.1%)    |
| 6   | F     | 0.58         | 0/838          | 0.86        | 0/1129           |
| 7   | G     | 0.51         | 0/1563         | 0.84        | 3/2124 (0.1%)    |
| 8   | H     | 0.52         | 0/1070         | 0.88        | 0/1449           |
| 9   | I     | 0.53         | 0/765          | 0.87        | 0/1030           |
| 10  | J     | 0.59         | 0/578          | 0.98        | 6/775 (0.8%)     |
| 11  | K     | 0.54         | 0/804          | 0.93        | 6/1083 (0.6%)    |
| 12  | L     | 0.50         | 0/354          | 0.91        | 2/468 (0.4%)     |
| 13  | M     | 0.49         | 0/872          | 0.87        | 0/1170           |
| 14  | N     | 0.48         | 0/1172         | 0.88        | 1/1580 (0.1%)    |
| 15  | O     | 0.45         | 1/3996 (0.0%)  | 0.82        | 2/5401 (0.0%)    |
| 16  | P     | 0.38         | 0/4822         | 0.74        | 6/6525 (0.1%)    |
| 17  | Q     | 0.36         | 1/3502 (0.0%)  | 0.67        | 2/4727 (0.0%)    |
| 18  | R     | 0.34         | 0/2591         | 0.78        | 6/3483 (0.2%)    |
| All | All   | 0.54         | 3/48965 (0.0%) | 0.89        | 118/66117 (0.2%) |

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

| Mol | Chain | #Chirality outliers | #Planarity outliers |
|-----|-------|---------------------|---------------------|
| 1   | A     | 0                   | 15                  |
| 2   | B     | 0                   | 4                   |
| 3   | C     | 0                   | 1                   |
| 6   | F     | 0                   | 1                   |
| 12  | L     | 0                   | 2                   |

*Continued on next page...*

*Continued from previous page...*

| Mol | Chain | #Chirality outliers | #Planarity outliers |
|-----|-------|---------------------|---------------------|
| 13  | M     | 0                   | 3                   |
| 14  | N     | 0                   | 1                   |
| 15  | O     | 0                   | 4                   |
| 16  | P     | 0                   | 26                  |
| 17  | Q     | 0                   | 8                   |
| 18  | R     | 0                   | 8                   |
| All | All   | 0                   | 73                  |

All (3) bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z    | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|-------|------|-------------|----------|
| 1   | A     | 356 | PHE  | C-N   | 8.46 | 1.44        | 1.33     |
| 17  | Q     | 15  | CYS  | CA-C  | 7.52 | 1.62        | 1.52     |
| 15  | O     | 212 | THR  | CA-CB | 5.44 | 1.62        | 1.53     |

The worst 5 of 118 bond angle outliers are listed below:

| Mol | Chain | Res  | Type | Atoms  | Z      | Observed(°) | Ideal(°) |
|-----|-------|------|------|--------|--------|-------------|----------|
| 18  | R     | 205  | VAL  | N-CA-C | 18.78  | 128.18      | 110.53   |
| 2   | B     | 405  | GLY  | N-CA-C | -12.26 | 97.18       | 113.24   |
| 1   | A     | 1071 | ASP  | N-CA-C | -9.90  | 97.47       | 110.53   |
| 3   | C     | 223  | SER  | N-CA-C | -9.68  | 101.63      | 112.57   |
| 15  | O     | 376  | TYR  | N-CA-C | -9.46  | 100.42      | 113.56   |

There are no chirality outliers.

5 of 73 planarity outliers are listed below:

| Mol | Chain | Res | Type | Group   |
|-----|-------|-----|------|---------|
| 1   | A     | 405 | LYS  | Peptide |
| 1   | A     | 406 | LEU  | Peptide |
| 1   | A     | 407 | GLN  | Peptide |
| 1   | A     | 410 | LYS  | Peptide |
| 1   | A     | 411 | VAL  | Peptide |

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1   | A     | 11598 | 0        | 11666    | 1160    | 0            |
| 2   | B     | 9312  | 0        | 9200     | 1000    | 0            |
| 3   | C     | 2423  | 0        | 2412     | 144     | 0            |
| 4   | D     | 459   | 0        | 462      | 7       | 0            |
| 5   | E     | 1735  | 0        | 1764     | 32      | 0            |
| 6   | F     | 823   | 0        | 839      | 91      | 0            |
| 7   | G     | 1526  | 0        | 1534     | 151     | 0            |
| 8   | H     | 1052  | 0        | 1021     | 56      | 0            |
| 9   | I     | 755   | 0        | 728      | 105     | 0            |
| 10  | J     | 569   | 0        | 584      | 54      | 0            |
| 11  | K     | 793   | 0        | 790      | 57      | 0            |
| 12  | L     | 352   | 0        | 371      | 59      | 0            |
| 13  | M     | 856   | 0        | 852      | 60      | 0            |
| 14  | N     | 1151  | 0        | 1168     | 172     | 0            |
| 15  | O     | 3907  | 0        | 3902     | 418     | 0            |
| 16  | P     | 4729  | 0        | 4675     | 626     | 0            |
| 17  | Q     | 3421  | 0        | 3463     | 533     | 0            |
| 18  | R     | 2535  | 0        | 2600     | 741     | 0            |
| 19  | A     | 2     | 0        | 0        | 1       | 0            |
| 19  | B     | 1     | 0        | 0        | 0       | 0            |
| 19  | I     | 2     | 0        | 0        | 0       | 0            |
| 19  | J     | 1     | 0        | 0        | 0       | 0            |
| 19  | L     | 1     | 0        | 0        | 0       | 0            |
| 19  | Q     | 1     | 0        | 0        | 0       | 0            |
| All | All   | 48004 | 0        | 48031    | 3997    | 0            |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 42.

The worst 5 of 3997 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1          | Atom-2           | Interatomic distance (Å) | Clash overlap (Å) |
|-----------------|------------------|--------------------------|-------------------|
| 17:Q:20:TRP:CZ3 | 17:Q:22:ILE:CG2  | 1.76                     | 1.69              |
| 17:Q:381:MET:SD | 18:R:212:HIS:HB3 | 1.26                     | 1.68              |
| 3:C:272:LYS:HA  | 14:N:175:TYR:CE1 | 1.16                     | 1.63              |
| 1:A:995:TYR:CD2 | 2:B:708:ASP:HA   | 1.29                     | 1.62              |
| 1:A:545:SER:HB2 | 17:Q:34:VAL:CG2  | 1.13                     | 1.61              |

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed        | Favoured   | Allowed  | Outliers | Percentiles |     |
|-----|-------|-----------------|------------|----------|----------|-------------|-----|
| 1   | A     | 1449/1664 (87%) | 1375 (95%) | 60 (4%)  | 14 (1%)  | 12          | 48  |
| 2   | B     | 1164/1203 (97%) | 1095 (94%) | 51 (4%)  | 18 (2%)  | 8           | 40  |
| 3   | C     | 303/335 (90%)   | 287 (95%)  | 13 (4%)  | 3 (1%)   | 12          | 48  |
| 4   | D     | 54/137 (39%)    | 50 (93%)   | 2 (4%)   | 2 (4%)   | 2           | 20  |
| 5   | E     | 210/215 (98%)   | 197 (94%)  | 11 (5%)  | 2 (1%)   | 12          | 48  |
| 6   | F     | 98/155 (63%)    | 95 (97%)   | 2 (2%)   | 1 (1%)   | 12          | 48  |
| 7   | G     | 187/326 (57%)   | 173 (92%)  | 12 (6%)  | 2 (1%)   | 11          | 46  |
| 8   | H     | 127/146 (87%)   | 121 (95%)  | 6 (5%)   | 0        | 100         | 100 |
| 9   | I     | 91/125 (73%)    | 80 (88%)   | 8 (9%)   | 3 (3%)   | 3           | 21  |
| 10  | J     | 67/70 (96%)     | 63 (94%)   | 4 (6%)   | 0        | 100         | 100 |
| 11  | K     | 99/142 (70%)    | 92 (93%)   | 7 (7%)   | 0        | 100         | 100 |
| 12  | L     | 42/70 (60%)     | 36 (86%)   | 4 (10%)  | 2 (5%)   | 2           | 16  |
| 13  | M     | 106/415 (26%)   | 96 (91%)   | 8 (8%)   | 2 (2%)   | 6           | 32  |
| 14  | N     | 139/233 (60%)   | 123 (88%)  | 13 (9%)  | 3 (2%)   | 5           | 29  |
| 15  | O     | 467/627 (74%)   | 426 (91%)  | 35 (8%)  | 6 (1%)   | 9           | 42  |
| 16  | P     | 569/894 (64%)   | 473 (83%)  | 76 (13%) | 20 (4%)  | 3           | 20  |
| 17  | Q     | 398/514 (77%)   | 347 (87%)  | 41 (10%) | 10 (2%)  | 4           | 26  |
| 18  | R     | 289/507 (57%)   | 240 (83%)  | 39 (14%) | 10 (4%)  | 3           | 20  |
| All | All   | 5859/7778 (75%) | 5369 (92%) | 392 (7%) | 98 (2%)  | 9           | 36  |

5 of 98 Ramachandran outliers are listed below:

| Mol | Chain | Res  | Type |
|-----|-------|------|------|
| 1   | A     | 408  | LYS  |
| 1   | A     | 479  | ALA  |
| 1   | A     | 1651 | THR  |
| 2   | B     | 111  | ASP  |

*Continued on next page...*

*Continued from previous page...*

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 2   | B     | 895 | PHE  |

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed        | Rotameric  | Outliers  | Percentiles |    |
|-----|-------|-----------------|------------|-----------|-------------|----|
| 1   | A     | 1297/1465 (88%) | 1205 (93%) | 92 (7%)   | 13          | 35 |
| 2   | B     | 1025/1053 (97%) | 962 (94%)  | 63 (6%)   | 17          | 38 |
| 3   | C     | 269/296 (91%)   | 254 (94%)  | 15 (6%)   | 19          | 40 |
| 4   | D     | 55/116 (47%)    | 50 (91%)   | 5 (9%)    | 9           | 27 |
| 5   | E     | 194/197 (98%)   | 179 (92%)  | 15 (8%)   | 12          | 32 |
| 6   | F     | 90/137 (66%)    | 85 (94%)   | 5 (6%)    | 19          | 40 |
| 7   | G     | 171/291 (59%)   | 158 (92%)  | 13 (8%)   | 12          | 33 |
| 8   | H     | 115/128 (90%)   | 107 (93%)  | 8 (7%)    | 14          | 35 |
| 9   | I     | 89/110 (81%)    | 84 (94%)   | 5 (6%)    | 19          | 40 |
| 10  | J     | 64/65 (98%)     | 56 (88%)   | 8 (12%)   | 4           | 16 |
| 11  | K     | 91/130 (70%)    | 82 (90%)   | 9 (10%)   | 7           | 24 |
| 12  | L     | 39/57 (68%)     | 34 (87%)   | 5 (13%)   | 4           | 15 |
| 13  | M     | 98/371 (26%)    | 85 (87%)   | 13 (13%)  | 4           | 14 |
| 14  | N     | 135/220 (61%)   | 128 (95%)  | 7 (5%)    | 21          | 42 |
| 15  | O     | 439/576 (76%)   | 383 (87%)  | 56 (13%)  | 4           | 15 |
| 16  | P     | 539/828 (65%)   | 386 (72%)  | 153 (28%) | 0           | 2  |
| 17  | Q     | 383/476 (80%)   | 320 (84%)  | 63 (16%)  | 2           | 10 |
| 18  | R     | 286/474 (60%)   | 223 (78%)  | 63 (22%)  | 1           | 6  |
| All | All   | 5379/6990 (77%) | 4781 (89%) | 598 (11%) | 8           | 19 |

5 of 598 residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 16  | P     | 748 | GLU  |

*Continued on next page...*

*Continued from previous page...*

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 18  | R     | 266 | SER  |
| 17  | Q     | 101 | LYS  |
| 16  | P     | 736 | ILE  |
| 17  | Q     | 379 | LYS  |

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 71 such sidechains are listed below:

| Mol | Chain | Res  | Type |
|-----|-------|------|------|
| 16  | P     | 228  | ASN  |
| 16  | P     | 461  | HIS  |
| 18  | R     | 150  | GLN  |
| 2   | B     | 1163 | GLN  |
| 2   | B     | 893  | ASN  |

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

| Mol | Chain | Number of breaks |
|-----|-------|------------------|
| 1   | A     | 2                |
| 7   | G     | 1                |
| 18  | R     | 1                |
| 9   | I     | 1                |

All chain breaks are listed below:

| Model | Chain | Residue-1 | Atom-1 | Residue-2 | Atom-2 | Distance (Å) |
|-------|-------|-----------|--------|-----------|--------|--------------|
| 1     | G     | 24:VAL    | C      | 25:THR    | N      | 4.53         |
| 1     | R     | 206:ARG   | C      | 207:ASN   | N      | 3.86         |
| 1     | I     | 45:LEU    | C      | 46:LYS    | N      | 3.17         |
| 1     | A     | 438:ILE   | C      | 439:ASP   | N      | 2.97         |
| 1     | A     | 991:LYS   | C      | 992:PRO   | N      | 2.86         |


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-3592. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

### 6.2 Central slices [i](#)


#### 6.2.1 Primary map



X Index: 120



Y Index: 120




Z Index: 120


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 109



Y Index: 102




Z Index: 123


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

## 6.5 Orthogonal surface views

This section was not generated.


## 6.6 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 442 nm<sup>3</sup>; this corresponds to an approximate mass of 399 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum ⓘ



\*Reported resolution corresponds to spatial frequency of 0.130 Å<sup>-1</sup>

## 8 Fourier-Shell correlation ⓘ

This section was not generated. No FSC curve or half-maps provided.


## 9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-3592 and PDB model 5N60. Per-residue inclusion information can be found in section 3 on page 8.

### 9.1 Map-model overlay [i](#)

This section was not generated.

### 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

### 9.3 Atom inclusion mapped to coordinate model [i](#)

This section was not generated.

## 9.4 Atom inclusion [i](#)



At the recommended contour level, 70% of all backbone atoms, 52% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.023) and Q-score for the entire model and for each chain.

| Chain | Atom inclusion                                                                             | Q-score                                                                                    |
|-------|--------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------|
| All   |  0.5240   |  0.1020   |
| A     |  0.5990   |  0.1190   |
| B     |  0.5970   |  0.1100   |
| C     |  0.6350   |  0.1230   |
| D     |  0.6460   |  0.1470   |
| E     |  0.6680   |  0.1310   |
| F     |  0.6300   |  0.1280   |
| G     |  0.6200   |  0.1060   |
| H     |  0.6910   |  0.1320   |
| I     |  0.4580   |  0.0960   |
| J     |  0.6230   |  0.1170   |
| K     |  0.6060   |  0.1280   |
| L     |  0.5500   |  0.0870   |
| M     |  0.5550   |  0.0870   |
| N     |  0.5190  |  0.1180  |
| O     |  0.5970 |  0.1220 |
| P     |  0.2460 |  0.0390 |
| Q     |  0.1910 |  0.0430 |
| R     |  0.3430 |  0.0820 |

