



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 10:13 AM UTC

PDB ID : 5N6B / pdb_00005n6b
Title : Human Leukocyte Antigen Class I A02 Carrying LLWNPGMAV
Authors : Rizkallah, P.J.; Cole, D.K.; Fuller, A.; Bulek, A.M.; Bovay, A.
Deposited on : 2017-02-14
Resolution : 1.71 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.71 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1039 (1.72-1.72)
Clashscore	190562	1049 (1.72-1.72)
Ramachandran outliers	187476	1041 (1.72-1.72)
Sidechain outliers	187428	1041 (1.72-1.72)
RSRZ outliers	180081	1039 (1.72-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	276	 2% 89% 9%
1	D	276	 3% 88% 12%
2	B	100	 3% 93% 5%
2	E	100	 3% 88% 12%
3	C	9	 78% 22%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	9	78% 22%

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 7416 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class I histocompatibility antigen, A-2 alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	276	Total	C	N	O	S	0	8	0
			2313	1445	418	440	10			
1	D	276	Total	C	N	O	S	0	9	0
			2319	1449	419	441	10			

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	100	Total	C	N	O	S	0	5	0
			877	557	148	167	5			
2	E	100	Total	C	N	O	S	0	3	0
			857	546	143	164	4			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	initiating methionine	UNP P61769
E	0	MET	-	initiating methionine	UNP P61769

- Molecule 3 is a protein called NS5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	9	Total	C	N	O	S	0	0	0
			70	47	11	11	1			
3	F	9	Total	C	N	O	S	0	0	0
			70	47	11	11	1			

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		
4	E	1	Total	C	O	0	0
			4	2	2		
4	E	1	Total	C	O	0	0
			4	2	2		
4	F	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: C₈H₁₈N₂O₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	S	0	1
			30	16	4	8	2		
5	D	1	Total	C	N	O	S	0	1
			30	16	4	8	2		

- Molecule 6 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			7	4	3		
6	A	1	Total	C	O	0	0
			7	4	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	E	1	Total	C	O	0	0
			7	4	3		

- Molecule 7 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	O	S	0	0
			5	4	1		
7	A	1	Total	O	S	0	0
			5	4	1		
7	A	1	Total	O	S	0	0
			5	4	1		
7	A	1	Total	O	S	0	0
			5	4	1		
7	B	1	Total	O	S	0	0
			5	4	1		
7	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 8 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	E	1	Total	C	O	0	0
			6	3	3		

- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	243	Total	O	0	0
			243	243		
9	B	125	Total	O	0	0
			125	125		
9	C	9	Total	O	0	0
			9	9		
9	D	268	Total	O	0	0
			268	268		
9	E	97	Total	O	0	0
			97	97		
9	F	7	Total	O	0	0
			7	7		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: HLA class I histocompatibility antigen, A-2 alpha chain

Chain A: 





- Molecule 1: HLA class I histocompatibility antigen, A-2 alpha chain

Chain D: 



- Molecule 2: Beta-2-microglobulin

Chain B: 




- Molecule 2: Beta-2-microglobulin

Chain E: 



- Molecule 3: NS5

Chain C: 



- Molecule 3: NS5

Chain F:  78% 22%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	118.09Å 168.99Å 49.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	96.80 – 1.71 96.80 – 1.71	Depositor EDS
% Data completeness (in resolution range)	100.0 (96.80-1.71) 100.0 (96.80-1.71)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.91 (at 1.71Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.183 , 0.214 (Not available) , 0.218	Depositor DCC
R_{free} test set	5240 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å ²)	23.5	Xtriage
Anisotropy	0.280	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 30.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7416	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 41.25 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.4273e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EPE, EDO, GOL, SO4, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.12	3/2382 (0.1%)	1.08	5/3233 (0.2%)
1	D	1.14	5/2391 (0.2%)	1.03	2/3245 (0.1%)
2	B	1.19	1/903 (0.1%)	1.03	2/1219 (0.2%)
2	E	1.15	1/883 (0.1%)	1.08	4/1195 (0.3%)
3	C	1.36	0/72	1.65	2/97 (2.1%)
3	F	1.41	0/72	1.36	1/97 (1.0%)
All	All	1.15	10/6703 (0.1%)	1.07	16/9086 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	202	ARG	CD-NE	-8.30	1.34	1.46
1	D	39	ASP	CG-OD1	7.28	1.39	1.25
2	E	78	TYR	N-CA	6.15	1.53	1.45
2	B	1	ILE	C-O	5.75	1.31	1.24
1	D	76	VAL	N-CA	5.60	1.53	1.46
1	A	256	ARG	CD-NE	-5.49	1.38	1.46
1	D	275	GLU	N-CA	5.33	1.53	1.46
1	D	9	PHE	C-O	5.27	1.29	1.24
1	A	256	ARG	CZ-NH2	-5.14	1.26	1.33
1	D	19	GLU	CA-C	5.04	1.59	1.53

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	202	ARG	NE-CZ-NH1	-15.19	106.31	121.50
1	A	202	ARG	NE-CZ-NH2	14.75	132.48	119.20
1	A	202	ARG	CD-NE-CZ	7.58	135.02	124.40
3	C	5	GLY	CA-C-N	-7.07	112.50	119.78
3	C	5	GLY	C-N-CA	-7.07	112.50	119.78
1	D	274	TRP	N-CA-C	6.72	118.30	110.97
1	A	157	ARG	NE-CZ-NH2	6.66	125.20	119.20
2	E	81	ARG	CG-CD-NE	-6.08	98.63	112.00
2	E	74	GLU	CB-CA-C	-5.95	101.29	110.81
1	D	41	ALA	N-CA-C	5.51	116.97	111.07
2	B	74	GLU	CB-CA-C	-5.47	101.67	110.74
2	E	97	ARG	NE-CZ-NH2	5.37	124.03	119.20
2	E	0	MET	N-CA-CB	-5.20	101.67	110.50
2	B	99	MET	N-CA-C	5.12	125.33	111.00
1	A	157	ARG	NE-CZ-NH1	-5.08	116.42	121.50
3	F	5	GLY	N-CA-C	5.00	122.54	112.34

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	202	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2313	0	2157	23	0
1	D	2319	0	2165	22	0
2	B	877	0	842	4	0
2	E	857	0	824	6	0
3	C	70	0	73	1	0
3	F	70	0	73	1	0
4	A	20	0	30	0	0
4	B	8	0	12	0	0
4	D	4	0	6	0	0
4	E	8	0	12	0	0
4	F	4	0	6	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	30	0	35	5	0
5	D	30	0	36	8	0
6	A	14	0	20	1	0
6	E	7	0	10	1	0
7	A	20	0	0	0	0
7	B	5	0	0	0	0
7	D	5	0	0	0	0
8	E	6	0	8	2	0
9	A	243	0	0	6	0
9	B	125	0	0	1	0
9	C	9	0	0	0	0
9	D	268	0	0	3	0
9	E	97	0	0	1	0
9	F	7	0	0	0	0
All	All	7416	0	6309	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:255:GLN:HE22	1:D:275:GLU:HG2	1.43	0.82
2:E:29:GLY:O	8:E:101:GOL:H31	1.91	0.70
5:D:302[A]:EPE:S	5:D:302[A]:EPE:H61	2.33	0.67
1:A:115[B]:GLN:NE2	9:A:403:HOH:O	2.31	0.63
5:D:302[A]:EPE:O3S	5:D:302[A]:EPE:H62	1.99	0.62
1:A:248[A]:VAL:O	1:A:248[A]:VAL:HG13	2.02	0.60
1:A:227:ASP:HB3	1:A:248[A]:VAL:HG12	1.84	0.60
1:D:248[A]:VAL:HG13	1:D:248[A]:VAL:O	2.00	0.60
1:D:248[A]:VAL:O	1:D:248[A]:VAL:CG1	2.51	0.58
1:A:248[A]:VAL:O	1:A:248[A]:VAL:CG1	2.52	0.57
1:D:98:MET:HE2	6:E:104:PEG:H22	1.88	0.56
1:D:227:ASP:HB3	1:D:248[A]:VAL:HG12	1.88	0.56
1:A:17:ARG:NH1	9:A:404:HOH:O	2.32	0.55
5:D:302[A]:EPE:O3S	5:D:302[A]:EPE:C6	2.55	0.55
5:D:302[A]:EPE:S	5:D:302[A]:EPE:C6	2.96	0.52
1:D:17:ARG:NH2	9:D:405:HOH:O	2.42	0.51
1:D:74:HIS:CE1	1:D:97:ARG:HE	2.29	0.51
1:A:191:HIS:HD2	9:A:601:HOH:O	1.93	0.51
1:D:121:LYS:NZ	9:D:402:HOH:O	2.31	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:191:HIS:HE1	1:D:254:GLU:OE1	1.93	0.51
1:D:25[B]:VAL:HG13	1:D:32:GLN:NE2	2.26	0.50
2:E:31:HIS:HD2	9:E:212:HOH:O	1.92	0.50
1:A:202:ARG:HD3	1:A:244:TRP:CE3	2.46	0.50
1:A:152:VAL:HG22	3:C:7:MET:HG3	1.93	0.50
2:B:31:HIS:HD2	9:B:206:HOH:O	1.95	0.49
1:A:191:HIS:HE1	1:A:254:GLU:OE1	1.96	0.49
5:D:302[B]:EPE:H31	5:D:302[B]:EPE:H91	1.45	0.49
1:A:41:ALA:HB3	5:A:306[A]:EPE:H72	1.95	0.48
1:D:232[B]:GLU:HG2	2:E:8:GLN:NE2	2.28	0.48
1:D:191:HIS:HD2	9:D:625:HOH:O	1.95	0.48
5:D:302[A]:EPE:H82	5:D:302[A]:EPE:H51	1.32	0.48
1:A:11[B]:SER:OG	1:A:22:PHE:HD1	1.97	0.47
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.49	0.47
1:D:96:GLN:OE1	2:E:31:HIS:HE1	1.97	0.47
1:A:96:GLN:OE1	2:B:31:HIS:HE1	1.98	0.47
1:A:74:HIS:CE1	1:A:97:ARG:HE	2.32	0.47
1:D:255:GLN:NE2	1:D:275:GLU:HG2	2.23	0.47
1:D:219:ARG:HH22	1:D:253:GLN:HE22	1.63	0.46
1:A:11[B]:SER:OG	9:A:401:HOH:O	2.19	0.45
1:A:121:LYS:HB2	6:A:307:PEG:H32	1.98	0.45
1:A:35:ARG:HH22	5:A:306[A]:EPE:H92	1.80	0.45
1:D:232[B]:GLU:OE2	2:E:28:SER:OG	2.27	0.44
1:A:202:ARG:HD2	1:A:244:TRP:CD2	2.53	0.44
1:D:41:ALA:H	5:D:302[A]:EPE:H51	1.82	0.44
1:D:66:LYS:O	1:D:70:HIS:HD2	2.00	0.44
5:A:306[B]:EPE:H101	9:A:622:HOH:O	2.18	0.43
3:F:7:MET:HE3	3:F:7:MET:HB2	1.87	0.43
2:E:29:GLY:O	8:E:101:GOL:C3	2.66	0.42
1:A:230:LEU:HD13	9:A:549:HOH:O	2.20	0.42
2:B:0[B]:MET:HE2	2:B:0[B]:MET:CA	2.50	0.42
1:A:41:ALA:H	5:A:306[B]:EPE:C3	2.34	0.41
1:D:255:GLN:OE1	1:D:275:GLU:N	2.40	0.41
5:A:306[A]:EPE:H81	5:A:306[A]:EPE:H51	1.85	0.41
1:A:196:ASP:OD1	1:A:196:ASP:N	2.54	0.41
1:D:40:ALA:HB1	5:D:302[B]:EPE:H51	2.03	0.41
1:D:165[B]:VAL:HG12	1:D:169:ARG:CZ	2.51	0.40
1:D:74:HIS:HE1	1:D:97:ARG:HE	1.67	0.40
1:A:228:THR:HG23	1:A:229:GLU:N	2.37	0.40
1:A:74:HIS:HD2	1:A:95:VAL:HG11	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	282/276 (102%)	279 (99%)	3 (1%)	0	100	100
1	D	283/276 (102%)	279 (99%)	4 (1%)	0	100	100
2	B	102/100 (102%)	102 (100%)	0	0	100	100
2	E	101/100 (101%)	101 (100%)	0	0	100	100
3	C	7/9 (78%)	7 (100%)	0	0	100	100
3	F	7/9 (78%)	7 (100%)	0	0	100	100
All	All	782/770 (102%)	775 (99%)	7 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	240/232 (103%)	231 (96%)	9 (4%)	29	8
1	D	241/232 (104%)	234 (97%)	7 (3%)	37	14
2	B	100/95 (105%)	97 (97%)	3 (3%)	36	12
2	E	98/95 (103%)	95 (97%)	3 (3%)	35	12
3	C	7/7 (100%)	7 (100%)	0	100	100
3	F	7/7 (100%)	7 (100%)	0	100	100
All	All	693/668 (104%)	671 (97%)	22 (3%)	35	11

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11[A]	SER
1	A	11[B]	SER
1	A	35	ARG
1	A	177	GLU
1	A	196	ASP
1	A	219	ARG
1	A	228	THR
1	A	251	SER
1	A	268	LYS
2	B	1	ILE
2	B	98	ASP
2	B	99	MET
1	D	11	SER
1	D	35	ARG
1	D	181	ARG
1	D	196	ASP
1	D	251	SER
1	D	268	LYS
1	D	275	GLU
2	E	19	LYS
2	E	48	LYS
2	E	75	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (25) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	74	HIS
1	A	86	ASN
1	A	141	GLN
1	A	174	ASN
1	A	191	HIS
1	A	226	GLN
1	A	253	GLN
1	A	255	GLN
2	B	13	HIS
2	B	31	HIS
2	B	83	ASN
1	D	43	GLN
1	D	70	HIS
1	D	74	HIS
1	D	86	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	141	GLN
1	D	151	HIS
1	D	174	ASN
1	D	191	HIS
1	D	192	HIS
1	D	226	GLN
1	D	253	GLN
2	E	13	HIS
2	E	31	HIS
2	E	83	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

25 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
8	GOL	E	101	-	5,5,5	1.34	1 (20%)	5,5,5	1.63	2 (40%)
7	SO4	A	309	-	4,4,4	0.47	0	6,6,6	0.33	0
4	EDO	A	304	-	3,3,3	0.41	0	2,2,2	0.37	0
4	EDO	E	102	-	3,3,3	0.42	0	2,2,2	0.63	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	EPE	D	302[B]	-	15,15,15	2.22	1 (6%)	19,20,20	1.31	2 (10%)
7	SO4	A	311	-	4,4,4	0.58	0	6,6,6	0.33	0
5	EPE	A	306[A]	-	15,15,15	2.36	1 (6%)	19,20,20	1.65	5 (26%)
4	EDO	F	101	-	3,3,3	0.59	0	2,2,2	0.31	0
4	EDO	B	101	-	3,3,3	0.42	0	2,2,2	0.53	0
4	EDO	D	301	-	3,3,3	0.24	0	2,2,2	0.94	0
6	PEG	A	307	-	6,6,6	0.60	0	5,5,5	0.30	0
6	PEG	A	308	-	6,6,6	0.66	0	5,5,5	0.46	0
4	EDO	B	102	-	3,3,3	0.37	0	2,2,2	0.68	0
7	SO4	A	312	-	4,4,4	0.47	0	6,6,6	0.26	0
7	SO4	B	103	-	4,4,4	0.43	0	6,6,6	0.27	0
7	SO4	D	303	-	4,4,4	0.50	0	6,6,6	0.40	0
6	PEG	E	104	-	6,6,6	0.56	0	5,5,5	1.04	0
7	SO4	A	310	-	4,4,4	0.61	0	6,6,6	0.70	0
5	EPE	A	306[B]	-	15,15,15	2.01	1 (6%)	19,20,20	2.02	6 (31%)
4	EDO	A	301	-	3,3,3	0.49	0	2,2,2	0.17	0
5	EPE	D	302[A]	-	15,15,15	2.24	2 (13%)	19,20,20	1.82	4 (21%)
4	EDO	A	303	-	3,3,3	0.40	0	2,2,2	0.59	0
4	EDO	E	103	-	3,3,3	0.44	0	2,2,2	0.34	0
4	EDO	A	302	-	3,3,3	0.41	0	2,2,2	0.30	0
4	EDO	A	305	-	3,3,3	0.43	0	2,2,2	0.20	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	GOL	E	101	-	-	2/4/4/4	-
4	EDO	A	304	-	-	1/1/1/1	-
4	EDO	E	102	-	-	1/1/1/1	-
5	EPE	D	302[B]	-	-	9/9/19/19	0/1/1/1
5	EPE	A	306[A]	-	-	7/9/19/19	0/1/1/1
4	EDO	F	101	-	-	0/1/1/1	-
4	EDO	B	101	-	-	0/1/1/1	-
4	EDO	D	301	-	-	1/1/1/1	-
6	PEG	A	307	-	-	2/4/4/4	-
6	PEG	A	308	-	-	3/4/4/4	-
4	EDO	B	102	-	-	1/1/1/1	-
6	PEG	E	104	-	-	1/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EPE	A	306[B]	-	-	5/9/19/19	0/1/1/1
4	EDO	A	301	-	-	1/1/1/1	-
5	EPE	D	302[A]	-	-	7/9/19/19	0/1/1/1
4	EDO	A	303	-	-	0/1/1/1	-
4	EDO	E	103	-	-	0/1/1/1	-
4	EDO	A	302	-	-	0/1/1/1	-
4	EDO	A	305	-	-	0/1/1/1	-

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	306[A]	EPE	C10-S	-8.81	1.65	1.77
5	D	302[B]	EPE	C10-S	-8.35	1.65	1.77
5	D	302[A]	EPE	C10-S	-8.19	1.66	1.77
5	A	306[B]	EPE	C10-S	-7.33	1.67	1.77
8	E	101	GOL	O2-C2	2.60	1.50	1.43
5	D	302[A]	EPE	O2S-S	2.13	1.51	1.45

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	D	302[A]	EPE	O2S-S-C10	4.63	113.72	106.73
5	A	306[A]	EPE	C6-C5-N4	-3.87	102.84	110.65
5	D	302[A]	EPE	C9-N1-C6	-3.55	101.78	111.24
5	A	306[B]	EPE	C6-N1-C2	-3.38	101.57	108.84
5	D	302[B]	EPE	C3-C2-N1	-3.33	103.93	110.65
5	A	306[B]	EPE	C7-N4-C3	-3.33	102.36	111.24
5	A	306[B]	EPE	C5-C6-N1	-3.27	104.06	110.65
5	A	306[B]	EPE	O1S-S-C10	3.00	111.26	106.73
5	A	306[A]	EPE	O2S-S-C10	2.68	110.78	106.73
5	D	302[A]	EPE	C10-C9-N1	-2.68	102.23	112.36
5	A	306[B]	EPE	C3-C2-N1	-2.67	105.27	110.65
5	A	306[A]	EPE	C2-C3-N4	2.62	115.94	110.65
5	A	306[B]	EPE	C2-C3-N4	2.62	115.93	110.65
5	D	302[A]	EPE	C7-N4-C5	-2.60	104.32	111.24
8	E	101	GOL	O1-C1-C2	2.57	121.94	110.38
5	A	306[A]	EPE	C5-C6-N1	-2.24	106.14	110.65
5	D	302[B]	EPE	O1S-S-C10	2.02	109.78	106.73
8	E	101	GOL	O3-C3-C2	2.01	119.44	110.38
5	A	306[A]	EPE	C7-N4-C5	-2.01	105.89	111.24

There are no chirality outliers.

All (41) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	306[A]	EPE	C10-C9-N1-C2
5	A	306[A]	EPE	S-C10-C9-N1
5	A	306[A]	EPE	C9-C10-S-O2S
5	A	306[A]	EPE	C9-C10-S-O3S
5	A	306[B]	EPE	C9-C10-S-O3S
5	D	302[A]	EPE	S-C10-C9-N1
5	D	302[A]	EPE	C9-C10-S-O1S
5	D	302[A]	EPE	C9-C10-S-O3S
5	D	302[B]	EPE	C9-C10-S-O2S
5	A	306[B]	EPE	N4-C7-C8-O8
5	D	302[A]	EPE	C8-C7-N4-C5
5	A	306[A]	EPE	N4-C7-C8-O8
6	A	307	PEG	O1-C1-C2-O2
6	A	308	PEG	O2-C3-C4-O4
5	D	302[B]	EPE	N4-C7-C8-O8
8	E	101	GOL	O1-C1-C2-C3
6	A	307	PEG	O2-C3-C4-O4
8	E	101	GOL	O1-C1-C2-O2
5	D	302[A]	EPE	N4-C7-C8-O8
5	D	302[B]	EPE	C9-C10-S-O3S
5	A	306[B]	EPE	S-C10-C9-N1
6	A	308	PEG	O1-C1-C2-O2
5	A	306[A]	EPE	C10-C9-N1-C6
6	E	104	PEG	O1-C1-C2-O2
5	D	302[A]	EPE	C8-C7-N4-C3
5	D	302[B]	EPE	C8-C7-N4-C3
5	A	306[A]	EPE	C9-C10-S-O1S
5	A	306[B]	EPE	C9-C10-S-O1S
5	A	306[B]	EPE	C9-C10-S-O2S
5	D	302[A]	EPE	C9-C10-S-O2S
5	D	302[B]	EPE	C9-C10-S-O1S
5	D	302[B]	EPE	S-C10-C9-N1
5	D	302[B]	EPE	C10-C9-N1-C2
5	D	302[B]	EPE	C10-C9-N1-C6
5	D	302[B]	EPE	C8-C7-N4-C5
4	D	301	EDO	O1-C1-C2-O2
6	A	308	PEG	C4-C3-O2-C2
4	B	102	EDO	O1-C1-C2-O2
4	A	301	EDO	O1-C1-C2-O2
4	A	304	EDO	O1-C1-C2-O2
4	E	102	EDO	O1-C1-C2-O2

There are no ring outliers.

7 monomers are involved in 17 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	E	101	GOL	2	0
5	D	302[B]	EPE	2	0
5	A	306[A]	EPE	3	0
6	A	307	PEG	1	0
6	E	104	PEG	1	0
5	A	306[B]	EPE	2	0
5	D	302[A]	EPE	6	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	276/276 (100%)	0.23	6 (2%) 62 70	8, 27, 53, 71	8 (2%)
1	D	276/276 (100%)	0.17	7 (2%) 58 66	10, 27, 48, 64	9 (3%)
2	B	100/100 (100%)	0.15	3 (3%) 52 61	9, 24, 47, 60	5 (5%)
2	E	100/100 (100%)	0.14	3 (3%) 52 61	10, 27, 50, 70	3 (3%)
3	C	9/9 (100%)	-0.11	0 100 100	21, 24, 27, 29	0
3	F	9/9 (100%)	0.44	0 100 100	21, 22, 33, 38	0
All	All	770/770 (100%)	0.19	19 (2%) 58 66	8, 27, 50, 71	25 (3%)

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	276	PRO	3.9
1	A	228	THR	3.4
1	D	274	TRP	3.2
1	A	227	ASP	3.1
1	A	225	THR	3.0
2	B	99	MET	2.8
2	E	98	ASP	2.8
1	A	177	GLU	2.6
2	B	98	ASP	2.6
1	D	276	PRO	2.5
1	D	228	THR	2.5
1	A	230	LEU	2.4
1	D	227	ASP	2.3
2	E	99	MET	2.3
1	D	225	THR	2.2
2	E	0	MET	2.2
1	D	275	GLU	2.2
1	D	182	THR	2.1
2	B	1	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	EPE	D	302[A]	15/15	0.66	0.28	29,41,50,52	15
5	EPE	D	302[B]	15/15	0.66	0.28	32,44,58,59	15
6	PEG	A	308	7/7	0.70	0.24	52,58,60,62	0
7	SO4	A	312	5/5	0.71	0.18	98,101,106,107	0
7	SO4	A	311	5/5	0.75	0.16	65,67,81,86	0
7	SO4	B	103	5/5	0.75	0.16	94,95,100,101	0
4	EDO	A	302	4/4	0.77	0.19	62,63,64,66	0
4	EDO	E	103	4/4	0.78	0.19	58,59,60,61	0
5	EPE	A	306[A]	15/15	0.78	0.25	41,47,55,56	15
5	EPE	A	306[B]	15/15	0.78	0.25	24,34,47,47	15
8	GOL	E	101	6/6	0.80	0.21	17,20,21,23	6
4	EDO	B	102	4/4	0.81	0.19	45,46,48,49	0
4	EDO	A	303	4/4	0.82	0.20	40,42,44,50	0
6	PEG	A	307	7/7	0.83	0.18	28,43,55,55	0
4	EDO	D	301	4/4	0.83	0.16	40,40,41,41	0
7	SO4	D	303	5/5	0.84	0.10	62,63,65,73	0
4	EDO	A	304	4/4	0.86	0.15	54,55,56,58	0
7	SO4	A	310	5/5	0.87	0.11	46,49,53,53	0
6	PEG	E	104	7/7	0.87	0.14	38,40,42,43	0
4	EDO	F	101	4/4	0.88	0.12	36,40,42,46	0
4	EDO	A	301	4/4	0.90	0.12	38,39,40,42	0
4	EDO	E	102	4/4	0.91	0.12	31,36,39,43	0
4	EDO	B	101	4/4	0.93	0.10	31,33,35,38	0
7	SO4	A	309	5/5	0.93	0.09	42,45,49,52	0
4	EDO	A	305	4/4	0.96	0.10	25,26,28,30	0

6.5 Other polymers [i](#)

There are no such residues in this entry.