



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 11:57 PM UTC

PDB ID : 6N6R / pdb_00006n6r
Title : Crystal structure of ABIN-1 UBAN in complex with two M1-linked di-ubiquitins
Authors : Rahighi, S.; Dikic, I.; Wakatsuki, S.
Deposited on : 2018-11-27
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	157	32% 89% 6%
1	C	157	40% 88% 6%
2	B	70	30% 87% 10%
2	D	70	26% 86% 7%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3653 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ubiquitin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	148	Total	C	N	O	S	0	0	0
			1175	740	203	230	2			
1	C	150	Total	C	N	O	S	0	0	0
			1194	752	208	232	2			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	expression tag	UNP P0CG48
A	-3	SER	-	expression tag	UNP P0CG48
A	-2	GLY	-	expression tag	UNP P0CG48
A	-1	SER	-	expression tag	UNP P0CG48
C	-4	GLY	-	expression tag	UNP P0CG48
C	-3	SER	-	expression tag	UNP P0CG48
C	-2	GLY	-	expression tag	UNP P0CG48
C	-1	SER	-	expression tag	UNP P0CG48

- Molecule 2 is a protein called TNFAIP3-interacting protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	68	Total	C	N	O	S	0	0	0
			580	356	106	117	1			
2	D	67	Total	C	N	O	S	0	0	0
			569	350	102	116	1			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	42	Total	O	0	0
			42	42		
3	B	34	Total	O	0	0
			34	34		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	25	Total	O	0	0
			25	25		
3	C	34	Total	O	0	0
			34	34		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Ubiquitin




• Molecule 1: Ubiquitin



• Molecule 2: TNFAIP3-interacting protein 1



• Molecule 2: TNFAIP3-interacting protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	52.90Å 88.18Å 105.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.00 – 1.95 48.00 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.6 (48.00-1.95) 99.6 (48.00-1.95)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.64 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.243 , 0.278 0.250 , 0.285	Depositor DCC
R_{free} test set	1890 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å ²)	27.0	Xtriage
Anisotropy	0.090	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 39.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	3653	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.54	0/1188	0.77	0/1600
1	C	0.60	0/1207	0.86	3/1625 (0.2%)
2	B	0.55	0/581	0.91	0/769
2	D	0.53	0/570	0.99	2/755 (0.3%)
All	All	0.56	0/3546	0.86	5/4749 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	21	ASP	N-CA-C	-6.48	100.69	110.28
1	C	36	ILE	N-CA-C	5.53	113.96	107.77
1	C	63	LYS	N-CA-C	5.04	116.47	111.07
2	D	515	THR	N-CA-CB	5.03	117.61	110.16
2	D	523	THR	N-CA-CB	5.00	117.66	110.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1175	0	1226	5	0
1	C	1194	0	1250	3	0
2	B	580	0	599	1	0
2	D	569	0	586	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	42	0	0	0	0
3	B	34	0	0	1	0
3	C	34	0	0	1	0
3	D	25	0	0	0	0
All	All	3653	0	3661	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:20:SER:HA	1:C:55:THR:HB	1.74	0.68
2:D:519:ALA:O	2:D:523:THR:HG23	2.05	0.56
1:A:90:THR:HG22	2:B:494:GLU:HG2	1.88	0.55
3:B:622:HOH:O	2:D:496:MET:HE2	2.07	0.53
1:A:3:ILE:HD12	1:A:17:VAL:HG21	1.94	0.50
1:C:55:THR:O	1:C:58:ASP:HB2	2.14	0.47
1:A:42:ARG:NH1	2:D:481:ILE:HD11	2.31	0.46
1:A:42:ARG:CZ	2:D:481:ILE:HD11	2.46	0.46
2:D:511:GLN:O	2:D:515:THR:HG23	2.18	0.44
1:C:20:SER:HB3	3:C:201:HOH:O	2.18	0.43
1:A:98:THR:HA	1:A:131:THR:HA	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	146/157 (93%)	146 (100%)	0	0	100	100
1	C	148/157 (94%)	148 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	66/70 (94%)	66 (100%)	0	0	100	100
2	D	65/70 (93%)	65 (100%)	0	0	100	100
All	All	425/454 (94%)	425 (100%)	0	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	134/138 (97%)	131 (98%)	3 (2%)	45	40
1	C	136/138 (99%)	127 (93%)	9 (7%)	15	5
2	B	65/67 (97%)	59 (91%)	6 (9%)	8	2
2	D	64/67 (96%)	61 (95%)	3 (5%)	23	12
All	All	399/410 (97%)	378 (95%)	21 (5%)	20	9

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	THR
1	A	24	GLU
1	A	90	THR
2	B	464	ARG
2	B	468	LEU
2	B	474	LEU
2	B	481	ILE
2	B	498	GLU
2	B	508	GLU
2	D	515	THR
2	D	523	THR
2	D	529	LYS
1	C	9	THR
1	C	11	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	14	THR
1	C	20	SER
1	C	21	ASP
1	C	42	ARG
1	C	58	ASP
1	C	66	THR
1	C	71	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	477	GLN
2	B	487	GLN
2	D	471	GLN
1	C	2	GLN
1	C	138	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	148/157 (94%)	1.49	50 (33%) 1 0	13, 33, 64, 74	0
1	C	150/157 (95%)	1.82	63 (42%) 0 0	16, 39, 63, 79	0
2	B	68/70 (97%)	1.38	21 (30%) 1 1	17, 30, 66, 71	0
2	D	67/70 (95%)	1.44	18 (26%) 1 1	18, 28, 58, 85	0
All	All	433/454 (95%)	1.58	152 (35%) 1 0	13, 34, 65, 85	0

All (152) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	20	SER	6.1
1	A	10	GLY	5.9
1	C	12	THR	5.8
1	A	16	GLU	5.3
1	C	58	ASP	5.0
1	C	35	GLY	4.8
1	A	4	PHE	4.7
1	C	4	PHE	4.7
2	D	530	ALA	4.7
1	A	57	SER	4.7
1	C	13	ILE	4.6
1	C	22	THR	4.6
2	D	465	LYS	4.5
2	B	469	VAL	4.5
1	C	14	THR	4.2
2	B	466	GLN	4.2
2	B	473	GLU	4.2
1	A	47	GLY	4.1
1	C	10	GLY	4.1
1	A	85	THR	4.0
1	A	45	PHE	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	53	GLY	4.0
1	A	61	ILE	3.9
1	C	19	PRO	3.9
1	C	55	THR	3.7
1	C	61	ILE	3.7
1	A	19	PRO	3.6
1	A	1	MET	3.6
1	A	50	LEU	3.6
1	A	55	THR	3.6
2	B	528	GLU	3.6
1	C	63	LYS	3.6
2	D	466	GLN	3.5
2	D	528	GLU	3.5
1	A	148	ARG	3.5
1	C	115	ASP	3.5
2	D	469	VAL	3.5
1	C	59	TYR	3.4
1	A	8	LEU	3.4
1	C	3	ILE	3.3
1	C	15	LEU	3.3
2	D	468	LEU	3.3
2	D	508	GLU	3.3
1	A	17	VAL	3.3
1	A	12	THR	3.3
1	A	24	GLU	3.2
2	B	467	GLU	3.2
1	A	49	GLN	3.2
1	A	59	TYR	3.2
1	C	150	ARG	3.2
1	C	46	ALA	3.1
1	A	18	GLU	3.1
1	A	46	ALA	3.1
1	C	9	THR	3.1
1	A	15	LEU	3.1
1	C	2	GLN	3.0
1	A	64	GLU	3.0
1	A	9	THR	3.0
2	D	531	LYS	3.0
1	C	127	GLU	3.0
2	D	523	THR	3.0
1	C	51	GLU	3.0
1	A	136	ASN	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	149	LEU	2.9
1	A	130	ARG	2.9
1	C	1	MET	2.9
2	B	530	ALA	2.9
1	C	17	VAL	2.9
1	C	47	GLY	2.9
1	A	118	ARG	2.9
1	A	14	THR	2.9
1	C	60	ASN	2.9
1	A	54	ARG	2.9
1	A	100	GLU	2.8
1	A	63	LYS	2.8
1	C	21	ASP	2.8
2	B	468	LEU	2.8
1	A	51	GLU	2.8
2	D	467	GLU	2.8
1	A	3	ILE	2.8
1	C	26	VAL	2.8
1	C	62	GLN	2.7
2	D	476	LYS	2.7
2	D	473	GLU	2.7
1	C	8	LEU	2.7
1	A	20	SER	2.7
1	A	127	GLU	2.7
1	C	34	GLU	2.7
1	C	33	LYS	2.6
1	C	109	LYS	2.6
2	B	481	ILE	2.6
1	C	18	GLU	2.6
2	D	529	LYS	2.6
1	A	123	GLY	2.6
1	A	128	ASP	2.6
1	C	133	SER	2.6
2	D	496	MET	2.6
2	B	508	GLU	2.6
1	A	65	SER	2.6
1	C	113	PRO	2.5
2	B	470	THR	2.5
2	B	475	LEU	2.5
1	C	54	ARG	2.5
1	C	49	GLN	2.5
2	B	496	MET	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	31	GLN	2.5
2	D	480	LYS	2.5
2	B	531	LYS	2.4
1	A	5	VAL	2.4
1	A	62	GLN	2.4
1	A	52	ASP	2.4
1	C	24	GLU	2.4
1	A	44	ILE	2.4
2	B	502	GLU	2.4
1	C	57	SER	2.4
1	C	28	ALA	2.3
1	A	60	ASN	2.3
1	C	64	GLU	2.3
1	C	25	ASN	2.3
1	C	23	ILE	2.3
2	D	483	GLU	2.3
1	A	2	GLN	2.3
1	A	131	THR	2.3
1	C	131	THR	2.3
1	C	135	TYR	2.3
1	C	100	GLU	2.3
1	C	108	ASP	2.3
1	C	66	THR	2.3
2	B	529	LYS	2.3
1	C	7	THR	2.2
2	D	515	THR	2.2
1	A	21	ASP	2.2
1	C	130	ARG	2.2
2	D	470	THR	2.2
2	B	477	GLN	2.2
1	C	65	SER	2.2
1	A	108	ASP	2.2
1	C	128	ASP	2.2
2	B	524	LEU	2.2
2	B	472	ASN	2.1
1	C	94	GLU	2.1
1	A	89	ILE	2.1
1	C	96	SER	2.1
1	C	136	ASN	2.1
1	A	23	ILE	2.1
1	C	29	LYS	2.1
2	B	483	GLU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	484	GLU	2.1
1	C	56	LEU	2.1
1	C	112	ILE	2.1
2	B	464	ARG	2.1
1	C	124	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.