



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 12:07 PM UTC

PDB ID : 4N7M / pdb_00004n7m
Title : Tailoring Small Molecules for an Allosteric Site on Procaspase-6
Authors : Murray, J.M.; Steffek, M.
Deposited on : 2013-10-15
Resolution : 2.12 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.12 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	8290 (2.14-2.10)
Clashscore	190562	8817 (2.14-2.10)
Ramachandran outliers	187476	8738 (2.14-2.10)
Sidechain outliers	187428	8739 (2.14-2.10)
RSRZ outliers	180081	8294 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	282	9% 77% 9% • 9%
1	B	282	2% 78% 7% • 14%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PO4	A	403	-	X	-	-
3	PO4	B	401	-	X	-	-
3	PO4	B	402	-	X	-	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4298 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Caspase-6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	256	Total	C	N	O	S	0	1	0
			2044	1308	354	367	15			
1	B	243	Total	C	N	O	S	0	1	0
			1944	1243	335	352	14			

There are 26 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	21	MET	-	initiating methionine	UNP P55212
A	22	GLY	-	expression tag	UNP P55212
A	23	SER	-	expression tag	UNP P55212
A	163	ALA	CYS	engineered mutation	UNP P55212
A	294	GLY	-	expression tag	UNP P55212
A	295	ASN	-	expression tag	UNP P55212
A	296	SER	-	expression tag	UNP P55212
A	297	HIS	-	expression tag	UNP P55212
A	298	HIS	-	expression tag	UNP P55212
A	299	HIS	-	expression tag	UNP P55212
A	300	HIS	-	expression tag	UNP P55212
A	301	HIS	-	expression tag	UNP P55212
A	302	HIS	-	expression tag	UNP P55212
B	21	MET	-	initiating methionine	UNP P55212
B	22	GLY	-	expression tag	UNP P55212
B	23	SER	-	expression tag	UNP P55212
B	163	ALA	CYS	engineered mutation	UNP P55212
B	294	GLY	-	expression tag	UNP P55212
B	295	ASN	-	expression tag	UNP P55212
B	296	SER	-	expression tag	UNP P55212
B	297	HIS	-	expression tag	UNP P55212
B	298	HIS	-	expression tag	UNP P55212
B	299	HIS	-	expression tag	UNP P55212
B	300	HIS	-	expression tag	UNP P55212
B	301	HIS	-	expression tag	UNP P55212

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	302	HIS	-	expression tag	UNP P55212

- Molecule 2 is 3-(pyrrolidin-1-yl)isoquinolin-1(2H)-one (CCD ID: 2GQ) (formula: $C_{13}H_{14}N_2O$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			16	13	2	1		

- Molecule 3 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		
3	A	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	135	Total	O	0	0
			135	135		
4	B	134	Total	O	0	0
			134	134		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Caspase-6



• Molecule 1: Caspase-6



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	115.70Å 115.70Å 79.69Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	81.81 – 2.12 81.81 – 2.12	Depositor EDS
% Data completeness (in resolution range)	99.9 (81.81-2.12) 100.0 (81.81-2.12)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.28 (at 2.12Å)	Xtriage
Refinement program	PHENIX 1.8.2_1309	Depositor
R, R_{free}	0.159 , 0.199 0.170 , 0.204	Depositor DCC
R_{free} test set	1598 reflections (5.14%)	wwPDB-VP
Wilson B-factor (Å ²)	28.9	Xtriage
Anisotropy	0.041	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 48.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4298	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 2GQ, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.91	5/2096 (0.2%)	1.20	18/2827 (0.6%)
1	B	0.76	0/1992	0.91	4/2684 (0.1%)
All	All	0.84	5/4088 (0.1%)	1.07	22/5511 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	38	LYS	C-O	-6.14	1.16	1.24
1	A	268	SER	N-CA	-5.38	1.39	1.46
1	A	267	PRO	CA-C	-5.20	1.43	1.52
1	A	147	LYS	C-O	-5.07	1.18	1.24
1	A	202	ALA	C-O	-5.07	1.18	1.24

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	262	ASP	N-CA-C	13.25	125.72	111.28
1	A	265	LYS	N-CA-C	12.66	124.81	111.14
1	A	266	ASP	CA-C-N	11.44	131.87	119.28
1	A	266	ASP	C-N-CA	11.44	131.87	119.28
1	A	262	ASP	CA-C-N	11.03	141.56	121.70
1	A	262	ASP	C-N-CA	11.03	141.56	121.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	261	VAL	CA-C-N	10.47	134.31	120.28
1	A	261	VAL	C-N-CA	10.47	134.31	120.28
1	A	266	ASP	O-C-N	8.79	131.42	121.32
1	A	267	PRO	CA-C-N	-7.16	107.63	121.58
1	A	267	PRO	C-N-CA	-7.16	107.63	121.58
1	A	268	SER	N-CA-C	-6.70	105.07	113.18
1	B	125	ASN	N-CA-CB	-6.34	100.74	110.44
1	A	264	CYS	N-CA-C	6.08	117.91	111.28
1	A	167	GLN	N-CA-C	6.04	119.61	112.72
1	A	262	ASP	CA-C-O	-6.01	114.18	120.55
1	B	32	ASP	CA-C-N	5.84	125.54	119.82
1	B	32	ASP	C-N-CA	5.84	125.54	119.82
1	A	262	ASP	CB-CA-C	-5.81	101.14	110.79
1	A	202	ALA	N-CA-C	5.74	122.97	113.50
1	A	269	ALA	N-CA-C	-5.33	106.84	113.19
1	B	202	ALA	N-CA-C	5.15	122.47	114.64

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	261	VAL	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2044	0	1998	31	0
1	B	1944	0	1908	11	0
2	A	16	0	14	1	0
3	A	10	0	0	0	0
3	B	15	0	0	0	0
4	A	135	0	0	2	0
4	B	134	0	0	1	0
All	All	4298	0	3920	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 5.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:277:CYS:SG	4:A:620:HOH:O	2.34	0.85
1:A:261:VAL:HG23	1:A:262:ASP:H	1.52	0.75
1:A:285:LYS:NZ	4:A:601:HOH:O	2.22	0.73
1:A:57:TRP:CD1	1:A:173:ILE:HG23	2.30	0.67
1:B:268:SER:O	1:B:272:LYS:HE2	1.98	0.64
1:A:263:PHE:C	1:A:263:PHE:CD1	2.77	0.61
1:A:263:PHE:O	1:A:267:PRO:HD3	2.01	0.60
1:B:44:ARG:HD2	1:B:81:LEU:O	2.02	0.59
1:A:261:VAL:CG2	1:A:262:ASP:H	2.14	0.58
1:B:60:THR:O	1:B:60:THR:OG1	2.22	0.56
1:A:227:TRP:CH2	1:A:261:VAL:HG12	2.41	0.56
1:A:60:THR:O	1:A:167:GLN:HG2	2.06	0.55
1:A:268:SER:C	1:A:270:ILE:H	2.15	0.55
1:A:53:GLU:OE1	1:A:54:ARG:HG3	2.08	0.54
1:A:268:SER:C	1:A:270:ILE:N	2.67	0.52
1:A:275:VAL:HG12	1:B:281:MET:HG2	1.91	0.52
1:A:262:ASP:CB	1:A:265:LYS:H	2.23	0.51
1:A:202:ALA:H	1:A:281[A]:MET:HE3	1.76	0.49
1:A:231:ASP:CG	1:A:255:LYS:HG2	2.38	0.49
1:A:173:ILE:H	1:A:173:ILE:HD12	1.79	0.48
1:A:200:LEU:HD23	2:A:401:2GQ:C2	2.43	0.48
1:B:273:LYS:NZ	4:B:581:HOH:O	2.47	0.48
1:A:33:PRO:HG3	1:B:239:TYR:CZ	2.50	0.47
1:A:268:SER:O	1:A:272:LYS:HE2	2.16	0.45
1:A:30:MET:HE2	1:A:30:MET:HB2	1.78	0.45
1:A:281[B]:MET:HE2	1:B:275:VAL:HG11	2.00	0.44
1:A:266:ASP:HA	1:A:267:PRO:HD2	1.24	0.44
1:B:123:GLU:OE1	1:B:164:ARG:NH2	2.46	0.44
1:A:277:CYS:HB2	1:B:281:MET:HE2	1.99	0.44
1:A:261:VAL:CG2	1:A:262:ASP:N	2.77	0.44
1:A:201:PRO:HD3	1:B:199:THR:OG1	2.18	0.43
1:A:266:ASP:O	1:A:269:ALA:HB3	2.18	0.43
1:A:32:ASP:HA	1:A:33:PRO:HD3	1.90	0.43
1:A:60:THR:O	1:A:60:THR:OG1	2.36	0.42
1:A:261:VAL:HG23	1:A:262:ASP:N	2.27	0.42
1:A:239:TYR:CZ	1:B:33:PRO:HG3	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	253/282 (90%)	244 (96%)	9 (4%)	0	100	100
1	B	240/282 (85%)	229 (95%)	11 (5%)	0	100	100
All	All	493/564 (87%)	473 (96%)	20 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	218/247 (88%)	207 (95%)	11 (5%)	22	21
1	B	209/247 (85%)	201 (96%)	8 (4%)	29	30
All	All	427/494 (86%)	408 (96%)	19 (4%)	24	25

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	30	MET
1	A	48	LEU
1	A	60	THR
1	A	96	LEU
1	A	97	LEU
1	A	139	LEU
1	A	170	VAL
1	A	173	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	251	LEU
1	A	261	VAL
1	A	263	PHE
1	B	40	ASP
1	B	48	LEU
1	B	60	THR
1	B	97	LEU
1	B	139	LEU
1	B	249	LEU
1	B	251	LEU
1	B	281	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	126	HIS
1	A	161	GLN
1	A	166	ASN
1	B	126	HIS
1	B	161	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PO4	A	402	-	4,4,4	1.17	0	6,6,6	0.71	0
2	2GQ	A	401	-	17,18,18	0.88	2 (11%)	21,25,25	1.25	2 (9%)
3	PO4	B	403	-	4,4,4	1.03	0	6,6,6	0.79	0
3	PO4	B	401	-	4,4,4	2.34	3 (75%)	6,6,6	1.67	2 (33%)
3	PO4	B	402	-	4,4,4	2.13	2 (50%)	6,6,6	1.56	2 (33%)
3	PO4	A	403	-	4,4,4	2.88	4 (100%)	6,6,6	1.38	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	2GQ	A	401	-	-	0/0/11/11	0/3/3/3

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	403	PO4	P-O2	-3.21	1.45	1.54
3	A	403	PO4	P-O4	-3.15	1.45	1.54
3	A	403	PO4	P-O3	-2.96	1.46	1.54
3	B	401	PO4	P-O2	-2.76	1.46	1.54
3	B	401	PO4	P-O4	-2.72	1.46	1.54
3	B	402	PO4	P-O2	-2.57	1.47	1.54
3	B	401	PO4	P-O3	-2.57	1.47	1.54
3	B	402	PO4	P-O3	-2.41	1.47	1.54
2	A	401	2GQ	C6-C7	-2.14	1.34	1.38
2	A	401	2GQ	C9-N1	-2.10	1.42	1.47
3	A	403	PO4	P-O1	-2.06	1.45	1.50

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	2GQ	C6-C7-N1	-4.55	118.42	125.00
3	B	402	PO4	O4-P-O2	2.75	116.48	107.91

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	403	PO4	O4-P-O3	2.60	116.00	107.91
3	B	401	PO4	O2-P-O1	-2.36	102.61	110.95
3	B	402	PO4	O2-P-O1	-2.26	102.96	110.95
3	B	401	PO4	O4-P-O3	-2.12	101.33	107.91
2	A	401	2GQ	C10-C9-N1	-2.03	100.02	103.44

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	2GQ	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	256/282 (90%)	-0.09	25 (9%) 13 14	10, 27, 59, 97	1 (0%)
1	B	243/282 (86%)	-0.26	7 (2%) 53 56	14, 29, 57, 88	1 (0%)
All	All	499/564 (88%)	-0.17	32 (6%) 25 28	10, 28, 58, 97	2 (0%)

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	263	PHE	5.3
1	A	267	PRO	4.9
1	A	168	HIS	4.8
1	A	266	ASP	4.7
1	A	268	SER	4.5
1	A	261	VAL	4.0
1	A	262	ASP	4.0
1	B	263	PHE	3.8
1	A	146	ASP	3.8
1	B	262	ASP	3.7
1	A	171	PRO	3.5
1	A	28	ARG	3.4
1	A	173	ILE	3.4
1	A	26	TYR	3.3
1	A	170	VAL	3.2
1	A	27	LYS	3.2
1	A	29	GLU	3.1
1	B	30	MET	3.1
1	B	40	ASP	2.9
1	B	166	ASN	2.8
1	A	189	ILE	2.7
1	A	264	CYS	2.7
1	A	271	GLY	2.7
1	A	269	ALA	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	265	LYS	2.5
1	B	187	THR	2.5
1	B	167	GLN	2.5
1	A	174	PRO	2.4
1	A	172	VAL	2.4
1	A	187	THR	2.4
1	A	31	PHE	2.4
1	A	25	PHE	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	PO4	B	402	5/5	0.76	0.26	30,30,30,30	0
2	2GQ	A	401	16/16	0.79	0.16	29,36,43,54	5
3	PO4	B	401	5/5	0.90	0.18	30,30,30,30	0
3	PO4	A	403	5/5	0.92	0.17	30,30,30,30	0
3	PO4	A	402	5/5	0.99	0.03	19,20,23,25	0
3	PO4	B	403	5/5	0.99	0.04	22,22,23,24	0

6.5 Other polymers [i](#)

There are no such residues in this entry.