



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 02:39 AM UTC

PDB ID : 4N7Y / pdb_00004n7y
Title : Crystal structure of 14-3-3zeta in complex with a 8-carbon-linker cyclic peptide derived from ExoS
Authors : Bier, D.; Glas, A.; Hahne, G.; Grossmann, T.; Ottmann, C.
Deposited on : 2013-10-16
Resolution : 2.16 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.16 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2057 (2.16-2.16)
Clashscore	190562	2159 (2.16-2.16)
Ramachandran outliers	187476	2134 (2.16-2.16)
Sidechain outliers	187428	2133 (2.16-2.16)
RSRZ outliers	180081	2059 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	235	12% 77% 18% ..
1	B	235	11% 71% 20% 5% 5%
2	C	11	55% 64% 18% 9% 9%
2	D	11	82% 64% 36%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	2JN	C	422	-	-	X	-
2	MK8	C	425	-	-	X	-
2	2JN	D	422	-	-	X	-
2	MK8	D	425	-	-	X	-

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3774 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 14-3-3 protein zeta/delta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	230	Total	C	N	O	S	0	0	0
			1814	1140	306	358	10			
1	B	223	Total	C	N	O	S	0	0	0
			1751	1102	290	349	10			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	expression tag	UNP P63104
A	-3	ALA	-	expression tag	UNP P63104
A	-2	MET	-	expression tag	UNP P63104
A	-1	GLY	-	expression tag	UNP P63104
A	0	SER	-	expression tag	UNP P63104
B	-4	GLY	-	expression tag	UNP P63104
B	-3	ALA	-	expression tag	UNP P63104
B	-2	MET	-	expression tag	UNP P63104
B	-1	GLY	-	expression tag	UNP P63104
B	0	SER	-	expression tag	UNP P63104

- Molecule 2 is a protein called Exoenzyme S.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	11	Total	C	N	O	0	0	0
			82	53	12	17			
2	D	11	Total	C	N	O	0	0	0
			82	53	12	17			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	422	2JN	LEU	engineered mutation	UNP Q93SQ3
C	425	MK8	ALA	engineered mutation	UNP Q93SQ3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	422	2JN	LEU	engineered mutation	UNP Q93SQ3
D	425	MK8	ALA	engineered mutation	UNP Q93SQ3


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	28	Total O 28 28	0	0
3	B	16	Total O 16 16	0	0
3	D	1	Total O 1 1	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 14-3-3 protein zeta/delta




- Molecule 1: 14-3-3 protein zeta/delta



- Molecule 2: Exoenzyme S



- Molecule 2: Exoenzyme S



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	72.51Å 103.52Å 114.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.94 – 2.16 49.94 – 2.16	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.94-2.16) 100.0 (49.94-2.16)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.59 (at 2.16Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.213 , 0.256 0.227 , 0.265	Depositor DCC
R_{free} test set	2340 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	46.9	Xtriage
Anisotropy	0.329	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 33.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3774	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.26% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MK8, 2JN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.29	4/1839 (0.2%)	1.22	7/2476 (0.3%)
1	B	1.30	3/1773 (0.2%)	1.18	3/2389 (0.1%)
2	C	1.14	0/61	1.21	0/78
2	D	0.94	0/61	1.03	1/78 (1.3%)
All	All	1.29	7/3734 (0.2%)	1.20	11/5021 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	C	0	1
2	D	0	1
All	All	0	2

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	154	GLU	N-CA	6.33	1.54	1.46
1	B	187	LYS	CA-C	6.12	1.60	1.52
1	A	106	ILE	CA-C	6.07	1.59	1.52
1	A	187	LYS	CA-C	6.00	1.60	1.52
1	A	177	PHE	CA-C	5.91	1.60	1.52
1	B	173	ASN	CA-C	5.59	1.60	1.52
1	B	24	ALA	C-O	-5.23	1.18	1.24

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	21	ASP	N-CA-C	-7.06	103.67	111.36
2	D	429	ALA	N-CA-C	-6.87	99.55	109.31
1	A	132	VAL	N-CA-CB	-6.76	103.57	112.36
1	B	105	LEU	N-CA-C	-5.83	104.92	112.68
1	A	104	PHE	N-CA-C	5.74	123.03	110.80
1	A	143	ASP	N-CA-C	5.46	117.24	111.28
1	B	229	THR	CB-CA-C	-5.42	100.63	109.56
1	A	134	ALA	N-CA-CB	-5.29	103.53	111.56
1	A	189	CYS	CA-C-N	-5.06	112.08	120.68
1	A	189	CYS	C-N-CA	-5.06	112.08	120.68
1	B	203	LEU	N-CA-CB	-5.02	101.97	110.50

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	C	425	MK8	Mainchain
2	D	421	GLY	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1814	0	1793	34	0
1	B	1751	0	1695	70	0
2	C	82	0	81	16	0
2	D	82	0	78	21	0
3	A	28	0	0	0	0
3	B	16	0	0	3	0
3	D	1	0	0	0	0
All	All	3774	0	3647	124	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

All (124) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:422:2JN:CAI	2:C:425:MK8:HEA	1.38	1.50
1:B:206:LEU:CD1	1:B:211:TYR:N	1.74	1.50
2:C:422:2JN:H14	2:C:425:MK8:CE	1.05	1.50
1:B:206:LEU:HD11	1:B:211:TYR:N	1.35	1.33
1:B:87:GLU:HG2	1:B:132:VAL:HG11	1.30	1.14
2:C:422:2JN:H14	2:C:425:MK8:CD	1.82	1.09
1:B:206:LEU:HD13	1:B:211:TYR:N	1.50	1.09
2:C:422:2JN:H14	2:C:425:MK8:HE	1.28	1.07
2:C:422:2JN:CAI	2:C:425:MK8:HE	1.81	1.06
1:A:156:SER:HB2	1:A:167:ARG:HG3	1.37	1.05
2:C:422:2JN:CAK	2:C:425:MK8:HEA	1.86	1.05
1:B:52:VAL:HG12	1:B:93:ILE:CG2	1.89	1.02
1:B:52:VAL:HG12	1:B:93:ILE:HG21	1.44	0.99
2:C:422:2JN:CAK	2:C:425:MK8:CE	2.42	0.96
1:B:162:PRO:HG2	1:B:202:GLU:CD	1.91	0.95
1:A:164:HIS:HD2	1:A:166:ILE:H	1.12	0.95
1:B:162:PRO:HG2	1:B:202:GLU:HG3	1.53	0.91
2:C:422:2JN:CAM	2:C:425:MK8:HEA	2.00	0.91
1:B:162:PRO:HG2	1:B:202:GLU:CG	2.02	0.90
1:B:52:VAL:CG1	1:B:93:ILE:CG2	2.51	0.87
1:B:133:ALA:O	1:B:138:LYS:HB2	1.73	0.87
1:B:52:VAL:CG1	1:B:93:ILE:HG21	2.05	0.87
1:B:42:ASN:HB3	2:D:420:GLN:HE21	1.42	0.82
1:B:87:GLU:CG	1:B:132:VAL:HG11	2.06	0.82
1:B:48:TYR:O	1:B:52:VAL:HG22	1.81	0.80
2:C:422:2JN:H14	2:C:425:MK8:HEA	0.85	0.80
1:A:160:MET:HE1	1:A:166:ILE:HB	1.64	0.79
1:A:103:LYS:HD2	1:A:104:PHE:CZ	2.18	0.78
3:B:313:HOH:O	2:D:429:ALA:HB1	1.84	0.77
1:B:162:PRO:HD2	1:B:202:GLU:OE2	1.85	0.77
1:B:84:GLU:O	1:B:88:THR:HG23	1.86	0.76
1:B:87:GLU:CD	1:B:132:VAL:CG1	2.59	0.75
1:B:52:VAL:HG12	1:B:93:ILE:HG22	1.68	0.73
1:B:162:PRO:HG2	1:B:202:GLU:OE2	1.88	0.72
2:D:424:ASP:C	2:D:425:MK8:HNA	1.89	0.72
1:B:42:ASN:HB3	2:D:420:GLN:NE2	2.04	0.72
1:B:190:SER:O	1:B:194:THR:HG23	1.90	0.71
1:B:52:VAL:CG1	1:B:93:ILE:HG22	2.19	0.70
1:B:87:GLU:CD	1:B:132:VAL:HG12	2.17	0.70
3:B:313:HOH:O	2:D:429:ALA:CB	2.40	0.70
1:A:156:SER:CB	1:A:167:ARG:HG3	2.20	0.70
2:D:423:LEU:HD23	2:D:428:LEU:HD12	1.72	0.70

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:49:LYS:HG3	2:D:427:ASP:OD1	1.91	0.70
1:A:164:HIS:CD2	1:A:166:ILE:H	2.04	0.69
1:B:46:VAL:HG23	2:D:425:MK8:HB1A	1.74	0.68
1:A:134:ALA:HA	1:A:138:LYS:HB2	1.75	0.68
1:B:213:ASP:CG	2:D:422:2JN:H9	2.18	0.68
1:A:133:ALA:O	1:A:138:LYS:HG3	1.93	0.67
1:A:206:LEU:HG	1:A:211:TYR:CB	2.25	0.67
1:A:71:GLY:HA2	1:A:72:ALA:HB3	1.78	0.66
1:A:3:LYS:HE2	1:A:40:GLU:OE1	1.96	0.66
2:D:420:GLN:HG3	2:D:425:MK8:HB1	1.79	0.64
2:C:422:2JN:CAM	2:C:425:MK8:CE	2.74	0.64
1:B:213:ASP:HB3	2:D:422:2JN:H9	1.79	0.63
1:A:164:HIS:NE2	1:A:166:ILE:HG13	2.13	0.63
1:A:103:LYS:HD2	1:A:104:PHE:CE2	2.34	0.62
1:B:46:VAL:HG23	2:D:425:MK8:CB1	2.31	0.60
1:B:162:PRO:CD	1:B:202:GLU:OE2	2.49	0.60
1:B:2:ASP:OD1	1:B:5:GLU:HG2	2.01	0.60
1:B:52:VAL:HG11	1:B:125:TYR:HE2	1.67	0.60
1:B:42:ASN:CB	2:D:420:GLN:HE21	2.14	0.59
1:A:59:TRP:CE2	1:A:132:VAL:HG22	2.37	0.59
1:B:87:GLU:CG	1:B:132:VAL:CG1	2.81	0.58
1:B:162:PRO:CG	1:B:202:GLU:OE2	2.51	0.58
1:A:71:GLY:HA2	1:A:72:ALA:C	2.28	0.57
1:B:213:ASP:CB	2:D:422:2JN:H9	2.34	0.57
1:A:164:HIS:CD2	1:A:166:ILE:HG13	2.40	0.56
1:B:103:LYS:HB3	1:B:104:PHE:CE2	2.41	0.56
1:A:206:LEU:HG	1:A:211:TYR:HB3	1.88	0.56
1:A:65:ILE:O	1:A:69:THR:HG23	2.06	0.55
1:B:216:LEU:HD12	1:B:216:LEU:O	2.05	0.55
1:B:59:TRP:CE2	1:B:132:VAL:HG13	2.41	0.54
1:B:59:TRP:NE1	1:B:132:VAL:HG13	2.23	0.54
1:B:162:PRO:CG	1:B:202:GLU:HG3	2.32	0.54
1:B:137:ASP:O	1:B:141:ILE:HG12	2.08	0.54
1:B:112:ALA:CB	1:B:159:GLU:HG2	2.39	0.53
1:B:163:THR:HB	1:B:206:LEU:HD23	1.91	0.53
1:B:211:TYR:N	1:B:211:TYR:CD1	2.75	0.52
1:B:206:LEU:CD1	1:B:211:TYR:CA	2.79	0.52
1:A:66:GLU:O	1:A:66:GLU:HG2	2.10	0.51
1:B:211:TYR:N	1:B:211:TYR:HD1	2.08	0.51
1:A:160:MET:CE	1:A:166:ILE:HB	2.39	0.50
1:A:71:GLY:CA	1:A:72:ALA:C	2.86	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:160:MET:HE3	1:B:164:HIS:CG	2.48	0.49
1:A:2:ASP:O	1:A:3:LYS:C	2.56	0.49
1:B:103:LYS:CB	1:B:104:PHE:CE2	2.97	0.48
1:B:160:MET:HE2	1:B:167:ARG:HB2	1.94	0.48
1:A:172:LEU:HD23	2:C:428:LEU:HD22	1.96	0.47
1:B:41:ARG:HD2	1:B:117:PHE:CD1	2.50	0.47
1:B:133:ALA:O	1:B:138:LYS:CB	2.56	0.47
2:C:422:2JN:H14	2:C:425:MK8:HD	1.84	0.47
1:A:14:GLU:OE2	2:C:420:GLN:NE2	2.48	0.46
1:B:168:LEU:HD12	1:B:168:LEU:HA	1.80	0.46
1:B:52:VAL:HG13	1:B:93:ILE:HG21	1.95	0.46
1:B:59:TRP:NE1	1:B:132:VAL:CG1	2.79	0.46
1:A:14:GLU:HB2	1:A:22:MET:HE3	1.97	0.45
2:C:422:2JN:H10	2:C:425:MK8:HEA	1.92	0.45
1:A:75:LYS:HE3	1:A:75:LYS:HB2	1.75	0.45
1:B:87:GLU:OE2	1:B:132:VAL:CG1	2.64	0.45
1:A:158:LYS:HE3	1:A:158:LYS:HB2	1.76	0.44
1:B:213:ASP:HB3	2:D:422:2JN:CAM	2.48	0.44
1:A:160:MET:HE1	1:A:166:ILE:CB	2.43	0.44
1:B:103:LYS:HB2	1:B:104:PHE:CD2	2.52	0.44
2:C:422:2JN:CAK	2:C:425:MK8:HE	2.28	0.44
1:B:213:ASP:CG	2:D:422:2JN:CAM	2.87	0.43
1:B:85:LYS:HG2	3:B:316:HOH:O	2.17	0.43
2:C:422:2JN:H12	2:C:425:MK8:CE	2.42	0.43
1:A:206:LEU:HG	1:A:211:TYR:HB2	1.98	0.43
1:B:162:PRO:O	1:B:168:LEU:HD13	2.18	0.43
2:D:422:2JN:H10	2:D:425:MK8:HEB	1.73	0.43
1:B:42:ASN:OD1	2:D:425:MK8:HG	2.19	0.42
1:A:90:LEU:C	1:A:90:LEU:HD23	2.45	0.42
1:A:69:THR:OG1	1:A:76:GLN:HG3	2.19	0.42
1:A:133:ALA:O	1:A:138:LYS:CG	2.65	0.42
1:B:46:VAL:HG22	2:D:424:ASP:HB3	2.02	0.42
1:B:59:TRP:CZ2	1:B:132:VAL:HG13	2.55	0.42
1:B:12:LEU:HD23	1:B:12:LEU:HA	1.86	0.42
1:A:206:LEU:CD2	1:A:211:TYR:HB2	2.50	0.41
1:B:182:LEU:HD12	1:B:182:LEU:HA	1.77	0.41
1:B:103:LYS:CB	1:B:104:PHE:CD2	3.03	0.41
1:B:153:PHE:HA	1:B:170:LEU:HD21	2.03	0.41
1:B:120:LYS:NZ	2:D:426:LEU:O	2.44	0.40
1:A:134:ALA:HA	1:A:135:GLY:HA2	1.76	0.40
1:B:169:GLY:HA3	2:D:426:LEU:HD13	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	228/235 (97%)	220 (96%)	8 (4%)	0	100	100
1	B	215/235 (92%)	213 (99%)	2 (1%)	0	100	100
2	C	7/11 (64%)	6 (86%)	1 (14%)	0	100	100
2	D	7/11 (64%)	6 (86%)	1 (14%)	0	100	100
All	All	457/492 (93%)	445 (97%)	12 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	192/202 (95%)	176 (92%)	16 (8%)	10	6
1	B	183/202 (91%)	165 (90%)	18 (10%)	7	4
2	C	7/7 (100%)	7 (100%)	0	100	100
2	D	7/7 (100%)	5 (71%)	2 (29%)	0	0
All	All	389/418 (93%)	353 (91%)	36 (9%)	8	4

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	LYS
1	A	9	LYS
1	A	67	GLN
1	A	100	LEU
1	A	103	LYS
1	A	113	GLU
1	A	132	VAL
1	A	139	LYS
1	A	143	ASP
1	A	147	GLN
1	A	168	LEU
1	A	182	LEU
1	A	203	LEU
1	A	204	ASP
1	A	210	SER
1	A	229	THR
1	B	5	GLU
1	B	38	ASN
1	B	49	LYS
1	B	76	GLN
1	B	88	THR
1	B	100	LEU
1	B	102	GLU
1	B	115	LYS
1	B	132	VAL
1	B	159	GLU
1	B	168	LEU
1	B	182	LEU
1	B	183	ASN
1	B	184	SER
1	B	194	THR
1	B	202	GLU
1	B	206	LEU
1	B	211	TYR
2	D	420	GLN
2	D	430	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	146	GLN
1	A	164	HIS
1	A	224	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	50	ASN
1	B	76	GLN
1	B	146	GLN
1	B	150	GLN
2	C	420	GLN
2	D	420	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	MK8	C	425	2	5,8,9	3.93	1 (20%)	6,10,12	2.98	4 (66%)
2	MK8	D	425	2	5,8,9	3.00	1 (20%)	6,10,12	1.82	1 (16%)
2	2JN	C	422	2	5,8,9	0.85	0	6,10,12	1.81	1 (16%)
2	2JN	D	422	2	5,8,9	0.84	0	6,10,12	1.80	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MK8	C	425	2	-	2/6/8/11	-
2	MK8	D	425	2	-	1/6/8/11	-
2	2JN	C	422	2	-	4/6/8/11	-
2	2JN	D	422	2	-	0/6/8/11	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	425	MK8	CB-CA	8.40	1.64	1.55
2	D	425	MK8	CB-CA	6.53	1.62	1.55

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	425	MK8	CG-CB-CA	-4.40	108.90	115.41
2	C	422	2JN	CAM-CAO-CA	-4.02	109.47	115.41
2	D	422	2JN	CAM-CAO-CA	-4.00	109.50	115.41
2	C	425	MK8	CB1-CA-CB	3.82	117.26	110.97
2	D	425	MK8	CG-CB-CA	3.69	120.87	115.41
2	C	425	MK8	CB-CA-C	3.44	117.33	108.47
2	C	425	MK8	CB1-CA-C	2.23	118.67	109.75

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	422	2JN	O-C-CA-CAA
2	C	425	MK8	CE-CD-CG-CB
2	C	422	2JN	CAA-CA-CAO-CAM
2	C	422	2JN	CAK-CAM-CAO-CA
2	D	425	MK8	CE-CD-CG-CB
2	C	422	2JN	CAI-CAK-CAM-CAO
2	C	425	MK8	CB1-CA-CB-CG

There are no ring outliers.

4 monomers are involved in 25 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	425	MK8	14	0
2	D	425	MK8	6	0
2	C	422	2JN	14	0
2	D	422	2JN	6	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	D	1
2	C	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	D	424:ASP	C	425:MK8	N	1.68
1	C	424:ASP	C	425:MK8	N	1.65

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	230/235 (97%)	0.69	29 (12%) 8 8	25, 45, 92, 133	0
1	B	223/235 (94%)	0.65	27 (12%) 8 9	23, 45, 80, 114	0
2	C	9/11 (81%)	2.71	6 (66%) 0 0	45, 63, 90, 94	0
2	D	9/11 (81%)	5.10	9 (100%) 0 0	64, 79, 116, 129	0
All	All	471/492 (95%)	0.79	71 (15%) 5 6	23, 45, 92, 133	0

All (71) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	MET	8.3
1	A	203	LEU	8.1
1	A	134	ALA	7.8
2	D	423	LEU	7.4
1	B	134	ALA	7.0
1	B	104	PHE	6.9
1	B	211	TYR	6.9
1	A	72	ALA	6.6
2	D	428	LEU	6.5
1	B	136	ASP	6.2
2	D	430	SER	6.1
1	A	205	THR	5.8
1	A	202	GLU	5.7
2	D	429	ALA	5.4
1	B	135	GLY	5.4
2	D	426	LEU	5.2
1	A	70	GLU	5.2
1	B	202	GLU	5.1
2	D	420	GLN	5.0
1	A	135	GLY	5.0
1	B	133	ALA	5.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	71	GLY	4.7
1	B	73	GLU	4.6
1	A	230	SER	4.6
1	B	203	LEU	4.5
2	C	423	LEU	4.5
1	B	1	MET	4.5
1	B	137	ASP	4.5
1	A	74	LYS	4.4
2	C	430	SER	4.4
1	B	207	SER	4.4
1	B	52	VAL	4.3
1	A	208	GLU	4.3
1	A	204	ASP	4.2
1	A	133	ALA	4.2
2	D	421	GLY	4.2
1	B	132	VAL	4.1
1	A	2	ASP	4.1
1	B	75	LYS	4.1
1	A	210	SER	3.9
1	A	209	GLU	3.8
1	A	206	LEU	3.8
2	C	420	GLN	3.7
1	A	211	TYR	3.6
1	A	75	LYS	3.6
1	A	69	THR	3.5
1	B	230	SER	3.5
1	B	70	GLU	3.5
1	B	102	GLU	3.4
2	D	424	ASP	3.2
1	B	74	LYS	3.2
2	D	427	ASP	3.0
1	B	212	LYS	3.0
1	B	206	LEU	3.0
2	C	426	LEU	2.9
1	A	73	GLU	2.9
2	C	421	GLY	2.9
1	B	69	THR	2.8
1	A	166	ILE	2.6
1	A	67	GLN	2.5
1	B	80	ARG	2.4
2	C	429	ALA	2.4
1	A	68	LYS	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	68	LYS	2.2
1	B	103	LYS	2.2
1	A	104	PHE	2.1
1	A	207	SER	2.1
1	A	215	THR	2.1
1	A	139	LYS	2.1
1	B	41	ARG	2.0
1	B	229	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MK8	D	425	9/10	0.81	0.27	59,74,103,108	0
2	MK8	C	425	9/10	0.87	0.19	41,51,62,66	0
2	2JN	C	422	9/10	0.90	0.16	53,58,67,72	0
2	2JN	D	422	9/10	0.90	0.20	69,88,96,97	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.