



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 07:11 AM UTC

PDB ID : 6N71 / pdb_00006n71
Title : Crystal Structure of ATPase delta1-79 Spa47 R191E
Authors : Morales, Y.; Olsen, K.J.; Johnson, S.J.; Demler, H.J.; Dickenson, N.E.
Deposited on : 2018-11-27
Resolution : 2.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1190 (2.46-2.46)
Clashscore	190562	1229 (2.46-2.46)
Ramachandran outliers	187476	1218 (2.46-2.46)
Sidechain outliers	187428	1218 (2.46-2.46)
RSRZ outliers	180081	1190 (2.46-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	351	8% 76% 18% 6%
1	B	351	7% 83% 12% • 5%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5243 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable ATP synthase SpaL/MxiB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	331	Total	C	N	O	S	0	0	0
			2585	1643	438	495	9			
1	B	333	Total	C	N	O	S	0	0	0
			2605	1653	440	503	9			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	191	GLU	ARG	engineered mutation	UNP P0A1C1
B	191	GLU	ARG	engineered mutation	UNP P0A1C1


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	27	Total	O	0	0
			27	27		
2	B	26	Total	O	0	0
			26	26		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Probable ATP synthase SpaL/MxiB



- Molecule 1: Probable ATP synthase SpaL/MxiB



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	44.02Å 152.74Å 54.47Å 90.00° 109.92° 90.00°	Depositor
Resolution (Å)	48.55 – 2.45 48.55 – 2.45	Depositor EDS
% Data completeness (in resolution range)	93.3 (48.55-2.45) 90.2 (48.55-2.45)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.93 (at 2.42Å)	Xtriage
Refinement program	PHENIX 1.13_2998	Depositor
R, R_{free}	0.205 , 0.247 0.207 , 0.249	Depositor DCC
R_{free} test set	1170 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	44.3	Xtriage
Anisotropy	0.298	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 43.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5243	wwPDB-VP
Average B, all atoms (Å ²)	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.12	0/2626	0.33	0/3544
1	B	0.12	0/2647	0.31	0/3575
All	All	0.12	0/5273	0.32	0/7119

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	187	GLY	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2585	0	2601	41	0
1	B	2605	0	2611	29	0
2	A	27	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	26	0	0	0	0
All	All	5243	0	5212	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:307:GLU:OE2	1:B:313:ASP:HB3	1.17	1.28
1:B:307:GLU:OE2	1:B:313:ASP:CB	2.02	1.06
1:B:307:GLU:CD	1:B:313:ASP:HB3	2.08	0.79
1:A:186:ILE:HD12	1:A:254:TYR:HB2	1.72	0.72
1:A:411:ASP:HB3	1:A:414:LEU:HG	1.73	0.71
1:B:313:ASP:HB2	1:B:314:PRO:HD2	1.75	0.68
1:B:304:VAL:HG21	1:B:319:VAL:HG21	1.76	0.67
1:A:429:ILE:HG13	1:A:430:ARG:HD3	1.77	0.66
1:B:153:GLN:HG3	1:B:324:ASP:HB2	1.78	0.66
1:A:238:ARG:HD3	1:A:298:ILE:HG13	1.79	0.63
1:A:358:GLU:H	1:A:358:GLU:CD	2.07	0.63
1:A:332:ASN:HA	1:A:335:GLN:NE2	2.16	0.61
1:B:314:PRO:O	1:B:318:GLU:HG2	2.00	0.61
1:A:86:GLN:NE2	1:A:100:GLY:O	2.35	0.60
1:B:419:GLU:N	1:B:419:GLU:OE2	2.33	0.60
1:A:133:GLU:HG3	1:A:134:LYS:HD2	1.86	0.57
1:B:258:LEU:HD21	1:B:281:SER:OG	2.05	0.57
1:A:83:LEU:N	2:A:502:HOH:O	2.37	0.57
1:A:186:ILE:CD1	1:A:254:TYR:HB2	2.36	0.55
1:A:235:GLU:OE2	1:A:288:ARG:NE	2.36	0.54
1:A:153:GLN:HG3	1:A:324:ASP:HB2	1.90	0.53
1:A:380:ILE:HG23	1:A:385:GLU:HG3	1.91	0.53
1:A:283:PRO:HA	1:A:286:LEU:HD11	1.89	0.53
1:B:87:VAL:HG21	1:B:117:ARG:HG2	1.91	0.52
1:A:130:ALA:HB3	1:A:291:LYS:HG3	1.92	0.51
1:A:282:LEU:O	1:A:286:LEU:HG	2.10	0.51
1:B:308:ASP:O	1:B:310:ASP:N	2.44	0.50
1:A:186:ILE:HG22	1:A:226:ALA:HB3	1.93	0.49
1:A:378:ARG:NH2	1:A:405:GLU:OE2	2.45	0.49
1:A:154:ARG:HD2	1:A:286:LEU:O	2.13	0.48
1:A:430:ARG:HD3	1:A:430:ARG:N	2.28	0.48
1:B:159:ALA:HB2	1:B:329:LEU:HB2	1.97	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:252:THR:HG21	1:A:305:LEU:H	1.81	0.46
1:A:88:GLY:HA2	1:A:236:PHE:CE2	2.50	0.46
1:A:146:LEU:HD11	1:A:367:PHE:CD1	2.50	0.46
1:B:422:MET:HA	1:B:425:ILE:HG22	1.98	0.45
1:A:178:ALA:HA	1:A:243:LYS:HD3	1.97	0.45
1:B:193:VAL:O	1:B:197:VAL:HG23	2.16	0.45
1:A:159:ALA:HB2	1:A:329:LEU:HB2	1.99	0.45
1:A:249:ASP:HA	1:A:250:SER:HA	1.58	0.45
1:B:154:ARG:HD2	1:B:286:LEU:O	2.17	0.44
1:A:332:ASN:HA	1:A:335:GLN:HE21	1.79	0.44
1:B:165:LYS:NZ	1:B:249:ASP:OD1	2.51	0.44
1:B:333:LEU:HD22	1:B:338:GLN:OE1	2.17	0.44
1:A:97:ASN:OD1	1:A:101:GLU:HG3	2.18	0.44
1:B:282:LEU:HB3	1:B:283:PRO:HD3	2.00	0.43
1:A:124:PRO:O	1:A:129:ARG:NH1	2.36	0.43
1:A:304:VAL:HG11	1:A:319:VAL:HG21	2.00	0.43
1:B:188:GLU:O	1:B:215:THR:HA	2.18	0.43
1:B:249:ASP:HA	1:B:250:SER:HA	1.52	0.43
1:A:340:PRO:HB2	1:A:342:ILE:HG13	2.00	0.43
1:B:197:VAL:O	1:B:201:LYS:HE3	2.19	0.43
1:A:335:GLN:HE21	1:A:335:GLN:HB2	1.62	0.42
1:B:318:GLU:O	1:B:322:ILE:HD12	2.19	0.42
1:B:191:GLU:HG3	1:B:194:THR:OG1	2.19	0.42
1:A:248:ILE:HB	1:A:302:TYR:HD1	1.83	0.42
1:B:340:PRO:HB2	1:B:342:ILE:HG13	2.01	0.42
1:B:339:PHE:CE2	1:B:412:TYR:HB3	2.55	0.42
1:B:339:PHE:HE2	1:B:412:TYR:HB3	1.84	0.42
1:A:93:GLY:HA2	1:A:106:PHE:CE1	2.54	0.42
1:B:117:ARG:HD2	1:B:117:ARG:HA	1.86	0.42
1:B:425:ILE:O	1:B:428:THR:HG22	2.20	0.41
1:A:163:CYS:SG	1:A:331:ARG:HB2	2.59	0.41
1:A:187:GLY:HA2	1:A:253:ARG:HD3	2.02	0.41
1:A:282:LEU:HB3	1:A:283:PRO:HD3	2.02	0.41
1:A:154:ARG:NH2	1:A:324:ASP:OD1	2.54	0.41
1:A:318:GLU:O	1:A:322:ILE:HG13	2.21	0.41
1:B:280:ASP:O	1:B:283:PRO:HD2	2.21	0.41
1:A:169:MET:HE1	1:A:249:ASP:HB2	2.03	0.40
1:A:251:LEU:HD23	1:A:251:LEU:HA	1.94	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	325/351 (93%)	317 (98%)	8 (2%)	0	100	100
1	B	329/351 (94%)	319 (97%)	7 (2%)	3 (1%)	14	17
All	All	654/702 (93%)	636 (97%)	15 (2%)	3 (0%)	24	32

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	310	ASP
1	B	309	ASP
1	B	307	GLU

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	277/293 (94%)	277 (100%)	0	100	100
1	B	280/293 (96%)	280 (100%)	0	100	100
All	All	557/586 (95%)	557 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	202	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	335	GLN
1	B	111	ASN
1	B	202	ASN
1	B	332	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	331/351 (94%)	0.69	28 (8%) 16 14	28, 56, 102, 121	0
1	B	333/351 (94%)	0.69	26 (7%) 19 16	31, 52, 101, 167	0
All	All	664/702 (94%)	0.69	54 (8%) 18 16	28, 54, 102, 167	0

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	286	LEU	5.5
1	A	279	PHE	4.7
1	B	322	ILE	4.1
1	B	282	LEU	3.9
1	A	266	GLY	3.8
1	B	319	VAL	3.4
1	A	83	LEU	3.4
1	B	121	ASN	3.2
1	A	187	GLY	3.2
1	B	213	TYR	3.2
1	B	264	ALA	3.2
1	B	280	ASP	3.0
1	B	311	PHE	3.0
1	A	164	GLY	3.0
1	B	312	ALA	2.8
1	B	385	GLU	2.6
1	A	186	ILE	2.5
1	B	95	VAL	2.5
1	A	264	ALA	2.5
1	B	314	PRO	2.5
1	A	316	ALA	2.5
1	B	430	ARG	2.4
1	B	217	ASP	2.4
1	B	305	LEU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	315	LEU	2.3
1	A	204	GLU	2.3
1	B	83	LEU	2.3
1	B	163	CYS	2.3
1	A	371	LEU	2.2
1	B	111	ASN	2.2
1	A	251	LEU	2.2
1	A	280	ASP	2.2
1	A	163	CYS	2.2
1	A	315	LEU	2.2
1	A	321	SER	2.2
1	A	212	VAL	2.2
1	A	304	VAL	2.2
1	A	319	VAL	2.2
1	A	263	LEU	2.2
1	A	375	GLU	2.2
1	B	199	TYR	2.1
1	B	281	SER	2.1
1	B	262	ALA	2.1
1	B	263	LEU	2.1
1	B	189	ARG	2.1
1	A	123	PRO	2.1
1	B	254	TYR	2.1
1	A	338	GLN	2.1
1	A	122	ALA	2.1
1	A	282	LEU	2.1
1	A	101	GLU	2.1
1	A	328	TYR	2.1
1	B	310	ASP	2.1
1	A	131	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.