



Full wwPDB NMR Structure Validation Report ⓘ

Mar 19, 2026 – 08:24 PM UTC

PDB ID : 2N80 / pdb_00002n80
BMRB ID : 25829
Title : p75NTR DD:RhoGDI
Authors : Lin, Z.; Ibanez, C.F.
Deposited on : 2015-09-30

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 78%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	94	
2	B	174	

2 Ensemble composition and analysis

This entry contains 10 models. Model 10 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:334-A:421, B:68-B:203 (224)	0.70	10
2	B:34-B:47 (14)	1.47	10

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 1 single-model cluster was found.

Cluster number	Models
1	3, 4, 5, 6, 7, 9, 10
2	2, 8
Single-model clusters	1

3 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 4210 atoms, of which 2098 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Tumor necrosis factor receptor superfamily member 16.

Mol	Chain	Residues	Atoms						Trace
1	A	94	Total	C	H	N	O	S	0
			1413	444	699	126	142	2	

- Molecule 2 is a protein called Rho GDP-dissociation inhibitor 1.


Mol	Chain	Residues	Atoms						Trace
2	B	174	Total	C	H	N	O	S	0
			2797	887	1399	235	272	4	

4 Residue-property plots


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Tumor necrosis factor receptor superfamily member 16



- Molecule 2: Rho GDP-dissociation inhibitor 1




4.2 Scores per residue for each member of the ensemble


Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

- Molecule 1: Tumor necrosis factor receptor superfamily member 16



- Molecule 2: Rho GDP-dissociation inhibitor 1





4.2.2 Score per residue for model 2


- Molecule 1: Tumor necrosis factor receptor superfamily member 16

Chain A: 77% 17% 6%



- Molecule 2: Rho GDP-dissociation inhibitor 1

Chain B: 71% 14% 14%



4.2.3 Score per residue for model 3


- Molecule 1: Tumor necrosis factor receptor superfamily member 16

Chain A: 78% 16% 6%



- Molecule 2: Rho GDP-dissociation inhibitor 1

Chain B: 74% 12% 14%



4.2.4 Score per residue for model 4


- Molecule 1: Tumor necrosis factor receptor superfamily member 16

Chain A: 79% 14% 6%



- Molecule 2: Rho GDP-dissociation inhibitor 1

Chain B: 71% 14% 14%



4.2.5 Score per residue for model 5


- Molecule 1: Tumor necrosis factor receptor superfamily member 16

Chain A: 79% 15% 6%



- Molecule 2: Rho GDP-dissociation inhibitor 1


Chain B: 73% 13% 14%



4.2.6 Score per residue for model 6


- Molecule 1: Tumor necrosis factor receptor superfamily member 16


Chain A: 71% 22% 6%



- Molecule 2: Rho GDP-dissociation inhibitor 1


Chain B: 71% 16% 14%






4.2.7 Score per residue for model 7

- Molecule 1: Tumor necrosis factor receptor superfamily member 16




- Molecule 2: Rho GDP-dissociation inhibitor 1




4.2.8 Score per residue for model 8

- Molecule 1: Tumor necrosis factor receptor superfamily member 16




- Molecule 2: Rho GDP-dissociation inhibitor 1



4.2.9 Score per residue for model 9

- Molecule 1: Tumor necrosis factor receptor superfamily member 16





• Molecule 2: Rho GDP-dissociation inhibitor 1

Chain B: 72% 14% 14%



4.2.10 Score per residue for model 10 (medoid)


• Molecule 1: Tumor necrosis factor receptor superfamily member 16

Chain A: 74% 19% 6%



• Molecule 2: Rho GDP-dissociation inhibitor 1

Chain B: 70% 17% 14%



5 Refinement protocol and experimental data overview

The models were refined using the following method: *molecular dynamics*.

Of the 100 calculated structures, 10 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	
Amber	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	2
Total number of shifts	2849
Number of shifts mapped to atoms	2848
Number of unparsed shifts	0
Number of shifts with mapping errors	1
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	78%

6 Model quality

6.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.90±0.04	0±0/689 (0.0± 0.0%)	1.40±0.03	0±0/938 (0.0± 0.0%)
2	B	0.82±0.03	0±0/1239 (0.0± 0.0%)	1.22±0.04	0±0/1671 (0.0± 0.0%)
All	All	0.85	0/19280 (0.0%)	1.29	3/26090 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	0.1±0.3
2	B	0.0±0.0	0.4±0.5
All	All	0	5

There are no bond-length outliers.

All unique angle outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
2	B	100	GLN	OE1-CD-NE2	-5.69	116.91	122.60	10	3

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
2	B	175	TYR	Sidechain	4
1	A	384	ARG	Sidechain	1

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	675	659	658	0±0
2	B	1213	1202	1202	0±0
All	All	18880	18610	18600	7

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:346:GLU:CD	2:B:88:LEU:HD21	0.55	2.26	4	2
1:A:402:LEU:HD22	1:A:414:VAL:HG13	0.49	1.84	9	3
2:B:86:LEU:HD23	2:B:102:PHE:CD1	0.42	2.49	7	1
2:B:75:LEU:HD23	2:B:75:LEU:C	0.42	2.39	9	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	87/94 (93%)	84±1 (96±1%)	2±1 (3±1%)	1±1 (1±1%)	12	60
2	B	150/174 (86%)	139±2 (93±1%)	9±2 (6±2%)	2±1 (1±1%)	15	65
All	All	2370/2680 (88%)	2227 (94%)	117 (5%)	26 (1%)	14	63

All 8 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	396	ALA	7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
2	B	120	ARG	5
2	B	93	ASP	4
2	B	203	LYS	3
1	A	355	ALA	3
2	B	35	ILE	2
2	B	38	ILE	1
1	A	354	SER	1

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	71/76 (93%)	58±2 (82±3%)	13±2 (18±3%)	3	35
2	B	136/155 (88%)	113±3 (83±2%)	23±3 (17±2%)	4	38
All	All	2070/2310 (90%)	1707 (82%)	363 (18%)	4	37

All 99 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	350	LEU	10
1	A	386	LEU	10
2	B	123	VAL	10
2	B	174	SER	10
1	A	401	LEU	9
1	A	375	SER	8
1	A	398	LEU	8
1	A	421	SER	8
1	A	346	GLU	8
1	A	343	LYS	7
1	A	420	GLU	7
2	B	41	LEU	7
2	B	43	LYS	7
2	B	108	VAL	7
2	B	119	ASN	7
2	B	120	ARG	7
2	B	139	ILE	7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
2	B	203	LYS	7
1	A	392	THR	6
1	A	399	ASP	6
2	B	47	SER	6
2	B	141	LYS	6
2	B	170	LEU	6
2	B	179	SER	6
2	B	181	PHE	6
1	A	374	ASP	6
1	A	413	LEU	6
2	B	70	VAL	5
2	B	159	LEU	5
2	B	167	LYS	5
2	B	184	ASP	5
2	B	185	ASP	5
2	B	142	THR	5
2	B	96	SER	5
2	B	176	SER	5
2	B	36	GLN	4
2	B	80	SER	4
2	B	135	LYS	4
2	B	138	LYS	4
2	B	200	LYS	4
1	A	416	SER	4
2	B	89	ASP	4
2	B	126	MET	4
2	B	157	GLU	4
1	A	412	ASP	3
2	B	117	ARG	3
1	A	362	LEU	3
2	B	37	GLU	3
1	A	406	ARG	3
1	A	371	GLU	3
2	B	152	ARG	3
2	B	155	GLU	3
2	B	99	LYS	3
2	B	76	THR	2
2	B	95	GLU	2
2	B	145	MET	2
2	B	154	GLU	2
2	B	177	ILE	2
1	A	378	HIS	2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
2	B	44	ASP	2
1	A	372	HIS	2
2	B	34	SER	2
2	B	38	ILE	2
2	B	169	MET	2
2	B	39	GLN	2
2	B	134	ARG	2
2	B	140	ASP	2
2	B	46	GLU	2
2	B	186	LYS	2
2	B	75	LEU	1
2	B	77	LEU	1
2	B	91	THR	1
1	A	336	TYR	1
1	A	366	LEU	1
1	A	405	LEU	1
1	A	414	VAL	1
2	B	40	GLU	1
2	B	100	GLN	1
1	A	368	TYR	1
2	B	180	ARG	1
2	B	199	LYS	1
2	B	93	ASP	1
2	B	115	SER	1
2	B	132	THR	1
1	A	360	ARG	1
1	A	384	ARG	1
1	A	387	LEU	1
1	A	397	THR	1
2	B	106	GLU	1
2	B	127	LYS	1
2	B	42	ASP	1
1	A	389	SER	1
2	B	109	GLU	1
1	A	358	THR	1
2	B	86	LEU	1
2	B	102	PHE	1
2	B	188	ASP	1
1	A	345	GLU	1
2	B	172	ARG	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 78% for the well-defined parts and 78% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1011
Number of shifts mapped to atoms	1011
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	6

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	94	0.21 ± 0.09	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	89	1.07 ± 0.06	Should be checked
$^{13}\text{C}'$	0	—	None (insufficient data)
^{15}N	89	0.57 ± 0.21	Should be applied

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 30%, i.e. 957 atoms were assigned a chemical shift out of a possible 3229. 0 out of 40 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	348/1185 (29%)	176/481 (37%)	88/476 (18%)	84/228 (37%)
Sidechain	567/1808 (31%)	385/1171 (33%)	178/570 (31%)	4/67 (6%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	42/236 (18%)	21/116 (18%)	19/110 (17%)	2/10 (20%)
Overall	957/3229 (30%)	582/1768 (33%)	285/1156 (25%)	90/305 (30%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 28%, i.e. 1010 atoms were assigned a chemical shift out of a possible 3630. 0 out of 47 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	370/1332 (28%)	187/540 (35%)	94/536 (18%)	89/256 (35%)
Sidechain	598/2053 (29%)	404/1331 (30%)	190/644 (30%)	4/78 (5%)
Aromatic	42/245 (17%)	21/120 (18%)	19/115 (17%)	2/10 (20%)
Overall	1010/3630 (28%)	612/1991 (31%)	303/1295 (23%)	95/344 (28%)

7.1.4 Statistically unusual chemical shifts ⓘ


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	396	ALA	HB1	-0.20	0.14 – 2.58	-6.4
1	A	396	ALA	HB2	-0.20	0.14 – 2.58	-6.4
1	A	396	ALA	HB3	-0.20	0.14 – 2.58	-6.4
1	A	360	ARG	HG2	-0.08	0.26 – 2.87	-6.3
1	A	382	PRO	HB3	-0.14	0.25 – 3.76	-6.1
1	A	360	ARG	HG3	-0.02	0.15 – 2.94	-5.6

7.1.5 Random Coil Index (RCI) plots ⓘ

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:



7.2 Chemical shift list 2

File name: working_cs.cif

Chemical shift list name: *assigned_chem_shift_list_2*

7.2.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1838
Number of shifts mapped to atoms	1837
Number of unparsed shifts	0
Number of shifts with mapping errors	1
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	1

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

- No matching atom found in the structure. All 1 occurrences are reported below.

List ID	Chain	Res	Type	Atom	Shift Data		
					Value	Uncertainty	Ambiguity
2	B	31	ALA	H	8.464	.	1

7.2.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	171	0.69 ± 0.11	Should be applied
$^{13}\text{C}_\beta$	160	0.54 ± 0.11	Should be applied
$^{13}\text{C}'$	0	—	None (insufficient data)
^{15}N	153	-0.49 ± 0.35	None needed (< 0.5 ppm)

7.2.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 48%, i.e. 1564 atoms were assigned a chemical shift out of a possible 3229. 0 out of 40 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	563/1185 (48%)	286/481 (59%)	147/476 (31%)	130/228 (57%)
Sidechain	929/1808 (51%)	611/1171 (52%)	314/570 (55%)	4/67 (6%)
Aromatic	72/236 (31%)	36/116 (31%)	33/110 (30%)	3/10 (30%)
Overall	1564/3229 (48%)	933/1768 (53%)	494/1156 (43%)	137/305 (45%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 51%, i.e. 1837 atoms were assigned a chemical shift out of a possible 3630. 0 out of 47 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	658/1332 (49%)	334/540 (62%)	171/536 (32%)	153/256 (60%)
Sidechain	1103/2053 (54%)	728/1331 (55%)	369/644 (57%)	6/78 (8%)
Aromatic	76/245 (31%)	38/120 (32%)	35/115 (30%)	3/10 (30%)
Overall	1837/3630 (51%)	1100/1991 (55%)	575/1295 (44%)	162/344 (47%)

7.2.4 Statistically unusual chemical shifts [i](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
2	B	156	TYR	HB3	0.65	0.93 – 4.76	-5.7

7.2.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication

of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain B:

