



wwPDB NMR Structure Validation Summary Report ⓘ

Mar 1, 2026 – 10:47 AM UTC

PDB ID : 2N8M / pdb_00002n8m
BMRB ID : 25855
Title : Zipcode-binding-protein-1 KH3(DD)KH4 domains in complex with the KH4 RNA target
Authors : Nicastro, G.; Ramos, A.; Candel, A.; Hollingworth, D.
Deposited on : 2015-10-21

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
BMRB Restraints Analysis : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 58%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823
RNA backbone	8273	777

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	191	
2	B	7	

2 Ensemble composition and analysis

This entry contains 12 models. Model 2 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:23-A:60, A:67-A:97, A:103-A:142, A:146-A:183 (147)	1.17	2

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 3 single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 5, 6
2	9, 10
3	8, 11
Single-model clusters	4; 7; 12

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3150 atoms, of which 1553 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Insulin-like growth factor 2 mRNA-binding protein 1.

Mol	Chain	Residues	Atoms						Trace
1	A	191	Total	C	H	N	O	S	0
			2926	912	1477	261	272	4	

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	expression tag	UNP O42254
A	2	ALA	-	expression tag	UNP O42254
A	3	MET	-	expression tag	UNP O42254
A	4	GLY	-	expression tag	UNP O42254
A	14	PHE	TYR	engineered mutation	UNP O42254
A	40	ASP	LYS	engineered mutation	UNP O42254
A	41	ASP	LYS	engineered mutation	UNP O42254

- Molecule 2 is a RNA chain called RNA (5'-R(P*UP*CP*GP*GP*AP*CP*U)-3').


Mol	Chain	Residues	Atoms						Trace
2	B	7	Total	C	H	N	O	P	0
			224	66	76	25	50	7	

4 Residue-property plots


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Insulin-like growth factor 2 mRNA-binding protein 1




- Molecule 2: RNA (5'-R(P*UP*CP*GP*GP*AP*CP*U)-3')



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 2. Colouring as in section 4.1 above.

- Molecule 1: Insulin-like growth factor 2 mRNA-binding protein 1





● Molecule 2: RNA (5'-R(P*UP*CP*GP*GP*AP*CP*U)-3')



5 Refinement protocol and experimental data overview

The models were refined using the following method: *torsion angle dynamics*.

Of the 100 calculated structures, 12 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
ARIA	structure solution	
ARIA	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1436
Number of shifts mapped to atoms	1436
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	58%

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.43±0.02	0±0/1161 (0.0± 0.0%)	0.77±0.03	0±1/1566 (0.0± 0.1%)
2	B	0.25±0.02	0±0/164 (0.0± 0.0%)	0.81±0.04	0±0/253 (0.0± 0.0%)
All	All	0.41	0/15900 (0.0%)	0.77	6/21828 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	0.2±0.4
All	All	0	3

There are no bond-length outliers.

5 of 6 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	57	LYS	CA-C-N	5.74	128.33	120.35	6	1
1	A	57	LYS	C-N-CA	5.74	128.33	120.35	6	1
1	A	56	ILE	CA-C-N	5.15	131.11	122.15	11	1
1	A	56	ILE	C-N-CA	5.15	131.11	122.15	11	1
1	A	59	ALA	CA-C-N	5.13	123.35	119.66	3	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	57	LYS	Peptide	2
1	A	67	LYS	Peptide	1

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1141	1198	1193	43±10
2	B	148	76	76	9±2
All	All	15468	15288	15228	608

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

5 of 376 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:52:ALA:HA	1:A:82:LYS:HD2	0.90	1.43	2	2
2:B:4:G:H4'	2:B:5:A:O5'	0.88	1.68	7	12
1:A:130:LEU:HG	1:A:136:ALA:HB3	0.86	1.48	10	3
1:A:56:ILE:HD12	1:A:57:LYS:HG2	0.85	1.46	11	1
1:A:56:ILE:HD12	1:A:57:LYS:HE3	0.81	1.52	6	1

6.3 Torsion angles

6.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	147/191 (77%)	140±2 (95±1%)	6±2 (4±1%)	1±1 (1±1%)	20	70
All	All	1764/2292 (77%)	1680 (95%)	71 (4%)	13 (1%)	20	70

All 5 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	40	ASP	6
1	A	112	PRO	3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	183	LYS	2
1	A	147	ASP	1
1	A	141	PRO	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	120/154 (78%)	113±2 (94±2%)	7±2 (6±2%)	21	70
All	All	1440/1848 (78%)	1360 (94%)	80 (6%)	21	70

5 of 39 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	97	PHE	5
1	A	159	HIS	4
1	A	109	ILE	4
1	A	58	ILE	4
1	A	120	ILE	4

6.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers	Suiteness
2	B	6/7 (86%)	5±0 (83±0%)	2±1 (38±12%)	0.27±0.00
All	All	74/84 (88%)	60 (81%)	27 (36%)	0.27

The overall RNA backbone suiteness is 0.08.

All unique RNA backbone outliers are listed below:

Mol	Chain	Res	Type	Models (Total)
2	B	2	C	12
2	B	3	G	12
2	B	4	G	12
2	B	5	A	12

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
2	B	7	U	12

All unique RNA pucker outliers are listed below:

Mol	Chain	Res	Type	Models (Total)
2	B	4	G	12
2	B	3	G	7
2	B	2	C	6
2	B	1	U	2

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 58% for the well-defined parts and 53% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1436
Number of shifts mapped to atoms	1436
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	119	-0.29 ± 0.29	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	133	0.14 ± 0.10	None needed (< 0.5 ppm)
$^{13}\text{C}'$	0	—	None (insufficient data)
^{15}N	0	—	None (insufficient data)

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 58%, i.e. 1269 atoms were assigned a chemical shift out of a possible 2195. 0 out of 22 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	343/732 (47%)	238/298 (80%)	105/294 (36%)	0/140 (0%)
Sidechain	902/1214 (74%)	635/791 (80%)	267/369 (72%)	0/54 (0%)

Continued on next page...

Continued from previous page...

	Total	^1H	^{13}C	^{15}N
Aromatic	24/118 (20%)	24/58 (41%)	0/50 (0%)	0/10 (0%)
Sugar	0/77 (0%)	0/42 (0%)	0/35 (0%)	0/0 (—%)
Base	0/54 (0%)	0/33 (0%)	0/12 (0%)	0/9 (0%)
Overall	1269/2195 (58%)	897/1222 (73%)	372/760 (49%)	0/213 (0%)


7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:



8 NMR restraints analysis

8.1 Conformationally restricting restraints

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	2220
Intra-residue ($ i-j =0$)	1208
Sequential ($ i-j =1$)	430
Medium range ($ i-j >1$ and $ i-j <5$)	172
Long range ($ i-j \geq 5$)	355
Inter-chain	37
Hydrogen bond restraints	18
Disulfide bond restraints	0
Total dihedral-angle restraints	0
Number of unmapped restraints	0
Number of restraints per residue	11.2
Number of long range restraints per residue ¹	1.8

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	33.5	0.2
0.2-0.5 (Medium)	45.4	0.5
>0.5 (Large)	54.3	7.18

8.2.2 Average number of dihedral-angle violations per model [i](#)

Dihedral-angle violations less than 1° are not included in the calculation. There are no dihedral-angle violations

9 Distance violation analysis ⓘ


9.1 Summary of distance violations ⓘ

The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Restraints type	Count	% ¹	Violated ³			Consistently Violated ⁴		
			Count	% ²	% ¹	Count	% ²	% ¹
Intra-residue ($i-j =0$)	1208	54.4	116	9.6	5.2	3	0.2	0.1
Backbone-Backbone	99	4.5	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	642	28.9	77	12.0	3.5	2	0.3	0.1
Sidechain-Sidechain	467	21.0	39	8.4	1.8	1	0.2	0.0
Sequential ($i-j =1$)	430	19.4	70	16.3	3.2	4	0.9	0.2
Backbone-Backbone	155	7.0	12	7.7	0.5	0	0.0	0.0
Backbone-Sidechain	228	10.3	42	18.4	1.9	2	0.9	0.1
Sidechain-Sidechain	47	2.1	16	34.0	0.7	2	4.3	0.1
Medium range ($i-j >1$ & $i-j <5$)	172	7.7	56	32.6	2.5	2	1.2	0.1
Backbone-Backbone	46	2.1	8	17.4	0.4	0	0.0	0.0
Backbone-Sidechain	79	3.6	24	30.4	1.1	1	1.3	0.0
Sidechain-Sidechain	47	2.1	24	51.1	1.1	1	2.1	0.0
Long range ($i-j \geq 5$)	355	16.0	167	47.0	7.5	7	2.0	0.3
Backbone-Backbone	33	1.5	10	30.3	0.5	0	0.0	0.0
Backbone-Sidechain	121	5.5	53	43.8	2.4	3	2.5	0.1
Sidechain-Sidechain	201	9.1	104	51.7	4.7	4	2.0	0.2
Inter-chain	37	1.7	10	27.0	0.5	1	2.7	0.0
Backbone-Backbone	5	0.2	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	19	0.9	7	36.8	0.3	0	0.0	0.0
Sidechain-Sidechain	13	0.6	3	23.1	0.1	1	7.7	0.0
Hydrogen bond	18	0.8	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	2220	100.0	419	18.9	18.9	17	0.8	0.8
Backbone-Backbone	338	15.2	30	8.9	1.4	0	0.0	0.0
Backbone-Sidechain	1101	49.6	203	18.4	9.1	8	0.7	0.4
Sidechain-Sidechain	781	35.2	186	23.8	8.4	9	1.2	0.4

¹ percentage calculated with respect to the total number of distance restraints, ² percentage calculated with respect to the number of restraints in a particular restraint category, ³ violated in at least one model, ⁴ violated in all the models

9.1.1 Bar chart : Distribution of distance restraints and violations [i](#)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfid bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model [i](#)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

Model ID	Number of violations						Mean (Å)	Max (Å)	SD ⁶ (Å)	Median (Å)
	IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total				
1	29	20	24	64	2	139	0.65	4.93	0.75	0.38
2	33	18	25	53	2	131	0.7	4.87	0.78	0.41
3	42	24	21	70	3	160	0.65	4.7	0.78	0.38
4	26	16	20	55	2	119	0.71	5.8	0.94	0.36
5	39	22	30	63	1	155	0.69	7.18	0.97	0.37
6	36	19	22	64	4	145	0.82	5.73	0.93	0.47
7	31	19	17	42	3	112	0.67	5.69	0.87	0.35
8	33	22	24	53	5	137	0.66	6.56	0.88	0.4
9	41	19	23	57	1	141	0.74	6.89	1.03	0.43
10	35	18	21	55	4	133	0.83	6.9	1.09	0.39


Continued on next page...

Continued from previous page...

Model ID	Number of violations						Mean (Å)	Max (Å)	SD ⁶ (Å)	Median (Å)
	IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total				
11	42	23	19	71	7	162	0.7	7.02	0.93	0.37
12	29	19	17	43	2	110	0.48	2.37	0.45	0.3

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints, ⁵Inter-chain restraints, ⁶Standard deviation

9.2.1 Bar graph : Distance Violation statistics for each model [i](#)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right

9.3 Distance violation statistics for the ensemble [i](#)


Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 1783(IR:1092, SQ:360, MR:116, LR:188, IC:27) restraints are not violated in the ensemble.

Number of violated restraints						Fraction of the ensemble	
IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total	Count ⁶	%
37	23	13	42	3	118	1	8.3
26	16	8	31	2	83	2	16.7
9	6	2	21	2	40	3	25.0
12	9	11	16	0	48	4	33.3
8	3	3	9	1	24	5	41.7
4	3	3	12	1	23	6	50.0
4	2	4	6	0	16	7	58.3
4	2	1	9	0	16	8	66.7
1	1	2	3	0	7	9	75.0
5	1	4	4	0	14	10	83.3
3	0	3	7	0	13	11	91.7
3	4	2	7	1	17	12	100.0

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints,

⁵Inter-chain restraints, ⁶ Number of models with violations


9.3.1 Bar graph : Distance violation statistics for the ensemble [i](#)



9.4 Most violated distance restraints in the ensemble [i](#)

9.4.1 Histogram : Distribution of mean distance violations [i](#)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble



9.4.2 Table: Most violated distance restraints [i](#)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Models ¹	Mean (Å)	SD ¹ (Å)	Median (Å)
(2,14)	1:93:A:GLU:H	1:160:A:PHE:HB2	12	3.72	1.54	3.69
(1,1141)	1:102:A:GLU:HB3	1:135:A:ALA:HB1	12	3.18	1.16	3.08
(1,1141)	1:102:A:GLU:HB3	1:135:A:ALA:HB2	12	3.18	1.16	3.08
(1,1141)	1:102:A:GLU:HB3	1:135:A:ALA:HB3	12	3.18	1.16	3.08
(1,1264)	1:174:A:LEU:HD21	1:179:A:GLN:HG3	12	1.56	1.64	0.41
(1,1264)	1:174:A:LEU:HD22	1:179:A:GLN:HG3	12	1.56	1.64	0.41
(1,1264)	1:174:A:LEU:HD23	1:179:A:GLN:HG3	12	1.56	1.64	0.41
(1,1264)	1:174:A:LEU:HD21	1:179:A:GLN:HG2	12	1.56	1.64	0.41
(1,1264)	1:174:A:LEU:HD22	1:179:A:GLN:HG2	12	1.56	1.64	0.41

Continued on next page...

Continued from previous page...


Key	Atom-1	Atom-2	Models ¹	Mean (Å)	SD ¹ (Å)	Median (Å)
(1,1264)	1:174:A:LEU:HD23	1:179:A:GLN:HG2	12	1.56	1.64	0.41
(1,1357)	1:140:A:VAL:HG21	1:117:A:GLY:HA3	12	1.46	0.33	1.49
(1,1357)	1:140:A:VAL:HG22	1:117:A:GLY:HA3	12	1.46	0.33	1.49
(1,1357)	1:140:A:VAL:HG23	1:117:A:GLY:HA3	12	1.46	0.33	1.49
(2,102)	1:142:A:ARG:NH1	2:5:B:A:OP1	12	1.26	0.65	0.97
(1,1216)	1:150:A:GLU:HG3	1:149:A:ASN:HA	12	1.18	0.21	1.17
(1,1806)	1:48:A:LEU:HB2	1:58:A:ILE:HG21	12	1.14	0.32	1.27
(1,1806)	1:48:A:LEU:HB2	1:58:A:ILE:HG22	12	1.14	0.32	1.27
(1,1806)	1:48:A:LEU:HB2	1:58:A:ILE:HG23	12	1.14	0.32	1.27
(1,538)	1:45:A:ILE:H	1:37:A:ILE:HD11	12	0.93	0.36	0.91
(1,538)	1:45:A:ILE:H	1:37:A:ILE:HD12	12	0.93	0.36	0.91
(1,538)	1:45:A:ILE:H	1:37:A:ILE:HD13	12	0.93	0.36	0.91
(1,1191)	1:105:A:LEU:HD11	1:104:A:LYS:HD2	12	0.83	0.24	0.88
(1,1191)	1:105:A:LEU:HD12	1:104:A:LYS:HD2	12	0.83	0.24	0.88
(1,1191)	1:105:A:LEU:HD13	1:104:A:LYS:HD2	12	0.83	0.24	0.88
(1,1807)	1:48:A:LEU:HD11	1:86:A:ARG:HG2	12	0.82	0.54	0.68

¹Number of violated models, ²Standard deviation

9.5 All violated distance restraints [i](#)

9.5.1 Histogram : Distribution of distance violations [i](#)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



9.5.2 Table : All distance violations [i](#)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(2,32)	1:63:A:THR:HG21	1:58:A:ILE:HA	5	7.18
(2,32)	1:63:A:THR:HG22	1:58:A:ILE:HA	5	7.18
(2,32)	1:63:A:THR:HG23	1:58:A:ILE:HA	5	7.18
(2,32)	1:63:A:THR:HG21	1:58:A:ILE:HA	11	7.02
(2,32)	1:63:A:THR:HG22	1:58:A:ILE:HA	11	7.02
(2,32)	1:63:A:THR:HG23	1:58:A:ILE:HA	11	7.02
(2,32)	1:63:A:THR:HG21	1:58:A:ILE:HA	10	6.9
(2,32)	1:63:A:THR:HG22	1:58:A:ILE:HA	10	6.9
(2,32)	1:63:A:THR:HG23	1:58:A:ILE:HA	10	6.9
(2,32)	1:63:A:THR:HG21	1:58:A:ILE:HA	9	6.89
(2,32)	1:63:A:THR:HG22	1:58:A:ILE:HA	9	6.89
(2,32)	1:63:A:THR:HG23	1:58:A:ILE:HA	9	6.89
(2,32)	1:63:A:THR:HG21	1:58:A:ILE:HA	8	6.56
(2,32)	1:63:A:THR:HG22	1:58:A:ILE:HA	8	6.56
(2,32)	1:63:A:THR:HG23	1:58:A:ILE:HA	8	6.56
(2,59)	1:102:A:GLU:HB2	1:136:A:ALA:HA	4	5.8
(2,14)	1:93:A:GLU:H	1:160:A:PHE:HB2	6	5.73

Continued on next page...

Continued from previous page...

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(2,59)	1:102:A:GLU:HB2	1:136:A:ALA:HA	5	5.72
(2,14)	1:93:A:GLU:H	1:160:A:PHE:HB2	7	5.69
(1,1365)	1:32:A:GLN:HG2	1:96:A:PHE:HD1	9	5.67

10 Dihedral-angle violation analysis ⓘ

No dihedral-angle restraints found