



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 18, 2026 – 02:56 PM UTC

PDB ID : 3N84 / pdb_00003n84
Title : Crystal Structure of the Grb2 SH2 Domain in Complex with a 23-Membered Macrocyclic Ligand Having the Sequence pYVNVP
Authors : Clements, J.H.; Martin, S.F.
Deposited on : 2010-05-27
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	112	 86% 9% . .
1	B	112	 85% 6% . . 5%
1	C	112	 93% . . .
1	D	112	 82% 6% . 10%
1	E	112	 86% . . . 9%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	112	 84% 5% • 10%
2	G	6	 100%
2	H	6	 83% 17%
2	I	6	 100%
2	J	6	 67% 33%
2	K	6	 83% 17%
2	L	6	 33% 50% 17%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	F	7	-	-	X	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6278 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Growth factor receptor-bound protein 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	109	Total	C	N	O	S	0	0	0
			898	576	160	160	2			
1	B	106	Total	C	N	O	S	0	0	0
			873	559	156	156	2			
1	C	109	Total	C	N	O	S	0	0	0
			898	576	160	160	2			
1	D	101	Total	C	N	O	S	0	0	0
			834	535	149	148	2			
1	E	102	Total	C	N	O	S	0	0	0
			843	541	150	149	3			
1	F	101	Total	C	N	O	S	0	0	0
			835	536	149	148	2			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	52	MET	-	expression tag	UNP P62993
B	52	MET	-	expression tag	UNP P62993
C	52	MET	-	expression tag	UNP P62993
D	52	MET	-	expression tag	UNP P62993
E	52	MET	-	expression tag	UNP P62993
F	52	MET	-	expression tag	UNP P62993

- Molecule 2 is a protein called 23-membered peptide-like macrocyclic ligand.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	G	6	Total	C	N	O	P	0	0	0
			54	35	7	11	1			
2	H	6	Total	C	N	O	P	0	0	0
			54	35	7	11	1			
2	I	6	Total	C	N	O	P	0	0	0
			54	35	7	11	1			

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	J	6	Total	C	N	O	P	0	0	0
			54	35	7	11	1			
2	K	6	Total	C	N	O	P	0	0	0
			54	35	7	11	1			
2	L	6	Total	C	N	O	P	0	0	0
			54	35	7	11	1			

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Cl	0	0
			2	2		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		
4	D	1	Total	C	O	0	0
			6	3	3		
4	E	1	Total	C	O	0	0
			6	3	3		
4	F	1	Total	C	O	0	0
			6	3	3		
4	F	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	F	1	Total	C	O	0	0
			6	3	3		
4	K	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	1	Total	Mg	0	0
			1	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	139	Total	O	0	0
			139	139		
6	B	99	Total	O	0	0
			99	99		
6	C	124	Total	O	0	0
			124	124		
6	D	100	Total	O	0	0
			100	100		
6	E	108	Total	O	0	0
			108	108		
6	F	105	Total	O	0	0
			105	105		
6	G	12	Total	O	0	0
			12	12		
6	H	10	Total	O	0	0
			10	10		
6	I	11	Total	O	0	0
			11	11		
6	J	8	Total	O	0	0
			8	8		
6	K	10	Total	O	0	0
			10	10		
6	L	2	Total	O	0	0
			2	2		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Growth factor receptor-bound protein 2

Chain A:  86% 9% . .



- Molecule 1: Growth factor receptor-bound protein 2

Chain B:  85% 6% . . 5%



- Molecule 1: Growth factor receptor-bound protein 2

Chain C:  93% . . .



- Molecule 1: Growth factor receptor-bound protein 2

Chain D:  82% 6% . 10%



- Molecule 1: Growth factor receptor-bound protein 2

Chain E:  86% . . . 9%



- Molecule 1: Growth factor receptor-bound protein 2

Chain F:  84% 5% . 10%



- Molecule 2: 23-membered peptide-like macrocyclic ligand

Chain G: 100%

There are no outlier residues recorded for this chain.

- Molecule 2: 23-membered peptide-like macrocyclic ligand

Chain H: 83% 17%



- Molecule 2: 23-membered peptide-like macrocyclic ligand

Chain I: 100%

There are no outlier residues recorded for this chain.

- Molecule 2: 23-membered peptide-like macrocyclic ligand

Chain J: 67% 33%



- Molecule 2: 23-membered peptide-like macrocyclic ligand

Chain K: 83% 17%



- Molecule 2: 23-membered peptide-like macrocyclic ligand

Chain L: 33% 50% 17%



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	83.22Å 141.32Å 62.45Å 90.00° 89.99° 90.00°	Depositor
Resolution (Å)	50.00 – 2.00 50.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	94.6 (50.00-2.00) 94.6 (50.00-2.00)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.08 (at 1.70Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.180 , 0.223 0.180 , 0.167	Depositor DCC
R_{free} test set	3629 reflections (4.57%)	wwPDB-VP
Wilson B-factor (Å ²)	20.5	Xtriage
Anisotropy	0.136	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 32.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.012 for -1/2*h+1/2*k,3/2*h+1/2*k,-l 0.013 for -1/2*h-1/2*k,-3/2*h+1/2*k,-l 0.012 for 1/2*h+1/2*k,3/2*h-1/2*k,-l 0.013 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.457 for -h,-k,l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6278	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 011, MG, CL, GOL, PTR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.86	0/923	0.99	2/1243 (0.2%)
1	B	0.94	2/897 (0.2%)	1.35	12/1208 (1.0%)
1	C	0.89	0/923	1.00	2/1243 (0.2%)
1	D	0.85	0/856	0.95	2/1150 (0.2%)
1	E	1.04	2/865 (0.2%)	1.15	5/1161 (0.4%)
1	F	0.88	0/857	0.95	0/1151
2	G	0.72	0/29	1.22	0/40
2	H	0.61	0/29	1.12	0/40
2	I	0.79	0/29	1.09	0/40
2	J	0.57	0/29	1.19	1/40 (2.5%)
2	K	0.81	0/29	1.11	0/40
2	L	0.69	0/29	1.14	0/40
All	All	0.91	4/5495 (0.1%)	1.08	24/7396 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	E	0	1
All	All	0	2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	52	MET	SD-CE	-15.84	1.40	1.79
1	B	55	MET	SD-CE	5.57	1.93	1.79
1	E	73	MET	SD-CE	5.24	1.92	1.79
1	B	105	VAL	CA-CB	5.02	1.60	1.54

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	157	GLN	CA-C-N	12.62	135.62	119.84
1	B	157	GLN	C-N-CA	12.62	135.62	119.84
1	E	52	MET	CA-C-N	12.07	143.70	121.97
1	E	52	MET	C-N-CA	12.07	143.70	121.97
1	B	156	GLN	CA-C-N	10.23	137.23	120.86

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	156	GLN	Mainchain
1	E	52	MET	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	898	0	878	11	0
1	B	873	0	853	4	0
1	C	898	0	878	4	0
1	D	834	0	816	5	0
1	E	843	0	827	6	0
1	F	835	0	818	8	0
2	G	54	0	49	0	0
2	H	54	0	48	1	0
2	I	54	0	48	0	0
2	J	54	0	48	1	0
2	K	54	0	48	1	0
2	L	54	0	48	4	0
3	A	2	0	0	0	0
4	B	6	0	8	0	0
4	D	6	0	8	0	0
4	E	6	0	8	2	0
4	F	18	0	24	9	0
4	K	6	0	8	1	0
5	C	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	139	0	0	7	2
6	B	99	0	0	3	0
6	C	124	0	0	3	1
6	D	100	0	0	3	1
6	E	108	0	0	1	0
6	F	105	0	0	3	1
6	G	12	0	0	0	0
6	H	10	0	0	0	0
6	I	11	0	0	0	0
6	J	8	0	0	1	0
6	K	10	0	0	0	0
6	L	2	0	0	0	0
All	All	6278	0	5415	43	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 43 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:52:MET:N	6:E:720:HOH:O	1.83	1.07
4:F:7:GOL:H31	2:L:1:PTR:O3P	1.75	0.87
1:F:53:ILE:N	6:F:274:HOH:O	2.16	0.79
1:D:73:MET:HE1	1:D:151:ILE:HD13	1.75	0.69
1:F:95:PHE:H	4:F:4:GOL:H32	1.59	0.68

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:673:HOH:O	6:D:437:HOH:O[2_555]	2.14	0.06
6:C:580:HOH:O	6:F:579:HOH:O[1_655]	2.16	0.04
6:A:507:HOH:O	6:A:561:HOH:O[2_556]	2.18	0.02

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	107/112 (96%)	106 (99%)	1 (1%)	0	100	100
1	B	104/112 (93%)	98 (94%)	3 (3%)	3 (3%)	3	1
1	C	107/112 (96%)	104 (97%)	3 (3%)	0	100	100
1	D	99/112 (88%)	98 (99%)	1 (1%)	0	100	100
1	E	100/112 (89%)	98 (98%)	1 (1%)	1 (1%)	12	8
1	F	99/112 (88%)	97 (98%)	2 (2%)	0	100	100
2	G	4/6 (67%)	4 (100%)	0	0	100	100
2	H	4/6 (67%)	4 (100%)	0	0	100	100
2	I	4/6 (67%)	4 (100%)	0	0	100	100
2	J	4/6 (67%)	4 (100%)	0	0	100	100
2	K	4/6 (67%)	4 (100%)	0	0	100	100
2	L	4/6 (67%)	4 (100%)	0	0	100	100
All	All	640/708 (90%)	625 (98%)	11 (2%)	4 (1%)	21	17

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	155	PRO
1	B	156	GLN
1	B	158	PRO
1	E	53	ILE

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	97/100 (97%)	92 (95%)	5 (5%)	21	18
1	B	95/100 (95%)	91 (96%)	4 (4%)	26	25
1	C	97/100 (97%)	93 (96%)	4 (4%)	27	26
1	D	90/100 (90%)	87 (97%)	3 (3%)	33	34
1	E	91/100 (91%)	90 (99%)	1 (1%)	65	73
1	F	90/100 (90%)	88 (98%)	2 (2%)	45	50
2	G	4/4 (100%)	4 (100%)	0	100	100
2	H	4/4 (100%)	4 (100%)	0	100	100
2	I	4/4 (100%)	4 (100%)	0	100	100
2	J	4/4 (100%)	4 (100%)	0	100	100
2	K	4/4 (100%)	4 (100%)	0	100	100
2	L	4/4 (100%)	4 (100%)	0	100	100
All	All	584/624 (94%)	565 (97%)	19 (3%)	33	34

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	78	ARG
1	F	78	ARG
1	F	153	GLN
1	E	153	GLN
1	B	157	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	79	HIS
1	F	79	HIS
1	C	143	ASN
1	F	153	GLN
1	E	135	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

12 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PTR	I	1	2	15,16,17	0.96	0	17,22,24	0.81	0
2	011	I	6	2	7,8,9	0.40	0	6,7,9	0.44	0
2	PTR	K	1	2	15,16,17	0.97	1 (6%)	17,22,24	1.02	1 (5%)
2	PTR	H	1	2	15,16,17	0.97	0	17,22,24	0.76	0
2	PTR	L	1	2	15,16,17	0.93	0	17,22,24	1.05	1 (5%)
2	011	K	6	2	7,8,9	0.46	0	6,7,9	0.50	0
2	011	L	6	2	7,8,9	0.36	0	6,7,9	0.52	0
2	011	H	6	2	7,8,9	0.40	0	6,7,9	0.45	0
2	PTR	G	1	2	15,16,17	0.98	0	17,22,24	0.81	0
2	011	G	6	2	7,8,9	0.36	0	6,7,9	0.66	0
2	PTR	J	1	2	15,16,17	0.97	0	17,22,24	0.80	0
2	011	J	6	2	7,8,9	0.38	0	6,7,9	0.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PTR	I	1	2	-	0/10/11/13	0/1/1/1
2	011	I	6	2	-	1/6/6/7	-
2	PTR	K	1	2	-	2/10/11/13	0/1/1/1
2	PTR	H	1	2	-	3/10/11/13	0/1/1/1
2	PTR	L	1	2	-	0/10/11/13	0/1/1/1
2	011	K	6	2	-	1/6/6/7	-
2	011	L	6	2	-	1/6/6/7	-
2	011	H	6	2	-	1/6/6/7	-
2	PTR	G	1	2	-	0/10/11/13	0/1/1/1
2	011	G	6	2	-	4/6/6/7	-
2	PTR	J	1	2	-	2/10/11/13	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	011	J	6	2	-	1/6/6/7	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	K	1	PTR	OH-CZ	-2.06	1.36	1.40

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	L	1	PTR	OH-CZ-CE2	2.13	125.59	119.22
2	K	1	PTR	OH-CZ-CE2	2.10	125.51	119.22

There are no chirality outliers.

5 of 16 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	H	1	PTR	O-C-CA-CB
2	H	1	PTR	CZ-OH-P-O1P
2	J	1	PTR	CZ-OH-P-O1P
2	G	6	011	O-C-C2-C3
2	H	6	011	O-C-C2-C3

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	K	1	PTR	1	0
2	L	1	PTR	3	0
2	L	6	011	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 3 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul

statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	F	7	-	5,5,5	0.55	0	5,5,5	0.42	0
4	GOL	D	3	-	5,5,5	0.23	0	5,5,5	0.53	0
4	GOL	F	2	-	5,5,5	0.29	0	5,5,5	0.60	0
4	GOL	B	6	-	5,5,5	0.43	0	5,5,5	0.29	0
4	GOL	E	1	-	5,5,5	0.31	0	5,5,5	0.97	0
4	GOL	F	4	-	5,5,5	0.56	0	5,5,5	1.25	1 (20%)
4	GOL	K	7	-	5,5,5	0.39	0	5,5,5	0.69	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	F	7	-	-	0/4/4/4	-
4	GOL	D	3	-	-	0/4/4/4	-
4	GOL	F	2	-	-	0/4/4/4	-
4	GOL	B	6	-	-	0/4/4/4	-
4	GOL	E	1	-	-	0/4/4/4	-
4	GOL	F	4	-	-	0/4/4/4	-
4	GOL	K	7	-	-	0/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	F	4	GOL	C3-C2-C1	-2.09	104.11	111.80

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	F	7	GOL	4	0
4	F	2	GOL	3	0
4	E	1	GOL	2	0
4	F	4	GOL	2	0
4	K	7	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	109/112 (97%)	-1.61	0 100 100	10, 16, 34, 39	0
1	B	106/112 (94%)	-1.58	0 100 100	11, 17, 37, 43	0
1	C	109/112 (97%)	-1.59	0 100 100	10, 16, 33, 38	0
1	D	101/112 (90%)	-1.59	0 100 100	12, 17, 30, 41	0
1	E	102/112 (91%)	-1.60	0 100 100	10, 16, 28, 39	0
1	F	101/112 (90%)	-1.60	0 100 100	9, 16, 27, 40	0
2	G	4/6 (66%)	-1.81	0 100 100	11, 11, 12, 12	0
2	H	4/6 (66%)	-1.66	0 100 100	13, 13, 16, 20	0
2	I	4/6 (66%)	-1.76	0 100 100	11, 12, 13, 13	0
2	J	4/6 (66%)	-1.68	0 100 100	11, 13, 14, 20	0
2	K	4/6 (66%)	-1.64	0 100 100	14, 14, 15, 18	0
2	L	4/6 (66%)	-1.68	0 100 100	12, 13, 16, 17	0
All	All	652/708 (92%)	-1.60	0 100 100	9, 16, 32, 43	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	011	G	6	9/10	0.99	0.03	13,16,19,20	0
2	011	H	6	9/10	0.99	0.03	14,16,18,19	0
2	011	I	6	9/10	0.99	0.03	13,15,16,17	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	011	K	6	9/10	0.99	0.03	13,15,17,18	0
2	PTR	K	1	16/17	1.00	0.02	12,14,17,20	0
2	PTR	L	1	16/17	1.00	0.03	12,16,18,20	0
2	PTR	G	1	16/17	1.00	0.02	11,13,16,16	0
2	PTR	H	1	16/17	1.00	0.02	12,13,18,19	0
2	PTR	I	1	16/17	1.00	0.02	10,16,18,19	0
2	011	J	6	9/10	1.00	0.02	13,16,18,18	0
2	PTR	J	1	16/17	1.00	0.02	13,16,18,19	0
2	011	L	6	9/10	1.00	0.02	13,16,17,18	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	CL	A	10	1/1	0.98	0.12	94,94,94,94	0
4	GOL	B	6	6/6	0.98	0.04	36,36,37,38	0
4	GOL	D	3	6/6	0.98	0.04	34,35,36,37	0
4	GOL	F	2	6/6	0.98	0.04	30,33,36,36	0
4	GOL	K	7	6/6	0.98	0.04	33,36,36,38	0
3	CL	A	9	1/1	0.99	0.03	20,20,20,20	0
4	GOL	F	4	6/6	0.99	0.04	27,29,31,31	0
4	GOL	F	7	6/6	0.99	0.04	33,35,36,37	0
4	GOL	E	1	6/6	0.99	0.03	25,28,30,32	0
5	MG	C	8	1/1	1.00	0.03	13,13,13,13	0

6.5 Other polymers [i](#)

There are no such residues in this entry.