



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 06:04 PM UTC

PDB ID : 5N83 / pdb_00005n83
Title : Structure of the distal domain of mouse adenovirus 2 fibre, methylmercury chloride derivative
Authors : Singh, A.K.; van Raaij, M.J.
Deposited on : 2017-02-22
Resolution : 2.76 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1009 (2.76-2.76)
Clashscore	190562	1044 (2.76-2.76)
Ramachandran outliers	187476	1024 (2.76-2.76)
Sidechain outliers	187428	1024 (2.76-2.76)
RSRZ outliers	180081	1009 (2.76-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	237	5% 66% 14% • 18%
1	B	237	6% 71% 10% • 18%
1	C	237	7% 69% 12% • 17%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	HG	C	802	-	-	X	-

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4578 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fiber.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	194	Total	C	N	O	S	0	0	0
			1505	957	242	295	11			
1	B	195	Total	C	N	O	S	0	0	0
			1510	960	243	296	11			
1	C	196	Total	C	N	O	S	0	0	0
			1514	962	244	297	11			

There are 105 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	551	GLY	-	expression tag	UNP E7CH51
A	552	SER	-	expression tag	UNP E7CH51
A	553	SER	-	expression tag	UNP E7CH51
A	554	HIS	-	expression tag	UNP E7CH51
A	555	HIS	-	expression tag	UNP E7CH51
A	556	HIS	-	expression tag	UNP E7CH51
A	557	HIS	-	expression tag	UNP E7CH51
A	558	HIS	-	expression tag	UNP E7CH51
A	559	HIS	-	expression tag	UNP E7CH51
A	560	SER	-	expression tag	UNP E7CH51
A	561	SER	-	expression tag	UNP E7CH51
A	562	GLY	-	expression tag	UNP E7CH51
A	563	LEU	-	expression tag	UNP E7CH51
A	564	VAL	-	expression tag	UNP E7CH51
A	565	PRO	-	expression tag	UNP E7CH51
A	566	ARG	-	expression tag	UNP E7CH51
A	567	GLY	-	expression tag	UNP E7CH51
A	568	SER	-	expression tag	UNP E7CH51
A	569	HIS	-	expression tag	UNP E7CH51
A	570	MET	-	expression tag	UNP E7CH51
A	571	ALA	-	expression tag	UNP E7CH51
A	572	SER	-	expression tag	UNP E7CH51
A	573	MET	-	expression tag	UNP E7CH51

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	574	THR	-	expression tag	UNP E7CH51
A	575	GLY	-	expression tag	UNP E7CH51
A	576	GLY	-	expression tag	UNP E7CH51
A	577	GLN	-	expression tag	UNP E7CH51
A	578	GLN	-	expression tag	UNP E7CH51
A	579	MET	-	expression tag	UNP E7CH51
A	580	GLY	-	expression tag	UNP E7CH51
A	581	ARG	-	expression tag	UNP E7CH51
A	582	GLY	-	expression tag	UNP E7CH51
A	583	SER	-	expression tag	UNP E7CH51
A	584	GLU	-	expression tag	UNP E7CH51
A	585	PHE	-	expression tag	UNP E7CH51
B	551	GLY	-	expression tag	UNP E7CH51
B	552	SER	-	expression tag	UNP E7CH51
B	553	SER	-	expression tag	UNP E7CH51
B	554	HIS	-	expression tag	UNP E7CH51
B	555	HIS	-	expression tag	UNP E7CH51
B	556	HIS	-	expression tag	UNP E7CH51
B	557	HIS	-	expression tag	UNP E7CH51
B	558	HIS	-	expression tag	UNP E7CH51
B	559	HIS	-	expression tag	UNP E7CH51
B	560	SER	-	expression tag	UNP E7CH51
B	561	SER	-	expression tag	UNP E7CH51
B	562	GLY	-	expression tag	UNP E7CH51
B	563	LEU	-	expression tag	UNP E7CH51
B	564	VAL	-	expression tag	UNP E7CH51
B	565	PRO	-	expression tag	UNP E7CH51
B	566	ARG	-	expression tag	UNP E7CH51
B	567	GLY	-	expression tag	UNP E7CH51
B	568	SER	-	expression tag	UNP E7CH51
B	569	HIS	-	expression tag	UNP E7CH51
B	570	MET	-	expression tag	UNP E7CH51
B	571	ALA	-	expression tag	UNP E7CH51
B	572	SER	-	expression tag	UNP E7CH51
B	573	MET	-	expression tag	UNP E7CH51
B	574	THR	-	expression tag	UNP E7CH51
B	575	GLY	-	expression tag	UNP E7CH51
B	576	GLY	-	expression tag	UNP E7CH51
B	577	GLN	-	expression tag	UNP E7CH51
B	578	GLN	-	expression tag	UNP E7CH51
B	579	MET	-	expression tag	UNP E7CH51
B	580	GLY	-	expression tag	UNP E7CH51

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	581	ARG	-	expression tag	UNP E7CH51
B	582	GLY	-	expression tag	UNP E7CH51
B	583	SER	-	expression tag	UNP E7CH51
B	584	GLU	-	expression tag	UNP E7CH51
B	585	PHE	-	expression tag	UNP E7CH51
C	551	GLY	-	expression tag	UNP E7CH51
C	552	SER	-	expression tag	UNP E7CH51
C	553	SER	-	expression tag	UNP E7CH51
C	554	HIS	-	expression tag	UNP E7CH51
C	555	HIS	-	expression tag	UNP E7CH51
C	556	HIS	-	expression tag	UNP E7CH51
C	557	HIS	-	expression tag	UNP E7CH51
C	558	HIS	-	expression tag	UNP E7CH51
C	559	HIS	-	expression tag	UNP E7CH51
C	560	SER	-	expression tag	UNP E7CH51
C	561	SER	-	expression tag	UNP E7CH51
C	562	GLY	-	expression tag	UNP E7CH51
C	563	LEU	-	expression tag	UNP E7CH51
C	564	VAL	-	expression tag	UNP E7CH51
C	565	PRO	-	expression tag	UNP E7CH51
C	566	ARG	-	expression tag	UNP E7CH51
C	567	GLY	-	expression tag	UNP E7CH51
C	568	SER	-	expression tag	UNP E7CH51
C	569	HIS	-	expression tag	UNP E7CH51
C	570	MET	-	expression tag	UNP E7CH51
C	571	ALA	-	expression tag	UNP E7CH51
C	572	SER	-	expression tag	UNP E7CH51
C	573	MET	-	expression tag	UNP E7CH51
C	574	THR	-	expression tag	UNP E7CH51
C	575	GLY	-	expression tag	UNP E7CH51
C	576	GLY	-	expression tag	UNP E7CH51
C	577	GLN	-	expression tag	UNP E7CH51
C	578	GLN	-	expression tag	UNP E7CH51
C	579	MET	-	expression tag	UNP E7CH51
C	580	GLY	-	expression tag	UNP E7CH51
C	581	ARG	-	expression tag	UNP E7CH51
C	582	GLY	-	expression tag	UNP E7CH51
C	583	SER	-	expression tag	UNP E7CH51
C	584	GLU	-	expression tag	UNP E7CH51
C	585	PHE	-	expression tag	UNP E7CH51

- Molecule 2 is MERCURY (II) ION (CCD ID: HG) (formula: Hg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total Hg 2 2	0	0
2	B	1	Total Hg 1 1	0	0
2	C	2	Total Hg 2 2	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	22	Total O 22 22	0	0
3	B	9	Total O 9 9	0	0
3	C	13	Total O 13 13	0	0

- Molecule 1: Fiber



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	92.12Å 164.28Å 96.43Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	82.14 – 2.76 82.14 – 2.76	Depositor EDS
% Data completeness (in resolution range)	99.6 (82.14-2.76) 99.7 (82.14-2.76)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.03 (at 2.77Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.203 , 0.248 (Not available) , 0.231	Depositor DCC
R_{free} test set	985 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å ²)	57.7	Xtriage
Anisotropy	0.208	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 38.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.016 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.026 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4578	wwPDB-VP
Average B, all atoms (Å ²)	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section:
HG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.69	1/1540 (0.1%)	0.86	4/2091 (0.2%)
1	B	0.62	0/1545	0.77	0/2098
1	C	0.64	0/1549	0.76	1/2103 (0.0%)
All	All	0.65	1/4634 (0.0%)	0.80	5/6292 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
All	All	0	3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	725	SER	CA-C	6.75	1.59	1.53

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	725	SER	N-CA-C	10.54	125.89	108.08
1	A	727	TYR	N-CA-C	8.37	122.91	110.52
1	A	663	PHE	CB-CA-C	-7.33	97.99	110.16
1	C	723	ASP	N-CA-C	5.82	119.89	111.56
1	A	637	THR	N-CA-C	-5.00	103.48	109.83

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	728	LYS	Peptide
1	B	727	TYR	Peptide
1	C	723	ASP	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1505	0	1456	17	0
1	B	1510	0	1461	11	0
1	C	1514	0	1464	16	0
2	A	2	0	0	2	0
2	B	1	0	0	1	0
2	C	2	0	0	3	0
3	A	22	0	0	1	0
3	B	9	0	0	0	0
3	C	13	0	0	1	0
All	All	4578	0	4381	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 43 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:736:CYS:SG	2:C:802:HG:HG	1.46	1.31
1:A:696:CYS:SG	2:A:801:HG:HG	1.57	1.22
1:B:748:CYS:SG	2:B:801:HG:HG	1.73	1.05
1:A:748:CYS:SG	2:A:802:HG:HG	1.89	0.91
1:C:748:CYS:SG	2:C:801:HG:HG	1.95	0.84

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	192/237 (81%)	178 (93%)	11 (6%)	3 (2%)	7	14
1	B	193/237 (81%)	179 (93%)	12 (6%)	2 (1%)	12	24
1	C	194/237 (82%)	180 (93%)	13 (7%)	1 (0%)	24	43
All	All	579/711 (81%)	537 (93%)	36 (6%)	6 (1%)	12	24

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	721	THR
1	B	679	ALA
1	B	721	THR
1	C	721	THR
1	A	728	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	168/198 (85%)	152 (90%)	16 (10%)	8	16
1	B	168/198 (85%)	159 (95%)	9 (5%)	20	38
1	C	168/198 (85%)	154 (92%)	14 (8%)	10	19
All	All	504/594 (85%)	465 (92%)	39 (8%)	12	22

5 of 39 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	683	GLN
1	C	767	THR
1	C	704	MET
1	C	727	TYR
1	C	776	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	636	GLN
1	A	783	GLN
1	C	636	GLN
1	C	783	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 5 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	194/237 (81%)	0.31	11 (5%) 29 29	33, 52, 100, 143	0
1	B	195/237 (82%)	0.36	15 (7%) 19 19	35, 55, 111, 137	0
1	C	196/237 (82%)	0.39	16 (8%) 17 18	36, 58, 116, 144	0
All	All	585/711 (82%)	0.36	42 (7%) 21 21	33, 56, 110, 144	0

The worst 5 of 42 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	727	TYR	6.0
1	B	596	PHE	4.2
1	A	726	VAL	4.0
1	B	593	ALA	3.9
1	B	679	ALA	3.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	HG	A	802	1/1	0.97	0.10	83,83,83,83	1
2	HG	C	802	1/1	0.97	0.06	88,88,88,88	1
2	HG	C	801	1/1	0.99	0.08	81,81,81,81	1
2	HG	B	801	1/1	0.99	0.03	47,47,47,47	1
2	HG	A	801	1/1	1.00	0.02	49,49,49,49	1

6.5 Other polymers [i](#)

There are no such residues in this entry.