



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 28, 2026 – 08:08 PM UTC

PDB ID : 4N99 / pdb_00004n99
Title : E. coli sliding clamp in complex with 6-chloro-2,3,4,9-tetrahydro-1H-carbazol
e-7-carboxylic acid
Authors : Yin, Z.; Oakley, A.J.
Deposited on : 2013-10-19
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	366	5% 80% 16% ..
1	B	366	6% 89% 10%

2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 6068 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase III subunit beta.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	356	Total	C	N	O	S	0	1	0
			2732	1725	468	519	20			
1	B	365	Total	C	N	O	S	0	2	0
			2818	1771	489	538	20			

- Molecule 2 is 6-chloro-2,3,4,9-tetrahydro-1H-carbazole-7-carboxylic acid (CCD ID: 2J1) (formula: C₁₃H₁₂ClNO₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Cl	N	O	0	0
			17	13	1	1	2		

- Molecule 3 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: C₄H₁₀O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	4	3		
3	A	1	Total	C	O	0	0
			7	4	3		
3	A	1	Total	C	O	0	0
			7	4	3		
3	A	1	Total	C	O	0	0
			7	4	3		
3	A	1	Total	C	O	0	0
			7	4	3		
3	A	1	Total	C	O	0	0
			7	4	3		
3	A	1	Total	C	O	0	0
			7	4	3		
3	B	1	Total	C	O	0	0
			7	4	3		
3	B	1	Total	C	O	0	0
			7	4	3		
3	B	1	Total	C	O	0	0
			7	4	3		
3	B	1	Total	C	O	0	0
			7	4	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			7	4	3		
3	B	1	Total	C	O	0	0
			7	4	3		
3	B	1	Total	C	O	0	0
			7	4	3		
3	B	1	Total	C	O	0	0
			7	4	3		
3	B	1	Total	C	O	0	0
			7	4	3		

- Molecule 4 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			13	8	5		
4	A	1	Total	C	O	0	0
			13	8	5		
4	A	1	Total	C	O	0	0
			13	8	5		
4	A	1	Total	C	O	0	0
			13	8	5		
4	B	1	Total	C	O	0	0
			13	8	5		
4	B	1	Total	C	O	0	0
			13	8	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			13	8	5		
4	B	1	Total	C	O	0	0
			13	8	5		
4	B	1	Total	C	O	0	0
			13	8	5		
4	B	1	Total	C	O	0	0
			13	8	5		

- Molecule 5 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	4	Total	Ca	0	0
			4	4		
5	B	1	Total	Ca	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	120	Total	O	0	0
			120	120		
6	B	113	Total	O	0	0
			113	113		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA polymerase III subunit beta



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	41.01Å 65.13Å 73.24Å 72.95° 85.06° 85.24°	Depositor
Resolution (Å)	40.95 – 2.30 40.95 – 2.30	Depositor EDS
% Data completeness (in resolution range)	92.9 (40.95-2.30) 92.9 (40.95-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.15	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.89 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.261 , 0.338 0.258 , 0.334	Depositor DCC
R_{free} test set	1495 reflections (4.67%)	wwPDB-VP
Wilson B-factor (Å ²)	26.1	Xtriage
Anisotropy	0.023	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 38.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.87	EDS
Total number of atoms	6068	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PG4, 2J1, PEG, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	0/2781	0.73	0/3768
1	B	0.38	0/2871	0.72	0/3887
All	All	0.37	0/5652	0.72	0/7655

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2732	0	2719	33	0
1	B	2818	0	2821	23	0
2	A	17	0	11	1	0
3	A	63	0	90	0	0
3	B	70	0	100	4	0
4	A	52	0	72	2	0
4	B	78	0	108	9	0
5	A	4	0	0	0	0
5	B	1	0	0	0	0
6	A	120	0	0	0	0
6	B	113	0	0	0	0
All	All	6068	0	5921	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 59 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:32:ASN:HB3	1:A:69:THR:HG22	1.71	0.71
1:A:242:PRO:HB3	2:A:401:2J1:H7	1.75	0.67
1:B:105:ARG:HE	4:B:412:PG4:H22	1.60	0.67
1:A:256:LEU:HB3	1:A:338[B]:MET:HG3	1.80	0.63
1:A:252:PRO:HB2	1:A:339:MET:HB3	1.82	0.61

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	351/366 (96%)	340 (97%)	11 (3%)	0	100	100
1	B	363/366 (99%)	358 (99%)	5 (1%)	0	100	100
All	All	714/732 (98%)	698 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	297/313 (95%)	286 (96%)	11 (4%)	30	45
1	B	309/313 (99%)	303 (98%)	6 (2%)	50	69
All	All	606/626 (97%)	589 (97%)	17 (3%)	38	56

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	251	ASN
1	B	366	LEU
1	A	212	ASN
1	A	256	LEU
1	A	282	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	295	ASN
1	B	289	GLN
1	B	91	GLN
1	B	212	ASN
1	B	16	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 35 ligands modelled in this entry, 5 are monoatomic - leaving 30 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PEG	B	405	-	6,6,6	0.46	0	5,5,5	0.30	0
3	PEG	B	414	-	6,6,6	0.44	0	5,5,5	0.31	0
3	PEG	A	418	-	6,6,6	0.46	0	5,5,5	0.30	0
4	PG4	B	404	-	12,12,12	0.48	0	11,11,11	0.23	0
3	PEG	A	414	-	6,6,6	0.45	0	5,5,5	0.28	0
3	PEG	A	408	-	6,6,6	0.46	0	5,5,5	0.24	0
4	PG4	B	401	-	12,12,12	0.48	0	11,11,11	0.21	0
3	PEG	B	403	-	6,6,6	0.47	0	5,5,5	0.29	0
3	PEG	A	402	-	6,6,6	0.46	0	5,5,5	0.26	0
3	PEG	A	403	-	6,6,6	0.45	0	5,5,5	0.29	0
3	PEG	A	413	-	6,6,6	0.45	0	5,5,5	0.24	0
3	PEG	B	407	-	6,6,6	0.44	0	5,5,5	0.32	0
3	PEG	B	409	-	6,6,6	0.45	0	5,5,5	0.28	0
4	PG4	B	412	-	12,12,12	0.47	0	11,11,11	0.24	0
3	PEG	B	413	-	6,6,6	0.44	0	5,5,5	0.27	0
4	PG4	B	402	-	12,12,12	0.47	0	11,11,11	0.26	0
3	PEG	A	405	-	6,6,6	0.48	0	5,5,5	0.28	0
4	PG4	A	410	-	12,12,12	0.48	0	11,11,11	0.22	0
3	PEG	B	415	-	6,6,6	0.43	0	5,5,5	0.32	0
3	PEG	A	412	-	6,6,6	0.46	0	5,5,5	0.19	0
3	PEG	B	416	-	6,6,6	0.48	0	5,5,5	0.24	0
4	PG4	B	406	-	12,12,12	0.48	0	11,11,11	0.24	0
4	PG4	B	411	-	12,12,12	0.50	0	11,11,11	0.19	0
3	PEG	B	410	-	6,6,6	0.45	0	5,5,5	0.24	0
4	PG4	A	416	-	12,12,12	0.46	0	11,11,11	0.25	0
4	PG4	A	404	-	12,12,12	0.48	0	11,11,11	0.25	0
4	PG4	A	415	-	12,12,12	0.47	0	11,11,11	0.26	0
3	PEG	B	408	-	6,6,6	0.45	0	5,5,5	0.25	0
2	2J1	A	401	-	19,19,19	3.01	7 (36%)	28,28,28	2.30	9 (32%)
3	PEG	A	406	-	6,6,6	0.45	0	5,5,5	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PEG	B	405	-	-	3/4/4/4	-
3	PEG	B	414	-	-	2/4/4/4	-
3	PEG	A	418	-	-	3/4/4/4	-
4	PG4	B	404	-	-	6/10/10/10	-
3	PEG	A	414	-	-	1/4/4/4	-
3	PEG	A	408	-	-	3/4/4/4	-
4	PG4	B	401	-	-	6/10/10/10	-
3	PEG	B	403	-	-	3/4/4/4	-
3	PEG	A	402	-	-	2/4/4/4	-
3	PEG	A	403	-	-	3/4/4/4	-
3	PEG	A	413	-	-	1/4/4/4	-
3	PEG	B	407	-	-	4/4/4/4	-
3	PEG	B	409	-	-	3/4/4/4	-
4	PG4	B	412	-	-	6/10/10/10	-
3	PEG	B	413	-	-	1/4/4/4	-
4	PG4	B	402	-	-	4/10/10/10	-
3	PEG	A	405	-	-	4/4/4/4	-
4	PG4	A	410	-	-	7/10/10/10	-
3	PEG	B	415	-	-	2/4/4/4	-
3	PEG	A	412	-	-	2/4/4/4	-
3	PEG	B	416	-	-	3/4/4/4	-
4	PG4	B	406	-	-	7/10/10/10	-
4	PG4	B	411	-	-	9/10/10/10	-
3	PEG	B	410	-	-	2/4/4/4	-
4	PG4	A	416	-	-	4/10/10/10	-
4	PG4	A	404	-	-	7/10/10/10	-
4	PG4	A	415	-	-	8/10/10/10	-
3	PEG	B	408	-	-	4/4/4/4	-
2	2J1	A	401	-	-	4/4/11/11	0/3/3/3
3	PEG	A	406	-	-	4/4/4/4	-

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	2J1	C12-C1	7.42	1.49	1.39
2	A	401	2J1	C3-C4	6.24	1.49	1.41
2	A	401	2J1	C5-C6	6.00	1.48	1.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	2J1	O1-C13	3.70	1.33	1.22
2	A	401	2J1	O2-C13	-2.77	1.22	1.30

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	2J1	C10-C5-C6	-6.60	123.00	125.71
2	A	401	2J1	C11-C4-C3	-4.60	118.85	122.52
2	A	401	2J1	C2-C3-C4	3.36	122.66	119.39
2	A	401	2J1	C10-C5-N	3.27	127.48	124.62
2	A	401	2J1	C7-C6-C5	3.24	124.74	122.28

There are no chirality outliers.

5 of 118 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	411	PG4	O3-C5-C6-O4
3	B	408	PEG	O1-C1-C2-O2
3	B	403	PEG	O2-C3-C4-O4
3	B	410	PEG	O1-C1-C2-O2
3	B	416	PEG	O2-C3-C4-O4

There are no ring outliers.

13 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	414	PEG	1	0
4	B	404	PG4	1	0
4	B	401	PG4	4	0
3	B	409	PEG	1	0
4	B	412	PG4	2	0
4	A	410	PG4	1	0
3	B	415	PEG	1	0
3	B	416	PEG	1	0
4	B	411	PG4	2	0
3	B	410	PEG	1	0
4	A	404	PG4	1	0
3	B	408	PEG	1	0
2	A	401	2J1	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	356/366 (97%)	0.69	20 (5%) 30 32	13, 21, 29, 37	2 (0%)
1	B	365/366 (99%)	0.69	21 (5%) 29 31	8, 20, 35, 39	2 (0%)
All	All	721/732 (98%)	0.69	41 (5%) 29 31	8, 20, 31, 39	4 (0%)

The worst 5 of 41 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	190	SER	3.4
1	B	251	ASN	3.4
1	B	253	ASP	3.3
1	A	253	ASP	3.2
1	B	211	ASP	3.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	PEG	B	416	7/7	0.50	0.23	51,54,56,56	0
4	PG4	B	411	13/13	0.52	0.27	48,50,51,52	0
4	PG4	B	401	13/13	0.55	0.22	37,38,39,39	0
3	PEG	B	408	7/7	0.56	0.24	40,40,41,41	0
3	PEG	B	405	7/7	0.56	0.22	55,56,56,56	0
3	PEG	B	403	7/7	0.60	0.19	27,27,28,28	0
3	PEG	A	414	7/7	0.61	0.22	58,59,59,59	0
4	PG4	A	415	13/13	0.61	0.20	46,47,50,50	0
3	PEG	A	408	7/7	0.64	0.17	29,30,30,30	0
2	2J1	A	401	17/17	0.66	0.20	51,51,52,53	0
3	PEG	B	410	7/7	0.66	0.20	44,44,45,45	0
3	PEG	A	418	7/7	0.66	0.18	51,52,52,52	0
4	PG4	A	404	13/13	0.68	0.17	35,36,38,38	0
3	PEG	A	412	7/7	0.70	0.19	28,29,29,29	0
4	PG4	B	406	13/13	0.70	0.20	47,50,51,51	0
3	PEG	A	405	7/7	0.70	0.20	38,38,38,38	0
3	PEG	A	403	7/7	0.71	0.19	41,41,42,42	0
3	PEG	A	413	7/7	0.72	0.15	37,37,38,38	0
4	PG4	A	410	13/13	0.72	0.16	28,29,30,30	0
3	PEG	A	402	7/7	0.75	0.16	34,34,34,34	0
3	PEG	B	409	7/7	0.78	0.14	44,45,45,45	0
3	PEG	B	413	7/7	0.79	0.19	55,55,56,56	0
3	PEG	B	414	7/7	0.80	0.14	38,39,39,40	0
4	PG4	A	416	13/13	0.82	0.13	40,41,44,44	0
3	PEG	B	407	7/7	0.83	0.11	34,34,34,34	0
4	PG4	B	412	13/13	0.83	0.13	25,27,27,28	0
4	PG4	B	404	13/13	0.84	0.13	34,35,36,36	0
4	PG4	B	402	13/13	0.85	0.13	30,31,32,32	0
5	CA	A	411	1/1	0.86	0.08	54,54,54,54	0
3	PEG	A	406	7/7	0.87	0.12	31,31,31,31	0
5	CA	A	409	1/1	0.87	0.12	58,58,58,58	0
3	PEG	B	415	7/7	0.87	0.19	21,21,21,22	0
5	CA	A	417	1/1	0.91	0.11	52,52,52,52	0
5	CA	B	417	1/1	0.92	0.10	43,43,43,43	0
5	CA	A	407	1/1	0.97	0.04	31,31,31,31	0

6.5 Other polymers ⓘ

There are no such residues in this entry.