



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 07:49 PM UTC

PDB ID : 7N9G / pdb_00007n9g
Title : Crystal structure of the Abl 1b Kinase domain in complex with Dasatinib and Imatinib
Authors : Miller, D.J.; Xie, T.
Deposited on : 2021-06-17
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	271	5% 87% 5% 8%
1	B	271	4% 93% . .
1	C	271	21% 79% 7% 14%

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 6307 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tyrosine-protein kinase ABL1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	250	Total	C	N	O	S	0	0	0
			2017	1308	328	366	15			
1	B	264	Total	C	N	O	S	0	0	0
			2109	1361	340	391	17			
1	C	233	Total	C	N	O	S	0	0	0
			1760	1124	293	331	12			

- Molecule 2 is 4-(4-METHYL-PIPERAZIN-1-YLMETHYL)-N-[4-METHYL-3-(4-PYRIDIN-3-YL-PYRIMIDIN-2-YLAMINO)-PHENYL]-BENZAMIDE (CCD ID: STI) (formula: C₂₉H₃₁N₇O) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			37	29	7	1		
2	B	1	Total	C	N	O	0	0
			37	29	7	1		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	C	1	Total	C	N	O	0	0
			37	29	7	1		

- Molecule 3 is N-(2-CHLORO-6-METHYLPHENYL)-2-({6-[4-(2-HYDROXYETHYL)PIPERAZIN-1-YL]-2-METHYLPYRIMIDIN-4-YL}AMINO)-1,3-THIAZOLE-5-CARBOXAMIDE (CCD ID: 1N1) (formula: C₂₂H₂₆ClN₇O₂S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	Cl	N	O	S	0	0
			33	22	1	7	2	1		
3	B	1	Total	C	Cl	N	O	S	0	0
			33	22	1	7	2	1		
3	C	1	Total	C	Cl	N	O	S	0	0
			33	22	1	7	2	1		

- Molecule 4 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	P	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	84	Total	O	0	0
			84	84		
5	B	94	Total	O	0	0
			94	94		
5	C	28	Total	O	0	0
			28	28		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tyrosine-protein kinase ABL1



- Molecule 1: Tyrosine-protein kinase ABL1



- Molecule 1: Tyrosine-protein kinase ABL1



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	69.93Å 166.88Å 186.82Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.70 – 2.20 46.70 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.8 (46.70-2.20) 99.9 (46.70-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.20 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.8.0257	Depositor
R, R_{free}	0.217 , 0.251 (Not available) , 0.251	Depositor DCC
R_{free} test set	2704 reflections (4.66%)	wwPDB-VP
Wilson B-factor (Å ²)	47.3	Xtriage
Anisotropy	0.376	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 31.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6307	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: STI, PO4, 1N1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.94	0/2069	1.23	0/2804
1	B	0.96	0/2163	1.28	0/2935
1	C	1.04	0/1797	1.37	0/2443
All	All	0.98	0/6029	1.29	0/8182

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2017	0	1955	10	0
1	B	2109	0	2028	13	0
1	C	1760	0	1573	10	0
2	A	37	0	31	1	0
2	B	37	0	31	1	0
2	C	37	0	31	1	0
3	A	33	0	26	2	0
3	B	33	0	26	1	0
3	C	33	0	26	2	0
4	A	5	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	84	0	0	0	0
5	B	94	0	0	0	0
5	C	28	0	0	0	0
All	All	6307	0	5727	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:346:LEU:HD21	1:B:362:MET:HE1	1.69	0.75
1:B:292:LEU:HD23	1:B:299:VAL:HA	1.75	0.69
1:C:359:LEU:HA	1:C:362:MET:HE2	1.77	0.67
1:B:359:LEU:HA	1:B:362:MET:HE2	1.77	0.66
3:A:602:1N1:N4	3:A:602:1N1:S	2.69	0.66
3:B:602:1N1:S	3:B:602:1N1:N4	2.69	0.65
1:A:346:LEU:HD21	1:A:362:MET:HE1	1.84	0.60
1:A:359:LEU:HA	1:A:362:MET:HE2	1.84	0.59
1:C:346:LEU:HD21	1:C:362:MET:HE1	1.85	0.58
1:B:292:LEU:CD2	1:B:299:VAL:HA	2.34	0.57
1:C:309:MET:HE1	3:C:602:1N1:C7	2.38	0.53
3:C:602:1N1:S	3:C:602:1N1:N4	2.81	0.53
1:B:256:MET:CE	1:B:322:GLY:HA3	2.39	0.53
1:A:380:HIS:O	1:A:440:ASP:OD1	2.27	0.52
1:C:312:ILE:HD11	1:C:400:ASP:OD2	2.09	0.52
1:C:374:GLU:HG3	1:C:437:ILE:HG12	1.93	0.51
2:A:601:STI:H151	2:A:601:STI:O29	2.11	0.51
1:B:256:MET:HE3	1:B:322:GLY:HA3	1.91	0.51
1:B:292:LEU:HD12	1:B:293:LYS:N	2.27	0.49
1:A:381:ARG:HG2	1:A:402:GLY:O	2.13	0.49
1:C:490:LEU:HD11	1:C:508:ILE:HG12	1.94	0.48
1:B:292:LEU:HD12	1:B:293:LYS:H	1.77	0.48
2:B:601:STI:O29	2:B:601:STI:H151	2.14	0.48
1:A:359:LEU:HD23	1:A:362:MET:CE	2.43	0.48
1:A:359:LEU:HD23	1:A:362:MET:HE2	1.96	0.47
1:C:276:TYR:O	1:C:289:VAL:N	2.46	0.47
1:A:420:PHE:CB	1:A:421:PRO:HD2	2.46	0.46
1:C:401:PHE:C	1:C:401:PHE:CD1	2.97	0.43
1:B:311:GLU:O	1:B:376:LYS:NZ	2.52	0.43
1:A:420:PHE:HB2	1:A:421:PRO:HD2	2.01	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:256:MET:HE3	1:B:322:GLY:CA	2.50	0.42
1:B:437:ILE:HD12	1:B:437:ILE:HA	1.96	0.42
1:A:498:ASN:HB3	1:A:501:ASP:OD2	2.20	0.42
1:C:383:LEU:HD23	1:C:440:ASP:O	2.19	0.41
1:C:442:TRP:CE3	1:C:495:TRP:HA	2.56	0.41
3:A:602:1N1:N6	4:A:603:PO4:O2	2.54	0.41
1:A:385:ALA:N	1:A:447:LEU:HD13	2.36	0.41
1:B:264:LYS:HD2	1:B:279:VAL:HG23	2.03	0.40
1:B:292:LEU:HD21	1:B:299:VAL:N	2.36	0.40
2:C:601:STI:O29	2:C:601:STI:H151	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

There are no protein backbone outliers to report in this entry.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	212/237 (90%)	210 (99%)	2 (1%)	70	84
1	B	221/237 (93%)	220 (100%)	1 (0%)	81	90
1	C	166/237 (70%)	165 (99%)	1 (1%)	78	89
All	All	599/711 (84%)	595 (99%)	4 (1%)	76	87

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	406	LEU
1	A	420	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	292	LEU
1	C	422	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	433	ASN
1	B	415	HIS
1	B	510	GLN
1	C	316	ASN
1	C	466	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	1N1	C	602	-	36,36,36	0.34	0	50,50,50	0.81	1 (2%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	1N1	B	602	-	36,36,36	0.39	0	50,50,50	0.82	2 (4%)
2	STI	A	601	-	41,41,41	0.28	0	56,56,56	0.44	0
3	1N1	A	602	-	36,36,36	0.40	0	50,50,50	0.97	3 (6%)
2	STI	B	601	-	41,41,41	0.28	0	56,56,56	0.47	0
4	PO4	A	603	-	4,4,4	0.79	0	6,6,6	0.72	0
2	STI	C	601	-	41,41,41	0.29	0	56,56,56	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	1N1	C	602	-	-	1/19/29/29	0/4/4/4
3	1N1	B	602	-	-	0/19/29/29	0/4/4/4
2	STI	A	601	-	-	1/20/30/30	0/5/5/5
3	1N1	A	602	-	-	0/19/29/29	0/4/4/4
2	STI	B	601	-	-	4/20/30/30	0/5/5/5
2	STI	C	601	-	-	8/20/30/30	0/5/5/5

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	602	1N1	C5-C4-N2	-2.41	118.27	121.12
3	A	602	1N1	N-C11-N4	2.36	119.05	113.33
3	A	602	1N1	C14-N3-C13	2.30	118.59	116.05
3	C	602	1N1	N-C11-N4	2.23	118.72	113.33
3	B	602	1N1	C14-N3-C13	2.12	118.39	116.05
3	B	602	1N1	C14-N4-C11	2.02	118.20	115.67

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	601	STI	C19-C14-N13-C9
2	B	601	STI	C15-C14-N13-C9
2	C	601	STI	C19-C14-N13-C9
2	C	601	STI	C15-C14-N13-C9
2	B	601	STI	C27-C46-N48-C49

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	B	601	STI	C27-C46-N48-C53
2	C	601	STI	C27-C46-N48-C53
2	C	601	STI	C27-C46-N48-C49
2	C	601	STI	N21-C22-C23-C25
2	C	601	STI	O29-C22-C23-C25
2	C	601	STI	N21-C22-C23-C29
2	A	601	STI	C15-C14-N13-C9
2	C	601	STI	O29-C22-C23-C29
3	C	602	1N1	N4-C11-N-C


There are no ring outliers.

7 monomers are involved in 8 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	602	1N1	2	0
3	B	602	1N1	1	0
2	A	601	STI	1	0
3	A	602	1N1	2	0
2	B	601	STI	1	0
4	A	603	PO4	1	0
2	C	601	STI	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand 1N1 C 602




Bond lengths



Bond angles




Torsions




Rings


Ligand 1N1 B 602




Bond lengths



Bond angles




Torsions




Rings


Ligand STI A 601




Bond lengths



Bond angles




Torsions




Rings


Ligand 1N1 A 602




Bond lengths



Bond angles




Torsions




Rings


Ligand STI B 601




Bond lengths




Bond angles



Torsions



Rings



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	250/271 (92%)	0.33	13 (5%) 33 29	35, 48, 78, 110	0
1	B	264/271 (97%)	0.42	12 (4%) 38 35	37, 50, 74, 102	0
1	C	233/271 (85%)	1.23	57 (24%) 2 1	43, 68, 112, 125	0
All	All	747/813 (91%)	0.65	82 (10%) 10 8	35, 54, 98, 125	0

All (82) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	272	TYR	6.5
1	C	435	PHE	5.6
1	C	280	TRP	5.5
1	C	291	THR	5.2
1	A	406	LEU	5.1
1	C	378	PHE	5.1
1	A	293	LYS	5.0
1	A	297	MET	5.0
1	A	435	PHE	4.9
1	C	267	LEU	4.9
1	C	414	ALA	4.7
1	C	401	PHE	4.7
1	C	298	GLU	4.3
1	C	265	HIS	4.3
1	B	417	GLY	4.3
1	C	420	PHE	4.3
1	B	269	GLY	4.2
1	B	294	GLU	4.2
1	C	306	ALA	4.0
1	C	275	VAL	3.9
1	A	434	LYS	3.9
1	C	332	ILE	3.8
1	C	412	TYR	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	321	LEU	3.5
1	C	400	ASP	3.5
1	C	260	ASP	3.5
1	B	410	ASP	3.4
1	B	293	LYS	3.3
1	A	432	TYR	3.3
1	C	261	ILE	3.3
1	C	408	THR	3.2
1	C	276	TYR	3.2
1	C	518	GLU	3.2
1	C	256	MET	3.2
1	A	268	GLY	3.1
1	C	279	VAL	3.1
1	C	407	MET	3.0
1	A	517	GLN	3.0
1	C	302	PHE	3.0
1	C	254	TRP	3.0
1	A	420	PHE	2.9
1	C	290	LYS	2.9
1	C	402	GLY	2.8
1	C	286	THR	2.8
1	C	406	LEU	2.8
1	C	255	GLU	2.8
1	A	298	GLU	2.7
1	A	292	LEU	2.7
1	B	517	GLN	2.7
1	A	456	MET	2.6
1	B	292	LEU	2.6
1	C	259	THR	2.6
1	C	413	THR	2.6
1	C	264	LYS	2.6
1	C	320	LEU	2.5
1	C	379	ILE	2.5
1	C	312	ILE	2.5
1	B	418	ALA	2.5
1	A	252	ASP	2.5
1	C	333	ILE	2.4
1	C	300	GLU	2.4
1	C	305	GLU	2.4
1	B	463	ASP	2.3
1	C	299	VAL	2.3
1	B	408	THR	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	287	VAL	2.3
1	C	258	ARG	2.3
1	C	375	LYS	2.2
1	C	289	VAL	2.2
1	C	301	GLU	2.2
1	C	399	ALA	2.2
1	B	409	GLY	2.2
1	C	411	THR	2.2
1	C	437	ILE	2.2
1	C	318	VAL	2.1
1	C	505	PHE	2.1
1	C	338	THR	2.1
1	C	263	MET	2.1
1	C	382	ASP	2.1
1	C	262	THR	2.0
1	C	309	MET	2.0
1	C	288	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	1N1	C	602	33/33	0.82	0.15	69,82,101,103	0
2	STI	C	601	37/37	0.87	0.16	54,60,119,121	0
2	STI	A	601	37/37	0.91	0.12	40,47,83,88	0
2	STI	B	601	37/37	0.91	0.13	39,50,106,108	0
3	1N1	B	602	33/33	0.95	0.08	35,40,47,57	0
3	1N1	A	602	33/33	0.97	0.07	36,40,48,59	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	PO4	A	603	5/5	0.98	0.06	38,40,46,47	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 1N1 C 602:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around STI C 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around STI A 601:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around STI B 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 1N1 B 602:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 1N1 A 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.