



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2026 – 06:41 PM UTC

PDB ID : 1NAS / pdb_00001nas
Title : SEPIAPTERIN REDUCTASE COMPLEXED WITH N-ACETYL SEROTONIN
Authors : Auerbach, G.; Herrmann, A.; Bacher, A.; Huber, R.
Deposited on : 1998-03-26
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	259	 86% 12% •

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 2420 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SEPIAPTERIN REDUCTASE.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	259	1935	1211	339	374	11	0	0	0

- Molecule 2 is OXALOACETATE ION (CCD ID: OAA) (formula: $C_4H_3O_5$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	9	4	5	0	0

- Molecule 3 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (CCD ID: NAP) (formula: $C_{21}H_{28}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 48	C 21	N 7	O 17	P 3	0	0

- Molecule 4 is N-ACETYL SEROTONIN (CCD ID: ASE) (formula: $C_{12}H_{14}N_2O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			16	12	2	2		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	412	Total 412	O 412	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: SEPIAPTERIN REDUCTASE



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	116.96Å 116.96Å 105.21Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	8.00 – 2.10	Depositor
% Data completeness (in resolution range)	(Not available) (8.00-2.10)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.166 , 0.202	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2420	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAP, OAA, ASE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.44	0/1961	0.92	7/2655 (0.3%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	110	LYS	N-CA-C	9.03	124.43	113.41
1	A	95	ARG	N-CA-C	6.25	118.80	108.99
1	A	199	PRO	N-CA-C	5.21	120.45	114.20
1	A	159	LEU	N-CA-C	-5.10	106.16	112.38
1	A	216	ASP	N-CA-C	5.04	116.74	109.48
1	A	216	ASP	CA-C-N	5.02	124.19	118.97
1	A	216	ASP	C-N-CA	5.02	124.19	118.97

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1935	0	1973	29	3
2	A	9	0	2	2	0
3	A	48	0	25	2	0
4	A	16	0	14	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	412	0	0	14	7
All	All	2420	0	2014	31	7

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:190:PRO:CB	5:A:1205:HOH:O	1.91	1.16
1:A:190:PRO:CG	5:A:1205:HOH:O	2.08	0.96
1:A:190:PRO:HG3	5:A:1205:HOH:O	1.64	0.95
1:A:190:PRO:CA	5:A:1205:HOH:O	2.15	0.91
2:A:1:OAA:H21	4:A:801:ASE:HN3	1.40	0.84
1:A:190:PRO:HB3	5:A:1205:HOH:O	1.66	0.81
1:A:219:LEU:HD11	4:A:801:ASE:H151	1.63	0.80
1:A:150:SER:HB3	5:A:1177:HOH:O	1.89	0.70
1:A:8:CYS:H	1:A:94:GLN:HE21	1.39	0.69
2:A:1:OAA:H21	4:A:801:ASE:N13	2.07	0.69
1:A:52:LYS:HB2	1:A:66:LEU:HD21	1.75	0.68
1:A:155:ASN:HD21	1:A:178:ARG:HE	1.44	0.65
1:A:190:PRO:HA	5:A:1205:HOH:O	1.87	0.63
1:A:8:CYS:H	1:A:94:GLN:NE2	1.98	0.60
1:A:104:THR:H	1:A:128:ASN:HD21	1.50	0.59
1:A:107:ASP:HB3	1:A:110:LYS:NZ	2.22	0.54
1:A:43:ARG:HB2	3:A:800:NAP:O2X	2.09	0.51
1:A:91:GLU:HG3	1:A:92:GLY:H	1.77	0.50
1:A:175:LYS:HD2	5:A:864:HOH:O	2.11	0.49
1:A:215:LYS:HA	5:A:1103:HOH:O	2.11	0.49
1:A:9:ALA:HB2	1:A:95:ARG:HG2	1.95	0.49
1:A:157:SER:O	3:A:800:NAP:H6N	2.13	0.49
1:A:155:ASN:ND2	1:A:178:ARG:HE	2.10	0.48
1:A:18:ARG:NH1	5:A:1010:HOH:O	2.47	0.47
1:A:241:LYS:NZ	5:A:1071:HOH:O	2.48	0.46
1:A:73:THR:HG22	5:A:1068:HOH:O	2.16	0.46
1:A:107:ASP:HB3	1:A:110:LYS:HZ1	1.81	0.45
1:A:110:LYS:HB3	5:A:1116:HOH:O	2.16	0.44
1:A:30:ARG:NH1	5:A:855:HOH:O	2.52	0.43
1:A:90:PRO:HD2	1:A:93:LEU:HD13	2.04	0.40
1:A:42:ALA:O	1:A:68:ALA:HA	2.20	0.40

All (7) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:5:GLY:CA	5:A:1207:HOH:O[12_564]	0.46	1.74
5:A:1213:HOH:O	5:A:1213:HOH:O[9_555]	0.82	1.38
1:A:5:GLY:C	5:A:1207:HOH:O[12_564]	1.44	0.76
1:A:5:GLY:N	5:A:1207:HOH:O[12_564]	1.51	0.69
5:A:1202:HOH:O	5:A:1202:HOH:O[10_665]	2.04	0.16
5:A:1107:HOH:O	5:A:1211:HOH:O[10_665]	2.05	0.15
5:A:881:HOH:O	5:A:1204:HOH:O[9_555]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	257/259 (99%)	250 (97%)	5 (2%)	2 (1%)	16 12

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	91	GLU
1	A	147	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	208/208 (100%)	206 (99%)	2 (1%)	68 76

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	155	ASN
1	A	248	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	58	GLN
1	A	59	GLN
1	A	94	GLN
1	A	100	ASN
1	A	116	ASN
1	A	122	ASN
1	A	123	ASN
1	A	128	ASN
1	A	141	ASN
1	A	155	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	OAA	A	1	-	8,8,8	4.59	1 (12%)	8,10,10	1.66	3 (37%)
4	ASE	A	801	-	17,17,17	1.87	4 (23%)	23,23,23	2.76	6 (26%)
3	NAP	A	800	-	50,52,52	1.50	7 (14%)	71,80,80	1.77	13 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	OAA	A	1	-	-	0/8/8/8	-
4	ASE	A	801	-	-	2/6/6/6	0/2/2/2
3	NAP	A	800	-	-	6/35/67/67	0/5/5/5

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1	OAA	C3-C4	-12.43	1.34	1.53
3	A	800	NAP	C2N-N1N	5.96	1.41	1.35
4	A	801	ASE	C11-C7	4.24	1.58	1.50
3	A	800	NAP	C5A-N7A	-3.93	1.31	1.39
3	A	800	NAP	P2B-O2B	3.77	1.66	1.59
4	A	801	ASE	C6-N9	3.75	1.43	1.37
4	A	801	ASE	C14-N13	2.30	1.40	1.34
3	A	800	NAP	C4A-N9A	-2.28	1.32	1.37
3	A	800	NAP	O4D-C1D	2.20	1.43	1.40
4	A	801	ASE	C2-C3	2.19	1.43	1.39
3	A	800	NAP	C6N-N1N	2.16	1.40	1.35
3	A	800	NAP	C3N-C7N	2.10	1.53	1.50

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	801	ASE	C12-N13-C14	-10.84	106.55	122.56
3	A	800	NAP	N3A-C4A-N9A	5.79	137.01	127.17
3	A	800	NAP	C5A-C4A-N3A	-5.64	118.95	126.72
3	A	800	NAP	C4D-O4D-C1D	-4.50	105.80	109.92
3	A	800	NAP	N3A-C2A-N1A	-4.49	121.78	128.58

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	800	NAP	C6A-C5A-N7A	-3.52	125.31	132.09
4	A	801	ASE	C3-C7-C8	3.31	109.54	106.16
4	A	801	ASE	O10-C1-C4	-3.25	110.96	120.00
3	A	800	NAP	C2A-N3A-C4A	3.23	119.73	111.83
3	A	800	NAP	C6A-C5A-C4A	3.08	121.39	117.18
4	A	801	ASE	O10-C1-C2	2.88	127.40	119.85
3	A	800	NAP	C4A-N9A-C8A	2.80	108.68	105.74
2	A	1	OAA	O3-C3-C2	2.54	124.32	120.41
3	A	800	NAP	N6A-C6A-N1A	2.52	123.99	118.38
2	A	1	OAA	O4-C4-C3	-2.36	118.88	121.81
3	A	800	NAP	N9A-C8A-N7A	-2.29	110.69	113.94
3	A	800	NAP	C4A-C5A-N7A	2.24	113.14	110.58
2	A	1	OAA	O5-C4-C3	2.17	119.62	113.59
4	A	801	ASE	C5-C4-C1	2.16	122.16	119.88
3	A	800	NAP	O3X-P2B-O2X	2.15	115.85	107.80
4	A	801	ASE	C15-C14-N13	2.07	119.65	116.12
3	A	800	NAP	O4B-C1B-N9A	2.06	112.04	108.09

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	800	NAP	C5D-O5D-PN-O3
4	A	801	ASE	O16-C14-N13-C12
4	A	801	ASE	C15-C14-N13-C12
3	A	800	NAP	PA-O3-PN-O2N
3	A	800	NAP	C5D-O5D-PN-O1N
3	A	800	NAP	O4B-C4B-C5B-O5B
3	A	800	NAP	PA-O3-PN-O1N
3	A	800	NAP	C2B-O2B-P2B-O1X


There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1	OAA	2	0
4	A	801	ASE	3	0
3	A	800	NAP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In

addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.