



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 07:38 AM UTC

PDB ID : 3NA1 / pdb_00003na1
Title : Crystal structure of human CYP11A1 in complex with 20-hydroxycholesterol
Authors : Strushkevich, N.V.; MacKenzie, F.; Tempel, W.; Botchkarev, A.; Arrowsmith, C.H.; Edwards, A.M.; Bountra, C.; Weigelt, J.U.; Park, H.; Structural Genomics Consortium (SGC)
Deposited on : 2010-05-31
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1898 (2.26-2.26)
Clashscore	190562	2005 (2.26-2.26)
Ramachandran outliers	187476	1965 (2.26-2.26)
Sidechain outliers	187428	1966 (2.26-2.26)
RSRZ outliers	180081	1898 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	487	
1	B	487	
2	C	123	
2	D	123	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 8795 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cholesterol side-chain cleavage enzyme, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	472	Total	C	N	O	S	0	0	0
			3886	2518	669	683	16			
1	B	471	Total	C	N	O	S	0	0	0
			3880	2515	668	681	16			


There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	483	HIS	-	expression tag	UNP P05108
A	484	HIS	-	expression tag	UNP P05108
A	485	HIS	-	expression tag	UNP P05108
A	486	HIS	-	expression tag	UNP P05108
A	487	HIS	-	expression tag	UNP P05108
A	488	HIS	-	expression tag	UNP P05108
B	483	HIS	-	expression tag	UNP P05108
B	484	HIS	-	expression tag	UNP P05108
B	485	HIS	-	expression tag	UNP P05108
B	486	HIS	-	expression tag	UNP P05108
B	487	HIS	-	expression tag	UNP P05108
B	488	HIS	-	expression tag	UNP P05108

- Molecule 2 is a protein called Adrenodoxin, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	26	Total	C	N	O	S	0	0	0
			179	103	28	43	5			
2	D	24	Total	C	N	O	S	0	0	0
			167	97	26	39	5			


- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



-
- The chemical structure of Hedericholide (HCD) is shown with atom numbering and stereochemistry. The molecule features a complex polycyclic core with a side chain. Key stereochemical features include:
- C10(R):** Indicated by a wedge bond to the C9-C10 bond.
 - C14(S):** Indicated by a wedge bond to the C13-C14 bond.
 - C17(S):** Indicated by a wedge bond to the C16-C17 bond.
 - C20(S):** Indicated by a wedge bond to the C19-C20 bond.
 - C3(S):** Indicated by a wedge bond to the C2-C3 bond.
 - C8(S):** Indicated by a wedge bond to the C7-C8 bond.
 - C9(S):** Indicated by a wedge bond to the C8-C9 bond.
 - C11(R):** Indicated by a wedge bond to the C10-C11 bond.
 - C13(S):** Indicated by a wedge bond to the C12-C13 bond.
 - C15(S):** Indicated by a wedge bond to the C14-C15 bond.
 - C16(S):** Indicated by a wedge bond to the C15-C16 bond.
 - C18(S):** Indicated by a wedge bond to the C17-C18 bond.
 - C19(S):** Indicated by a wedge bond to the C18-C19 bond.
 - C21(S):** Indicated by a wedge bond to the C20-C21 bond.
 - C22(S):** Indicated by a wedge bond to the C21-C22 bond.
 - C23(S):** Indicated by a wedge bond to the C22-C23 bond.
 - C24(S):** Indicated by a wedge bond to the C23-C24 bond.
 - C25(S):** Indicated by a wedge bond to the C24-C25 bond.
 - C26(S):** Indicated by a wedge bond to the C25-C26 bond.
 - C27(S):** Indicated by a wedge bond to the C26-C27 bond.
- The structure also shows a double bond between C5 and C6, and a double bond between C12 and C13. The stereochemistry is defined by the relative configuration of the chiral centers.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total 29	C 27	O 2	0	0
4	B	1	Total 29	C 27	O 2	0	0

- Molecule 5 is FE2/S2 (INORGANIC) CLUSTER (CCD ID: FES) (formula: Fe_2S_2).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	Fe	S	0	0
			4	2	2		
5	D	1	Total	Fe	S	0	0
			4	2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	261	Total	O	0	0
			261	261		
6	B	261	Total	O	0	0
			261	261		
6	C	5	Total	O	0	0
			5	5		
6	D	4	Total	O	0	0
			4	4		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cholesterol side-chain cleavage enzyme, mitochondrial



- Molecule 1: Cholesterol side-chain cleavage enzyme, mitochondrial



- Molecule 2: Adrenodoxin, mitochondrial



LYS

THR

SER

● Molecule 2: Adrenodoxin, mitochondrial



SER	SER	GLU	ASP	LYS	THR	VAL	HIS	PHE	ILE	ASN	ARG	ASP	GLY	THR	LEU	THR	THR	LYS	GLY	LYS	VAL	GLY	ASP	SER	LEU	VAL	VAL	VAL	GLU	ASN	LYS	ASP	ILE	GLY	PHE	GLY	GLY	A45	C46	E47	G48	T49	L50	A51	C52	S53	T54	C55	HIS	LEU	ILE	PHE	GLU	ASP
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

HIS	ILE	TYR	GLU	LYS	LEU	ASP	ALA	THR	ASP	E73	E74	N75	D76	M77	L78	D79	L80	A81	TYR	GLY	LEU	THR	ASP	ARG	SER	ARG	L90	G91	C92	Q93	ILE	CYS	LEU	THR	LYS	SER	MET	ASN	MET	THR	VAL	ARG	VAL	PRO	GLU	THR	VAL	ALA	ASP	ALA	ARG	GLN	SER	ILE	PHE	ASP	VAL	GLY
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

LYS

THR

SER

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	82.81Å 115.15Å 85.73Å 90.00° 101.45° 90.00°	Depositor
Resolution (Å)	50.00 – 2.25 50.00 – 2.25	Depositor EDS
% Data completeness (in resolution range)	98.9 (50.00-2.25) 98.9 (50.00-2.25)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.69 (at 2.24Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.193 , 0.232 0.193 , 0.229	Depositor DCC
R_{free} test set	3714 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	24.1	Xtriage
Anisotropy	0.031	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 31.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.170 for l,-k,h	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8795	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, FES, HCD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.56	0/3996	0.80	2/5423 (0.0%)
1	B	0.56	1/3990 (0.0%)	0.82	3/5414 (0.1%)
2	C	0.51	0/176	0.83	0/233
2	D	0.57	0/164	0.69	0/217
All	All	0.56	1/8326 (0.0%)	0.81	5/11287 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	206	VAL	CA-CB	5.33	1.57	1.54

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	209	LEU	N-CA-C	8.46	120.12	111.07
1	B	353	VAL	CB-CA-C	-7.40	102.15	112.14
1	A	209	LEU	N-CA-C	7.30	119.87	111.11
1	A	216	PHE	CB-CA-C	-5.93	101.49	110.92
1	B	216	PHE	CB-CA-C	-5.65	101.94	110.92

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3886	0	3852	34	0
1	B	3880	0	3848	49	0
2	C	179	0	153	5	0
2	D	167	0	147	6	0
3	A	43	0	30	2	0
3	B	43	0	30	2	0
4	A	29	0	46	0	0
4	B	29	0	46	0	0
5	C	4	0	0	0	0
5	D	4	0	0	0	0
6	A	261	0	0	1	0
6	B	261	0	0	1	0
6	C	5	0	0	0	0
6	D	4	0	0	0	0
All	All	8795	0	8152	88	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (88) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:87:TRP:NE1	1:B:201:MET:HE1	1.63	1.10
1:B:87:TRP:HE1	1:B:201:MET:HE1	1.28	0.94
1:A:310:GLN:HE21	1:A:314:ARG:HH22	1.15	0.90
1:B:310:GLN:HE21	1:B:314:ARG:HH22	1.17	0.87
1:B:87:TRP:NE1	1:B:201:MET:CE	2.37	0.87
1:B:116:ASN:HB3	2:D:49:THR:HG21	1.55	0.86
1:B:310:GLN:HE22	1:B:446:VAL:H	1.24	0.83
1:A:310:GLN:HE22	1:A:446:VAL:H	1.25	0.81
1:B:418:TRP:HE1	2:D:80:LEU:HB3	1.44	0.80
1:B:201:MET:O	1:B:201:MET:HE2	1.84	0.77
1:B:389:PHE:O	1:B:412:ARG:NH1	2.20	0.74
1:B:87:TRP:CE2	1:B:201:MET:HE1	2.23	0.73
1:B:201:MET:CE	1:B:227:HIS:HE1	2.03	0.71
1:A:418:TRP:HE1	2:C:80:LEU:HB3	1.56	0.70
1:B:87:TRP:HE1	1:B:201:MET:CE	2.01	0.67
1:B:380:ILE:HD11	1:B:414:LEU:HD13	1.78	0.65
1:A:344:GLU:CD	1:A:347:ARG:NH1	2.55	0.63
1:A:14:SER:HB3	1:A:51:ARG:O	1.99	0.63
1:A:272:MET:HE2	1:A:276:ASP:HB3	1.84	0.60
1:B:201:MET:HE2	1:B:227:HIS:HE1	1.67	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:294:MET:HA	1:A:294:MET:HE2	1.85	0.59
1:B:272:MET:HE1	1:B:280:ASN:HD22	1.67	0.59
1:A:272:MET:HE1	1:A:280:ASN:HD22	1.68	0.59
1:A:422:GLN:NE2	1:A:426:ARG:HH11	2.01	0.58
1:B:294:MET:HA	1:B:294:MET:HE2	1.85	0.57
1:B:201:MET:HE3	1:B:227:HIS:CE1	2.40	0.56
3:A:601:HEM:HBB2	3:A:601:HEM:HMB2	1.88	0.55
1:B:201:MET:HE3	1:B:227:HIS:HE1	1.72	0.55
1:B:475:PHE:O	1:B:476:ASN:HB2	2.07	0.55
1:B:201:MET:CE	1:B:227:HIS:CE1	2.88	0.54
1:A:272:MET:HE1	1:A:280:ASN:ND2	2.23	0.54
1:A:418:TRP:NE1	2:C:80:LEU:HB3	2.23	0.53
1:B:257:HIS:HB2	6:B:591:HOH:O	2.08	0.53
1:A:378:VAL:HG12	1:A:380:ILE:HD12	1.90	0.53
1:A:422:GLN:HE21	1:A:426:ARG:HD2	1.73	0.52
1:A:380:ILE:HD11	1:A:414:LEU:HD13	1.91	0.52
1:B:312:MET:HE2	1:B:337:LEU:HD21	1.91	0.52
1:B:378:VAL:HG12	1:B:380:ILE:HD12	1.93	0.51
1:A:411:PHE:HA	2:C:80:LEU:CD2	2.42	0.50
1:B:256:HIS:CG	1:B:260:ARG:HD2	2.47	0.50
1:B:299:HIS:HE1	1:B:341:SER:OG	1.94	0.49
1:B:335:VAL:O	1:B:335:VAL:HG13	2.12	0.49
1:B:418:TRP:CD1	2:D:80:LEU:HD13	2.46	0.49
1:A:5:SER:HB2	1:A:6:PRO:HD2	1.95	0.49
1:B:418:TRP:NE1	2:D:80:LEU:HB3	2.21	0.48
1:A:100:VAL:HG22	1:A:108:TRP:CD1	2.48	0.48
1:A:335:VAL:HG13	1:A:335:VAL:O	2.14	0.48
1:B:310:GLN:NE2	1:B:314:ARG:HH22	2.00	0.47
1:A:312:MET:HE2	1:A:337:LEU:HD21	1.95	0.47
1:B:156:SER:HA	1:B:470:PHE:O	2.14	0.47
1:B:194:PHE:CE2	1:B:198:ILE:HD11	2.49	0.47
1:B:390:PHE:HZ	1:B:412:ARG:HB3	1.79	0.47
1:B:418:TRP:HD1	2:D:80:LEU:HD13	1.79	0.47
1:B:217:ARG:O	1:B:221:THR:OG1	2.33	0.47
1:A:372:ALA:O	1:A:373:LYS:HB2	2.15	0.47
1:A:390:PHE:HZ	1:A:412:ARG:HB3	1.80	0.46
1:B:380:ILE:HD11	1:B:414:LEU:CD1	2.45	0.46
3:A:601:HEM:HBB2	3:A:601:HEM:CMB	2.46	0.46
1:B:463:MET:HB2	1:B:464:PRO:HD2	1.97	0.46
1:A:169:GLU:HG2	1:A:182:MET:HB2	1.98	0.46
1:B:390:PHE:CZ	1:B:412:ARG:HB3	2.51	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:390:PHE:CZ	1:A:412:ARG:HB3	2.52	0.45
1:A:156:SER:HA	1:A:470:PHE:O	2.17	0.45
3:B:601:HEM:HBB2	3:B:601:HEM:HMB2	1.99	0.45
2:D:74:GLU:HG3	2:D:90:LEU:HD12	2.00	0.44
1:B:53:LYS:HA	1:B:58:GLU:HA	2.00	0.44
1:B:310:GLN:HE21	1:B:314:ARG:NH2	2.00	0.43
1:A:310:GLN:NE2	1:A:314:ARG:HH22	1.98	0.43
1:B:335:VAL:HG22	1:B:338:LEU:HB2	2.00	0.43
1:A:310:GLN:NE2	1:A:446:VAL:H	2.04	0.43
1:B:100:VAL:HG22	1:B:108:TRP:CD1	2.54	0.43
1:A:389:PHE:O	1:A:412:ARG:NH1	2.53	0.42
1:A:109:LYS:NZ	2:C:45:ALA:O	2.43	0.42
1:A:411:PHE:HA	2:C:80:LEU:HD23	2.01	0.42
1:B:448:ILE:HD12	1:B:452:SER:HB3	2.02	0.41
1:B:188:ASN:HA	1:B:189:PRO:HD3	1.97	0.41
1:B:201:MET:HE2	1:B:227:HIS:CE1	2.52	0.41
1:B:324:ALA:HB1	1:B:330:THR:HB	2.01	0.41
1:A:239:ASP:CG	1:A:278:LYS:HE3	2.45	0.41
1:B:463:MET:HB2	1:B:464:PRO:CD	2.51	0.41
1:A:344:GLU:CD	1:A:347:ARG:HH12	2.29	0.41
1:B:90:TYR:CZ	1:B:231:TRP:HB3	2.56	0.41
1:B:272:MET:HE1	1:B:280:ASN:ND2	2.35	0.40
1:A:53:LYS:HA	1:A:58:GLU:HA	2.03	0.40
1:A:269:ASP:HB3	6:A:662:HOH:O	2.20	0.40
1:B:72:PHE:HB3	1:B:418:TRP:CE3	2.57	0.40
3:B:601:HEM:HBB2	3:B:601:HEM:CMB	2.51	0.40
1:A:310:GLN:HE21	1:A:314:ARG:NH2	1.99	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	470/487 (96%)	460 (98%)	9 (2%)	1 (0%)	43	50
1	B	469/487 (96%)	455 (97%)	11 (2%)	3 (1%)	21	21
2	C	20/123 (16%)	14 (70%)	6 (30%)	0	100	100
2	D	18/123 (15%)	13 (72%)	5 (28%)	0	100	100
All	All	977/1220 (80%)	942 (96%)	31 (3%)	4 (0%)	30	31

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	451	LEU
1	B	452	SER
1	A	105	SER
1	B	254	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	420/434 (97%)	409 (97%)	11 (3%)	40	50
1	B	419/434 (96%)	403 (96%)	16 (4%)	29	36
2	C	20/108 (18%)	20 (100%)	0	100	100
2	D	19/108 (18%)	18 (95%)	1 (5%)	20	22
All	All	878/1084 (81%)	850 (97%)	28 (3%)	34	43

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	ARG
1	A	14	SER
1	A	184	GLU
1	A	187	VAL
1	A	335	VAL
1	A	351	ILE
1	A	353	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	369	MET
1	A	380	ILE
1	A	403	SER
1	A	460	LEU
1	B	62	VAL
1	B	73	LYS
1	B	119	VAL
1	B	174	VAL
1	B	187	VAL
1	B	201	MET
1	B	239	ASP
1	B	278	LYS
1	B	335	VAL
1	B	338	LEU
1	B	351	ILE
1	B	353	VAL
1	B	367	ASP
1	B	380	ILE
1	B	452	SER
1	B	460	LEU
2	D	49	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	38	HIS
1	A	78	ASN
1	A	310	GLN
1	A	333	GLN
1	A	422	GLN
1	B	25	HIS
1	B	154	ASN
1	B	299	HIS
1	B	310	GLN
1	B	377	GLN
2	C	75	ASN
2	D	93	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	FES	D	151	2	0,4,4	-	-	-		
4	HCD	B	602	-	32,32,32	0.56	0	49,51,51	0.94	2 (4%)
3	HEM	B	601	6,1	50,50,50	1.86	11 (22%)	67,82,82	1.33	7 (10%)
4	HCD	A	602	-	32,32,32	0.53	0	49,51,51	0.92	3 (6%)
5	FES	C	150	2	0,4,4	-	-	-		
3	HEM	A	601	6,1	50,50,50	1.90	9 (18%)	67,82,82	1.38	9 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	FES	D	151	2	-	-	0/1/1/1
4	HCD	B	602	-	-	0/13/71/71	0/4/4/4
3	HEM	B	601	6,1	-	4/14/54/54	-
4	HCD	A	602	-	-	0/13/71/71	0/4/4/4
5	FES	C	150	2	-	-	0/1/1/1
3	HEM	A	601	6,1	-	3/14/54/54	-

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	601	HEM	C3D-C2D	7.62	1.53	1.36
3	A	601	HEM	C3D-C2D	7.34	1.52	1.36
3	A	601	HEM	FE-ND	5.00	2.10	1.94
3	A	601	HEM	FE-NC	4.90	2.11	1.95
3	B	601	HEM	FE-ND	4.63	2.09	1.94
3	B	601	HEM	FE-NC	3.83	2.07	1.95
3	A	601	HEM	FE-NB	3.55	2.05	1.94
3	B	601	HEM	FE-NA	3.30	2.06	1.95
3	A	601	HEM	CAB-C3B	2.97	1.55	1.47
3	B	601	HEM	CAB-C3B	2.85	1.55	1.47
3	B	601	HEM	FE-NB	2.71	2.03	1.94
3	A	601	HEM	CAC-C3C	2.65	1.54	1.47
3	B	601	HEM	CAC-C3C	2.60	1.54	1.47
3	A	601	HEM	CMC-C2C	2.54	1.56	1.50
3	B	601	HEM	CMC-C2C	2.42	1.55	1.50
3	A	601	HEM	CMB-C2B	2.40	1.55	1.50
3	B	601	HEM	CMB-C2B	2.17	1.55	1.50
3	A	601	HEM	CMA-C3A	2.16	1.55	1.50
3	B	601	HEM	CMA-C3A	2.16	1.55	1.50
3	B	601	HEM	CMD-C2D	2.12	1.55	1.50

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	601	HEM	C4D-ND-C1D	5.47	111.69	105.21
3	A	601	HEM	C4D-ND-C1D	5.34	111.53	105.21
3	A	601	HEM	CBD-CAD-C3D	-3.06	104.06	112.53
4	B	602	HCD	C4-C5-C10	2.81	120.02	116.42
4	A	602	HCD	C4-C5-C10	2.74	119.93	116.42
3	B	601	HEM	CBD-CAD-C3D	-2.72	105.02	112.53
3	B	601	HEM	CBA-CAA-C2A	-2.53	105.54	112.53
3	A	601	HEM	CAD-C3D-C4D	2.37	128.83	124.70
3	A	601	HEM	CHD-C1D-ND	2.35	126.95	124.42
3	B	601	HEM	CHD-C1D-ND	2.31	126.91	124.42
3	A	601	HEM	C4C-C3C-C2C	2.30	108.81	106.81
4	A	602	HCD	C4-C5-C6	-2.29	117.46	120.57
3	A	601	HEM	CBA-CAA-C2A	-2.27	106.27	112.53
4	B	602	HCD	C4-C5-C6	-2.22	117.56	120.57
3	B	601	HEM	O2D-CGD-CBD	2.19	120.93	114.00
4	A	602	HCD	C3-C4-C5	-2.16	108.61	112.05
3	A	601	HEM	C2A-C1A-NA	-2.10	107.82	110.15
3	B	601	HEM	C1B-NB-C4B	2.08	107.68	105.21
3	A	601	HEM	CMD-C2D-C1D	2.07	128.26	125.03

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	601	HEM	C1B-NB-C4B	2.04	107.63	105.21
3	B	601	HEM	CHC-C4B-NB	2.02	126.59	124.42

There are no chirality outliers.

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	601	HEM	C2C-C3C-CAC-CBC
3	B	601	HEM	C2C-C3C-CAC-CBC
3	A	601	HEM	C4C-C3C-CAC-CBC
3	B	601	HEM	C4C-C3C-CAC-CBC
3	B	601	HEM	CAD-CBD-CGD-O2D
3	A	601	HEM	CAD-CBD-CGD-O2D
3	B	601	HEM	CAD-CBD-CGD-O1D


There are no ring outliers.


2 monomers are involved in 4 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	601	HEM	2	0
3	A	601	HEM	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	472/487 (96%)	-0.10	11 (2%) 61 61	13, 22, 37, 47	0
1	B	471/487 (96%)	-0.06	14 (2%) 52 53	12, 22, 37, 47	0
2	C	26/123 (21%)	4.00	25 (96%) 0 0	78, 81, 98, 99	0
2	D	24/123 (19%)	4.37	24 (100%) 0 0	79, 92, 97, 98	0
All	All	993/1220 (81%)	0.13	74 (7%) 20 18	12, 22, 47, 99	0

All (74) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	78	LEU	8.4
2	C	78	LEU	8.3
2	D	81	ALA	7.5
2	D	51	ALA	6.3
2	C	90	LEU	6.3
2	C	81	ALA	6.2
1	B	451	LEU	6.0
2	C	80	LEU	6.0
2	D	45	ALA	5.8
2	D	73	GLU	5.5
1	B	254	SER	5.5
2	C	51	ALA	5.3
2	C	50	LEU	5.0
2	D	46	CYS	5.0
2	C	93	GLN	4.9
2	D	90	LEU	4.8
2	C	45	ALA	4.8
2	C	92	CYS	4.7
2	D	76	ASP	4.6
2	D	80	LEU	4.6
2	D	79	ASP	4.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	50	LEU	4.6
2	D	92	CYS	4.5
2	C	54	THR	4.4
2	C	44	GLY	4.3
2	D	49	THR	4.2
2	D	54	THR	4.1
2	C	49	THR	4.0
2	C	46	CYS	4.0
2	D	53	SER	3.6
2	D	77	MET	3.6
2	C	48	GLY	3.5
1	A	269	ASP	3.4
1	A	254	SER	3.4
2	C	55	CYS	3.4
1	B	450	HIS	3.3
1	B	407	ASN	3.3
1	A	404	LYS	3.2
1	A	476	ASN	3.2
2	D	48	GLY	3.2
1	B	257	HIS	3.2
2	C	75	ASN	3.2
2	D	93	GLN	3.1
2	D	52	CYS	3.0
2	C	53	SER	3.0
2	C	47	GLU	3.0
2	C	73	GLU	3.0
2	D	75	ASN	3.0
2	D	47	GLU	3.0
1	B	404	LYS	2.9
2	C	77	MET	2.9
2	C	72	ASP	2.8
2	D	74	GLU	2.7
2	C	79	ASP	2.6
1	A	217	ARG	2.6
2	C	52	CYS	2.6
2	D	91	GLY	2.6
1	B	476	ASN	2.6
1	A	418	TRP	2.5
1	A	93	TYR	2.4
1	A	253	GLY	2.4
1	A	5	SER	2.4
1	A	323	GLN	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	55	CYS	2.3
1	B	323	GLN	2.3
2	C	91	GLY	2.3
1	B	418	TRP	2.3
1	B	6	PRO	2.2
1	B	93	TYR	2.1
1	B	449	GLN	2.1
2	C	76	ASP	2.1
1	B	250	ARG	2.1
1	A	450	HIS	2.1
1	B	269	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	FES	C	150	4/4	0.84	0.12	99,99,99,100	0
5	FES	D	151	4/4	0.85	0.12	110,110,110,110	0
4	HCD	B	602	29/29	0.96	0.07	9,11,16,17	0
4	HCD	A	602	29/29	0.97	0.05	9,11,17,17	0
3	HEM	A	601	43/43	0.98	0.06	13,14,16,17	0
3	HEM	B	601	43/43	0.98	0.06	12,14,17,18	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around HCD B 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around HCD A 602:**


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM A 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.