



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 14, 2026 – 11:04 AM UTC

PDB ID : 6NA4 / pdb_00006na4
Title : Co crystal structure of ECR with Butryl-CoA
Authors : DeMirici, H.
Deposited on : 2018-12-05
Resolution : 1.72 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.72 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1039 (1.72-1.72)
Clashscore	190562	1049 (1.72-1.72)
Ramachandran outliers	187476	1041 (1.72-1.72)
Sidechain outliers	187428	1041 (1.72-1.72)
RSRZ outliers	180081	1039 (1.72-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	448	91% 8%
1	B	448	94% 5%
1	C	448	2% 92% 7%
1	D	448	2% 90% 8%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAP	A	501	X	-	-	-
2	NAP	B	501	X	-	-	-
2	NAP	D	501	X	-	-	-
3	VES	C	502	-	-	X	-
3	VES	D	504	-	-	X	-
4	2EC	A	503	-	X	-	-
6	NDP	C	503	X	-	-	-

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 16338 atoms, of which 25 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Putative crotonyl-CoA reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	446	Total	C	N	O	S	0	10	0
			3525	2212	636	663	14			
1	B	448	Total	C	N	O	S	7	10	0
			3546	2225	639	668	14			
1	C	447	Total	C	N	O	S	0	9	0
			3519	2209	638	658	14			
1	D	445	Total	C	N	O	S	0	6	0
			3477	2186	625	652	14			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLU	-	expression tag	UNP E4N096
A	-2	GLY	-	expression tag	UNP E4N096
A	-1	ARG	-	expression tag	UNP E4N096
A	0	HIS	-	expression tag	UNP E4N096
B	-3	GLU	-	expression tag	UNP E4N096
B	-2	GLY	-	expression tag	UNP E4N096
B	-1	ARG	-	expression tag	UNP E4N096
B	0	HIS	-	expression tag	UNP E4N096
C	-3	GLU	-	expression tag	UNP E4N096
C	-2	GLY	-	expression tag	UNP E4N096
C	-1	ARG	-	expression tag	UNP E4N096
C	0	HIS	-	expression tag	UNP E4N096
D	-3	GLU	-	expression tag	UNP E4N096
D	-2	GLY	-	expression tag	UNP E4N096
D	-1	ARG	-	expression tag	UNP E4N096
D	0	HIS	-	expression tag	UNP E4N096

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (CCD ID: NAP) (formula: C₂₁H₂₈N₇O₁₇P₃).



-
- A chemical structure diagram of a five-membered ring. The ring is represented by a black pentagon. The vertices are labeled with green text: C1 (top right), C2 (bottom right), C3 (bottom left), C4 (top left), and N5 (top center). A double bond is shown between the top vertex (N5) and the top-right vertex (C1). The double bond is represented by two parallel blue lines. The label N5 is positioned above the top vertex, and the label C1 is positioned to the right of the top-right vertex. The label C2 is positioned to the right of the bottom-right vertex, C3 is positioned to the left of the bottom-left vertex, and C4 is positioned to the left of the top-left vertex. The label VES is positioned above the N5 label.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	N	0	0
			5	4	1		



Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	N	0	0
			5	4	1		
3	C	1	Total	C	N	0	0
			5	4	1		
3	D	1	Total	C	N	0	0
			5	4	1		
3	D	1	Total	C	N	0	0
			5	4	1		

- Molecule 4 is 9-ETHYL-9H-PURIN-6-YLAMINE (CCD ID: 2EC) (formula: $C_7H_9N_5$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	N	0	0
			11	6	5		
4	B	1	Total	C	N	0	0
			12	7	5		

- Molecule 5 is {S}-[2-[3-[(2 {R})-4-[[[(2 {S}),3 {R}),4 {S}),5 {R}]-5-(6-aminopurin-9-yl)-4-oxidanyl-3-phosphonooxy-oxolan-2-yl]methoxy-oxidanyl-phosphoryl]oxy-oxidanyl-phosphoryl]oxy-3,3-dimethyl-2-oxidanyl-butanoyl]amino]propanoylamino]ethyl] butanethioate (CCD ID: YAS) (formula: $C_{25}H_{42}N_7O_{17}P_3S$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
5	C	1	Total	C	N	O	P	S	1	0
			53	25	7	17	3	1		
5	D	1	Total	C	N	O	P	S	1	0
			53	25	7	17	3	1		

- Molecule 6 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (CCD ID: NDP) (formula: C₂₁H₃₀N₇O₁₇P₃).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
6	C	1	Total	C	H	N	O	P	0	0
			73	21	25	7	17	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	444	Total 454	O 454	0	6
7	B	496	Total 500	O 500	0	4
7	C	484	Total 491	O 491	0	6
7	D	448	Total 455	O 455	0	7


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Putative crotonyl-CoA reductase




- Molecule 1: Putative crotonyl-CoA reductase



- Molecule 1: Putative crotonyl-CoA reductase



- Molecule 1: Putative crotonyl-CoA reductase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	109.34Å 78.83Å 138.79Å 90.00° 108.06° 90.00°	Depositor
Resolution (Å)	39.41 – 1.72 39.41 – 1.72	Depositor EDS
% Data completeness (in resolution range)	95.8 (39.41-1.72) 87.9 (39.41-1.72)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.05 (at 1.72Å)	Xtriage
Refinement program	PHENIX 1.11.1 _2575	Depositor
R, R_{free}	0.170 , 0.198 0.170 , 0.198	Depositor DCC
R_{free} test set	1997 reflections (0.84%)	wwPDB-VP
Wilson B-factor (Å ²)	20.2	Xtriage
Anisotropy	0.280	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 43.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	16338	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 56.81 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.5949e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: NAP, VES, YAS, 2EC, NDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.29	0/3607	0.49	0/4887
1	B	0.32	1/3628 (0.0%)	0.50	0/4917
1	C	0.28	0/3601	0.48	0/4881
1	D	0.31	1/3559 (0.0%)	0.50	3/4825 (0.1%)
All	All	0.30	2/14395 (0.0%)	0.49	3/19510 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	355	TRP	C-N	-5.58	1.26	1.34
1	B	81	ASN	C-O	-5.02	1.17	1.24

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	78	VAL	CA-C-N	6.19	133.22	122.82
1	D	78	VAL	C-N-CA	6.19	133.22	122.82
1	D	78	VAL	O-C-N	-5.30	117.58	123.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3525	0	3448	34	0
1	B	3546	0	3471	24	0
1	C	3519	0	3446	30	0
1	D	3477	0	3406	47	0
2	A	48	0	22	1	0
2	B	48	0	24	7	0
2	D	48	0	23	7	0
3	A	5	0	9	1	0
3	B	5	0	9	0	0
3	C	5	0	9	5	0
3	D	10	0	18	7	0
4	A	11	0	4	0	0
4	B	12	0	9	0	0
5	C	53	0	0	1	0
5	D	53	0	0	3	0
6	C	48	25	24	3	0
7	A	454	0	0	13	0
7	B	500	0	0	4	0
7	C	491	0	0	7	0
7	D	455	0	0	15	0
All	All	16313	25	13922	149	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 149 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:501:NAP:O4D	2:B:501:NAP:C4D	1.66	1.24
2:B:501:NAP:C4B	2:B:501:NAP:O4B	1.65	1.18
5:D:502:YAS:O2A	7:D:601:HOH:O	1.67	1.12
1:A:145:CYS:SG	7:A:1027:HOH:O	2.26	0.94
1:D:395:GLU:OE1	7:D:602:HOH:O	1.89	0.88

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	455/448 (102%)	441 (97%)	11 (2%)	3 (1%)	18	7
1	B	457/448 (102%)	442 (97%)	12 (3%)	3 (1%)	18	7
1	C	454/448 (101%)	439 (97%)	13 (3%)	2 (0%)	30	16
1	D	449/448 (100%)	433 (96%)	14 (3%)	2 (0%)	30	16
All	All	1815/1792 (101%)	1755 (97%)	50 (3%)	10 (1%)	21	9

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	158	ASP
1	A	317	HIS
1	B	158	ASP
1	B	317	HIS
1	C	158	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	369/359 (103%)	368 (100%)	1 (0%)	86	79
1	B	371/359 (103%)	366 (99%)	5 (1%)	61	43
1	C	367/359 (102%)	363 (99%)	4 (1%)	65	50
1	D	363/359 (101%)	358 (99%)	5 (1%)	59	40
All	All	1470/1436 (102%)	1455 (99%)	15 (1%)	68	53

5 of 15 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	355	TRP
1	D	355	TRP
1	C	388	LEU
1	D	388	LEU
1	D	25	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	81	ASN
1	D	157	ASN
1	C	207	ASN
1	D	207	ASN
1	C	407	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAP	D	501	-	50,52,52	4.11	19 (38%)	71,80,80	2.72	23 (32%)
2	NAP	A	501	-	50,52,52	3.92	29 (58%)	71,80,80	2.94	19 (26%)
3	VES	D	504	-	5,5,5	0.40	0	5,5,5	1.60	1 (20%)
4	2EC	A	503	-	12,12,13	1.82	5 (41%)	17,17,18	2.58	8 (47%)
4	2EC	B	503	-	13,13,13	2.00	6 (46%)	18,18,18	2.74	7 (38%)
5	YAS	C	501	-	53,55,55	3.16	20 (37%)	75,81,81	1.90	17 (22%)
6	NDP	C	503	-	51,52,52	7.38	30 (58%)	71,80,80	4.57	27 (38%)
3	VES	B	502	-	5,5,5	0.27	0	5,5,5	1.46	1 (20%)
5	YAS	D	502	-	53,55,55	3.24	27 (50%)	75,81,81	2.47	31 (41%)
3	VES	D	503	-	5,5,5	0.29	0	5,5,5	1.48	1 (20%)
3	VES	A	502	-	5,5,5	0.31	0	5,5,5	1.51	1 (20%)
3	VES	C	502	-	5,5,5	0.25	0	5,5,5	1.47	1 (20%)
2	NAP	B	501	-	50,52,52	3.47	19 (38%)	71,80,80	2.30	21 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	VES	C	502	-	-	-	0/1/1/1
2	NAP	D	501	-	6/6/12/12	7/35/67/67	0/5/5/5
2	NAP	A	501	-	3/3/12/12	4/35/67/67	0/5/5/5
3	VES	D	504	-	-	-	0/1/1/1
4	2EC	A	503	-	-	-	0/2/2/2
4	2EC	B	503	-	-	0/2/2/2	0/2/2/2
6	NDP	C	503	-	2/2/17/17	11/34/77/77	0/5/5/5
3	VES	B	502	-	-	-	0/1/1/1
3	VES	D	503	-	-	-	0/1/1/1
3	VES	A	502	-	-	-	0/1/1/1
5	YAS	D	502	-	-	8/54/70/70	0/3/3/3
5	YAS	C	501	-	-	16/54/70/70	0/3/3/3
2	NAP	B	501	-	1/1/12/12	8/35/67/67	0/5/5/5

The worst 5 of 155 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	C	503	NDP	O4B-C4B	41.78	2.37	1.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	C	503	NDP	C3B-C4B	-13.98	1.17	1.53
5	C	501	YAS	C2B-C3B	-12.89	1.24	1.53
5	D	502	YAS	C2B-C3B	-11.75	1.27	1.53
2	A	501	NAP	O4D-C1D	11.65	1.56	1.40

The worst 5 of 158 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	NAP	C4D-O4D-C1D	-15.69	95.56	109.92
6	C	503	NDP	C4A-N9A-C8A	15.06	121.55	105.74
6	C	503	NDP	C5A-C4A-N9A	-14.99	89.47	105.81
6	C	503	NDP	O4B-C4B-C3B	-13.63	78.10	105.15
2	D	501	NAP	C4D-O4D-C1D	-12.42	98.55	109.92

5 of 12 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	501	NAP	C2D
2	A	501	NAP	C4D
2	A	501	NAP	C3D
2	B	501	NAP	C2B
2	D	501	NAP	C2B

5 of 54 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	501	NAP	O4D-C1D-N1N-C2N
2	B	501	NAP	O4D-C1D-N1N-C6N
2	B	501	NAP	C2D-C1D-N1N-C2N
2	D	501	NAP	C5B-O5B-PA-O3
5	C	501	YAS	C4B-C5B-O5B-P1A

There are no ring outliers.

10 monomers are involved in 34 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	501	NAP	7	0
2	A	501	NAP	1	0
3	D	504	VES	5	0
5	C	501	YAS	1	0
6	C	503	NDP	3	0


Continued on next page...


Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	502	YAS	3	0
3	D	503	VES	2	0
3	A	502	VES	1	0
3	C	502	VES	5	0
2	B	501	NAP	7	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.










Ligand 2EC B 503



Ligand YAS C 501







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	446/448 (99%)	-0.16	1 (0%) 91 93	7, 24, 39, 56	10 (2%)
1	B	448/448 (100%)	-0.29	1 (0%) 91 93	7, 21, 35, 53	10 (2%)
1	C	447/448 (99%)	-0.12	7 (1%) 70 76	8, 23, 40, 59	9 (2%)
1	D	445/448 (99%)	0.01	10 (2%) 62 70	9, 25, 43, 58	6 (1%)
All	All	1786/1792 (99%)	-0.14	19 (1%) 78 82	7, 23, 40, 59	35 (1%)

The worst 5 of 19 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	444	ASN	5.0
1	D	132	VAL	3.7
1	D	134	ALA	3.7
1	C	103	LEU	3.1
1	B	-3[A]	GLU	2.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	2EC	A	503	11/12	0.62	0.17	28,44,51,53	0
3	VES	D	504	5/5	0.64	0.22	44,44,49,50	0
4	2EC	B	503	12/12	0.65	0.18	31,46,54,54	0
3	VES	D	503	5/5	0.81	0.15	32,39,42,42	0
5	YAS	C	501	53/53	0.85	0.12	24,33,58,65	1
3	VES	B	502	5/5	0.86	0.13	29,36,38,42	0
3	VES	A	502	5/5	0.86	0.13	30,33,36,41	0
3	VES	C	502	5/5	0.87	0.14	34,42,43,52	0
5	YAS	D	502	53/53	0.88	0.11	20,28,54,73	1
2	NAP	D	501	48/48	0.94	0.10	22,33,49,61	0
6	NDP	C	503	48/48	0.94	0.10	20,39,59,64	0
2	NAP	A	501	48/48	0.96	0.07	17,21,33,46	0
2	NAP	B	501	48/48	0.97	0.06	15,19,24,32	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 2EC A 503:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 2EC B 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around YAS C 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around YAS D 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NAP D 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around NDP C 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NAP A 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.