



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 08:08 AM UTC

PDB ID : 3NBC / pdb_00003nbc
Title : Clitocybe nebularis ricin B-like lectin (CNL) in complex with lactose, crystallized at pH 4.4
Authors : Renko, M.; Pohleven, J.; Sabotic, J.; Kos, J.; Turk, D.
Deposited on : 2010-06-03
Resolution : 1.01 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.01 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1726 (1.06-0.98)
Clashscore	190562	1007 (1.04-1.00)
Ramachandran outliers	187476	1715 (1.06-0.98)
Sidechain outliers	187428	1716 (1.06-0.98)
RSRZ outliers	180081	1723 (1.06-0.98)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	148	5% 44% 54% .
1	B	148	6% 41% 58% .
2	C	2	100%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2936 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ricin B-like lectin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	148	Total	C	N	O	S	21	7	0
			1156	724	197	234	1			
1	B	148	Total	C	N	O	S	12	9	0
			1161	731	195	234	1			

- Molecule 2 is an oligosaccharide called beta-D-galactopyranose-(1-4)-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	C	2	Total	C	O	0	0	0
			23	12	11			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	349	Total	O	0	0
			349	349		
3	B	247	Total	O	0	0
			247	247		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ricin B-like lectin



- Molecule 1: Ricin B-like lectin



- Molecule 2: beta-D-galactopyranose-(1-4)-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	54.79Å 55.90Å 112.41Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	27.95 – 1.01 27.95 – 1.01	Depositor EDS
% Data completeness (in resolution range)	98.3 (27.95-1.01) 98.3 (27.95-1.01)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.73 (at 1.01Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.157 , 0.172 0.149 , 0.165	Depositor DCC
R_{free} test set	8906 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	11.2	Xtriage
Anisotropy	0.181	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 43.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.010 for k,h,-l	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	2936	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GAL, BGC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	3.03	85/1195 (7.1%)	1.03	4/1641 (0.2%)
1	B	3.13	100/1211 (8.3%)	1.04	3/1664 (0.2%)
All	All	3.08	185/2406 (7.7%)	1.03	7/3305 (0.2%)

All (185) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	111	LEU	CA-C	10.86	1.65	1.52
1	A	111	LEU	CA-C	8.78	1.63	1.52
1	B	19	ILE	CA-C	8.36	1.61	1.52
1	B	117	LEU	C-N	8.27	1.42	1.34
1	B	13	ALA	N-CA	8.04	1.56	1.46
1	B	5	GLY	N-CA	7.49	1.52	1.45
1	A	129	GLY	N-CA	7.44	1.52	1.45
1	A	68	GLN	CA-C	7.29	1.61	1.53
1	B	35	HIS	CG-ND1	7.29	1.46	1.38
1	B	34	GLY	CA-C	7.23	1.59	1.52
1	A	33	ILE	C-N	7.11	1.38	1.33
1	B	117	LEU	CA-C	7.10	1.61	1.52
1	A	101	SER	CA-C	7.05	1.62	1.52
1	B	121	ALA	C-O	7.02	1.32	1.23
1	A	19	ILE	CA-C	6.99	1.60	1.52
1	B	126	ILE	CA-CB	6.96	1.62	1.54
1	A	59	ILE	CA-C	6.94	1.61	1.52
1	A	35	HIS	CG-ND1	6.89	1.45	1.38
1	A	20	ASP	CA-C	6.88	1.60	1.52
1	B	37	LEU	CA-C	6.75	1.61	1.52
1	B	48	GLN	N-CA	6.71	1.54	1.46
1	B	20	ASP	CA-C	6.68	1.60	1.52
1	B	54	LEU	C-N	6.62	1.42	1.33
1	B	84	LEU	CA-C	6.61	1.61	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	84	LEU	CA-C	6.60	1.61	1.52
1	A	24	SER	CA-C	6.60	1.62	1.53
1	A	13	ALA	N-CA	6.60	1.54	1.46
1	B	138	GLN	N-CA	6.60	1.53	1.46
1	A	3	THR	CA-CB	6.58	1.63	1.53
1	B	134	THR	N-CA	6.58	1.54	1.46
1	B	79	VAL	N-CA	6.43	1.53	1.46
1	B	71	VAL	CA-CB	6.39	1.61	1.54
1	A	34	GLY	C-O	6.39	1.30	1.23
1	A	73	VAL	CA-CB	6.32	1.62	1.54
1	B	76	ASP	CA-C	6.29	1.61	1.53
1	B	15	THR	CA-C	6.28	1.61	1.52
1	A	85	VAL	C-N	6.28	1.39	1.33
1	A	31	LEU	C-O	6.28	1.31	1.23
1	B	4	PRO	N-CA	6.27	1.54	1.47
1	A	139	LEU	CA-C	6.25	1.60	1.52
1	A	61	THR	CA-C	6.24	1.60	1.52
1	B	59	ILE	CA-C	6.22	1.60	1.52
1	A	65	VAL	CA-C	6.22	1.60	1.52
1	B	78	LEU	N-CA	6.20	1.54	1.46
1	B	104	PHE	CA-C	6.19	1.60	1.52
1	A	1	SER	C-N	6.18	1.41	1.33
1	A	2	ILE	CA-CB	6.18	1.61	1.54
1	A	71	VAL	CA-CB	6.17	1.61	1.54
1	B	100	ASN	C-O	6.15	1.31	1.23
1	B	108	ILE	CA-CB	6.12	1.62	1.54
1	B	12	VAL	CA-C	6.12	1.61	1.52
1	A	132	ASP	N-CA	6.08	1.54	1.46
1	A	131	VAL	CA-CB	6.07	1.61	1.54
1	A	123	SER	CA-C	6.05	1.60	1.52
1	A	109	PRO	CA-CB	6.04	1.61	1.53
1	B	141	ALA	C-N	6.04	1.41	1.33
1	A	28	GLU	N-CA	6.00	1.53	1.46
1	A	121	ALA	CA-C	5.99	1.60	1.52
1	B	92	PRO	N-CA	5.99	1.54	1.47
1	A	12	VAL	N-CA	5.98	1.53	1.46
1	A	90	PRO	CA-C	5.97	1.59	1.52
1	B	99	GLY	C-O	5.96	1.31	1.23
1	B	104	PHE	N-CA	5.96	1.53	1.45
1	B	125	PRO	N-CD	5.96	1.56	1.47
1	B	93	VAL	C-O	5.95	1.30	1.24
1	B	63	GLN	CA-C	5.92	1.59	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	88	GLN	C-O	5.89	1.31	1.24
1	A	21	LEU	CA-C	5.85	1.60	1.52
1	A	11	ASN	CG-OD1	5.84	1.34	1.23
1	A	82	ALA	CA-C	5.83	1.60	1.52
1	B	60	TYR	N-CA	5.83	1.53	1.45
1	A	46	ASN	CA-C	5.83	1.60	1.52
1	B	83	ALA	CA-C	5.81	1.60	1.52
1	A	54	LEU	CA-C	5.81	1.59	1.52
1	B	134	THR	CA-C	5.79	1.60	1.52
1	B	131	VAL	CA-C	5.78	1.60	1.52
1	B	140	TRP	CA-C	5.76	1.59	1.52
1	B	62	MET	CA-C	5.76	1.59	1.52
1	A	71	VAL	CA-C	5.76	1.58	1.52
1	B	88	GLN	N-CA	5.75	1.53	1.46
1	B	22	THR	CA-CB	5.75	1.62	1.53
1	B	124	THR	CA-C	5.74	1.59	1.53
1	A	114	ALA	C-O	5.73	1.30	1.23
1	B	30	THR	CA-C	5.72	1.60	1.52
1	B	114	ALA	C-O	5.72	1.30	1.23
1	A	79	VAL	C-O	5.71	1.29	1.23
1	A	109	PRO	CA-C	5.69	1.59	1.52
1	A	55	PRO	CA-C	5.68	1.59	1.52
1	B	139	LEU	N-CA	5.67	1.52	1.45
1	A	105	ARG	CZ-NH2	5.64	1.40	1.33
1	A	88	GLN	N-CA	5.64	1.53	1.46
1	A	70	TYR	N-CA	5.64	1.53	1.45
1	B	95	ILE	C-O	5.62	1.30	1.24
1	B	96	GLU	CA-C	5.60	1.59	1.52
1	B	129	GLY	C-O	5.60	1.31	1.23
1	B	129	GLY	N-CA	5.60	1.50	1.44
1	B	43	GLY	N-CA	5.59	1.51	1.45
1	A	69	SER	CA-CB	5.57	1.61	1.53
1	B	122	ASN	C-O	5.56	1.30	1.23
1	B	61	THR	CA-C	5.54	1.59	1.52
1	A	27	ALA	C-O	5.51	1.30	1.23
1	A	95	ILE	N-CA	5.51	1.52	1.46
1	B	139	LEU	C-N	5.50	1.41	1.33
1	B	38	ASN	N-CA	5.48	1.52	1.46
1	B	107	LYS	C-O	5.48	1.30	1.23
1	B	4	PRO	C-O	5.46	1.30	1.23
1	B	14	TYR	C-N	5.46	1.41	1.33
1	B	71	VAL	CA-C	5.46	1.58	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	66	ASN	CA-CB	5.44	1.62	1.53
1	A	26	PRO	N-CD	5.42	1.55	1.47
1	B	87	SER	C-O	5.41	1.30	1.23
1	B	10	THR	C-N	5.41	1.40	1.33
1	B	52	VAL	CA-C	5.41	1.59	1.52
1	A	50	THR	CA-C	5.40	1.59	1.52
1	A	21	LEU	C-O	5.39	1.30	1.24
1	B	113	LEU	C-O	5.38	1.30	1.23
1	A	8	ASN	N-CA	5.38	1.52	1.45
1	A	113	LEU	C-O	5.37	1.30	1.23
1	A	108	ILE	C-N	5.36	1.39	1.33
1	B	53	GLN	N-CA	5.35	1.52	1.46
1	B	73	VAL	CA-C	5.35	1.58	1.52
1	A	45	GLY	C-N	5.35	1.41	1.33
1	B	108	ILE	C-N	5.31	1.39	1.33
1	B	42	SER	N-CA	5.31	1.52	1.46
1	B	137	ASN	CA-CB	5.31	1.62	1.53
1	B	80	ASP	C-O	5.31	1.30	1.23
1	A	91	THR	CA-C	5.30	1.58	1.52
1	A	140	TRP	N-CA	5.30	1.52	1.45
1	A	50	THR	N-CA	5.30	1.52	1.46
1	A	124	THR	N-CA	5.30	1.54	1.46
1	A	18	LEU	C-O	5.29	1.30	1.23
1	B	141	ALA	N-CA	5.28	1.52	1.45
1	A	53	GLN	N-CA	5.28	1.52	1.46
1	A	105	ARG	CA-CB	5.28	1.61	1.53
1	B	95	ILE	N-CA	5.27	1.52	1.46
1	A	130	GLU	C-N	5.27	1.40	1.33
1	B	145	VAL	CA-CB	5.26	1.59	1.53
1	B	115	LEU	CA-C	5.25	1.59	1.52
1	B	18	LEU	N-CA	5.25	1.52	1.45
1	A	5	GLY	N-CA	5.25	1.50	1.45
1	A	16	ASN	CG-OD1	5.25	1.33	1.23
1	A	77	ASN	CA-C	5.25	1.59	1.53
1	B	127	VAL	CA-CB	5.24	1.62	1.54
1	A	35	HIS	C-O	5.23	1.30	1.23
1	A	25	ASN	N-CA	5.22	1.53	1.46
1	A	22	THR	CA-C	5.20	1.59	1.52
1	A	69	SER	C-O	5.19	1.30	1.24
1	B	33	ILE	C-N	5.19	1.38	1.33
1	B	5	GLY	CA-C	5.19	1.57	1.52
1	A	141	ALA	CA-C	5.18	1.59	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	25	ASN	C-N	5.18	1.39	1.34
1	A	125	PRO	N-CA	5.18	1.53	1.47
1	A	47	GLN	CA-C	5.17	1.59	1.52
1	B	128	LEU	C-O	5.17	1.30	1.23
1	B	24	SER	CA-C	5.17	1.59	1.53
1	B	25	ASN	N-CA	5.16	1.54	1.46
1	B	32	ILE	CB-CG1	5.16	1.63	1.53
1	B	55	PRO	C-O	5.16	1.30	1.23
1	A	78	LEU	CA-CB	5.16	1.59	1.52
1	B	29	ASN	CA-C	5.15	1.59	1.52
1	B	56	HIS	N-CA	5.15	1.53	1.46
1	A	25	ASN	C-N	5.14	1.39	1.34
1	B	109	PRO	CA-C	5.14	1.58	1.52
1	A	13	ALA	C-O	5.13	1.30	1.24
1	A	142	PHE	CA-C	5.13	1.58	1.52
1	A	91	THR	N-CA	5.13	1.53	1.46
1	B	105	ARG	CZ-NH2	-5.11	1.26	1.33
1	B	85	VAL	CA-C	5.10	1.58	1.52
1	B	94	SER	CA-C	5.10	1.59	1.52
1	B	91	THR	CA-C	5.09	1.58	1.52
1	B	110	ASN	CA-C	5.08	1.59	1.52
1	B	92	PRO	CA-CB	5.08	1.60	1.53
1	B	75	ASP	N-CA	5.08	1.53	1.46
1	B	106	ILE	C-O	5.07	1.30	1.23
1	A	103	GLN	N-CA	5.07	1.51	1.45
1	B	8	ASN	N-CA	5.06	1.52	1.45
1	A	127	VAL	N-CA	5.06	1.52	1.46
1	B	12	VAL	N-CA	5.06	1.52	1.46
1	A	86	GLY	CA-C	5.05	1.57	1.52
1	A	63	GLN	N-CA	5.03	1.52	1.46
1	A	134	THR	C-O	5.02	1.30	1.24
1	A	70	TYR	C-N	5.01	1.39	1.33
1	A	136	THR	N-CA	5.01	1.52	1.46
1	B	36	HIS	CA-C	5.01	1.58	1.52
1	B	89	GLN	CA-C	5.01	1.58	1.52

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	100	ASN	N-CA-CB	7.36	120.94	109.69
1	A	100	ASN	N-CA-C	-6.88	99.85	110.10
1	B	20	ASP	CA-CB-CG	6.54	119.14	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	99	GLY	N-CA-C	6.11	121.24	113.24
1	A	99	GLY	CA-C-N	-5.28	113.38	120.82
1	A	99	GLY	C-N-CA	-5.28	113.38	120.82
1	B	88	GLN	CB-CG-CD	5.01	121.11	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1156	0	1148	10	0
1	B	1161	0	1173	3	0
2	C	23	0	21	0	0
3	A	349	0	0	7	0
3	B	247	0	0	2	0
All	All	2936	0	2342	13	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (13) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:36[A]:HIS:HD2	1:A:122:ASN:HD21	1.04	0.91
1:A:36[A]:HIS:CD2	1:A:122:ASN:HD21	1.93	0.84
1:A:89:GLN:HG3	3:A:542:HOH:O	1.78	0.83
1:B:68:GLN:HG3	3:B:554:HOH:O	1.78	0.83
1:A:89:GLN:CG	3:A:542:HOH:O	2.38	0.65
1:A:11:ASN:HD22	1:A:14:TYR:H	1.47	0.61
1:A:107[B]:LYS:HG2	3:A:486:HOH:O	2.04	0.58
1:A:89:GLN:CD	3:A:542:HOH:O	2.50	0.54
1:A:107[B]:LYS:HD2	3:A:518:HOH:O	2.14	0.47
1:B:116[B]:THR:HG21	3:B:441:HOH:O	2.16	0.45
1:A:107[B]:LYS:CD	3:A:486:HOH:O	2.67	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:2[A]:ILE:HD11	1:B:104:PHE:CE2	2.55	0.42
1:A:107[B]:LYS:CG	3:A:486:HOH:O	2.66	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	153/148 (103%)	148 (97%)	5 (3%)	0	100	100
1	B	154/148 (104%)	152 (99%)	2 (1%)	0	100	100
All	All	307/296 (104%)	300 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	134/127 (106%)	130 (97%)	4 (3%)	36	6
1	B	136/127 (107%)	135 (99%)	1 (1%)	76	50
All	All	270/254 (106%)	265 (98%)	5 (2%)	48	13

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	ASN
1	A	100	ASN
1	A	146	SER
1	A	148	VAL
1	B	100	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	11	ASN
1	A	16	ASN
1	A	48	GLN
1	B	68	GLN
1	B	88	GLN
1	B	103	GLN
1	B	137	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	BGC	C	1	2	12,12,12	1.48	2 (16%)	17,17,17	1.86	3 (17%)
2	GAL	C	2	2	11,11,12	2.24	4 (36%)	15,15,17	0.74	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	C	1	2	-	0/2/22/22	0/1/1/1
2	GAL	C	2	2	-	0/2/19/22	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	2	GAL	O5-C1	4.71	1.51	1.43
2	C	2	GAL	C2-C3	3.64	1.58	1.52
2	C	1	BGC	C4-C3	3.01	1.60	1.52
2	C	1	BGC	C4-C5	2.86	1.59	1.53
2	C	2	GAL	C4-C3	2.67	1.59	1.52
2	C	2	GAL	C4-C5	2.18	1.57	1.53

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1	BGC	O1-C1-O5	4.32	123.23	110.41
2	C	1	BGC	O5-C1-C2	4.05	117.43	110.30
2	C	1	BGC	O1-C1-C2	2.62	116.59	108.98


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	146/148 (98%)	0.14	7 (4%) 35 43	4, 10, 17, 31	9 (6%)
1	B	147/148 (99%)	0.28	9 (6%) 27 31	4, 11, 18, 30	10 (6%)
All	All	293/296 (98%)	0.21	16 (5%) 30 36	4, 11, 18, 31	19 (6%)

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	148	VAL	29.9
1	A	148	VAL	16.0
1	B	99	GLY	8.3
1	A	146	SER	7.8
1	A	44	TYR	7.4
1	A	99	GLY	4.8
1	B	101	SER	4.5
1	B	98	ALA	4.2
1	B	100	ASN	4.2
1	B	134	THR	4.1
1	A	101	SER	4.0
1	A	98	ALA	3.1
1	B	1[A]	SER	3.0
1	B	146	SER	2.5
1	B	133	GLU	2.1
1	A	133	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	BGC	C	1	12/12	0.94	0.09	11,16,20,27	12
2	GAL	C	2	11/12	0.96	0.07	8,10,12,13	11

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.